

Scientific note

Invasive centipede (*Scolopendra subspinipes*) preying on a house gecko (*Hemidactylus mercatorius*) in eastern Madagascar

(Arthropoda, Scolopendridae and Squamata, Gekkonidae)

David Prötzel, Lalaina O. Randriamanana & Frank Glaw




Fig. 1. Centipede (*Scolopendra subspinipes*) preying on a house gecko (*Hemidactylus mercatorius*). **A.** Clinging the tail and the hindlimbs of the gecko with the pairs of legs; **B.** Large parts of the gecko are already consumed.

Centipedes are predatory arthropods that use their venomous forcipules (a pair of modified legs) for painful defensive bites and also to kill their prey. The usual prey of centipedes consists of insects and arachnids, however large species are able to overpower small vertebrates including amphibians, birds, small mammals, and reptiles (e. g. Noronha et al. 2015, Halpin et al. 2021).

Scolopendra subspinipes Leach, 1815 is a cosmopolitan and variable species (Kronmüller 2012, Siritwut et al. 2016), which can also be found in anthropogenic habitats including houses. The nocturnal house gecko *Hemidactylus mercatorius* Gray, 1842 is often found in human settlements and is probably native, but not endemic to Madagascar (Vences et al. 2004).

Returning from a herpetological nightwalk we found a large invasive centipede (*Scolopendra subspinipes*) on the wooden fence of the terrace of our bungalow of the hotel Nirvana in Ambila-Lemaitso on the east coast of Madagascar (18°52'46.6" S, 49°08'25.3" E, 14 m a. s. l.) on 3 January 2016. With its forcipules and the first 10 pairs of legs it had gripped a *Hemidactylus mercatorius* (field number FGZC 5164), which was apparently freshly dead at the moment of our encounter at 10:57 pm (Fig. 1A). The centipede held the gecko upside down and was nibbling with its mandibles at the tail, which surprisingly was not autotomized during the attack. The lower jaw and the

front legs were already gone and most of the tissue of the body of the gecko was gnawed to the ribs (Fig. 1B). At 11:06 pm the centipede had consumed half of the muscle tissue of the tail leaving just the vertebrae. The remains of the gecko and the scolopender were preserved and stored in the Mention Zoologie et Biodiversité Animale, Antananarivo, Madagascar (UADBA). Considering another observation from the Comoro Islands on *S. subspinipes* eating the Malagasy day gecko *Phelsuma laticauda* (Hawllitschek et al. 2020) we suggest that this invasive centipede species might be a potential predator of other Malagasy reptiles as well.

References

- Halpin, L. R., Terrington, D. I., Jones, H. P., Mott, R., Wong, W. W., Dow, D. C., Carlile, N & Clarke, R. H. 2021. Arthropod predation of vertebrates structures trophic dynamics in island ecosystems. *The American Naturalist* 198: 540–550.
- Hawllitschek, O., Eudeline, R. & Rouillé, A. 2020. Indian Ocean Field Guide: Terrestrial fauna of the Comoros Archipelago – Guide de Terrain de l’Océan Indien: Faune terrestre de l’archipel des Comores. 338pp., Hamburg (self published).
- Kronmüller, C. 2012. Review of the subspecies of *Scolopendra subspinipes* Leach, 1815 with the new description of the South Chinese member of the genus *Scolopendra* Linnaeus, 1758 named *Scolopendra hainanum* spec. nov. *Spixiana* 35: 19–27.
- Noronha, J. d. C. d., Battirolo, L. D., Chagas Junior, A., Miranda, R. M. d., Carpanedo, R. d. S. & Rodrigues, D. d. J. 2015. Predation of bat (*Molossus molossus*: Molossidae) by the centipede *Scolopendra viridicornis* (Scolopendridae) in Southern Amazonia. *Acta Amazonica* 45: 333–336.
- Siritwut, W., Edgecombe, G. D., Sutcharit, C., Tongkerd, P. & Panha, S. 2016. A taxonomic review of the centipede genus *Scolopendra* Linnaeus, 1758 (Scolopendromorpha, Scolopendridae) in mainland Southeast Asia, with description of a new species from Laos. *ZooKeys* 590: 1–124.
- Vences, M., Wanke, S., Vieites, D. R., Branch, W. R., Glaw, F. & Meyer A. 2004. Natural colonization or introduction? Phylogeographical relationships and morphological differentiation of house geckos (*Hemidactylus*) from Madagascar. *Biological Journal of the Linnean Society* 83: 115–130.

David Prötzel, BIOTOPIA – Naturkundemuseum Bayern, Botanisches Institut, Menzinger Str. 67, 80638 München, Germany
Lalaina O. Randriamanana, Mention Zoologie et Biodiversité Animale, Université d’Antananarivo, BP 906, Antananarivo 101, Madagascar

Frank Glaw (corresponding author), Zoologische Staatssammlung München (ZSM–SNSB), Münchhausenstr. 21, 81247 München, Germany; e-mail: glaw@snsb.de

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Spixiana, Zeitschrift für Zoologie](#)

Jahr/Year: 2022

Band/Volume: [045](#)

Autor(en)/Author(s): Prötzel David, Randriamanana Lalaina O., Glaw Frank

Artikel/Article: [Invasive centipede \(*Scolopendra subspinipes*\) preying on a house gecko \(*Hemidactylus mercatorius*\) in eastern Madagascar \(Arthropoda, Scolopendridae and Squamata, Gekkonidae\) 276](#)