

BIBLIOGRAPHY

Univ.-Prof. Mag. Dr. Konrad FIEDLER

(as of: 21.12.2016)

- [1] Fiedler, K. (1985): Der erste Nachweis von *Alcis maculata bastelbergi* Hirschke 1908 im Spessart. – Mitt. int. entomol. Ver. Frankfurt 10: 1-3.
- [2] Fiedler, K. (1985): Auch in Hessen: Die neuerkannte *Mesapamea secalella* Remm, 1983 (Lep., Noctuidae). – Nachr. ent. Ver. Apollo Frankfurt, N. F. 6: 49-52.
- [3] Fiedler, K. (1985): Ein bemerkenswerter Nachweis des Flechtenbären *Eilema lutarella* Linné (Lepidoptera: Arctiidae, Lithosiinae) im Nordspessart. – Hess. faun. Briefe 5 (4): 69-71.
- [4] Fiedler, K. (1987): *Tetramorium caespitum* (Linnaeus, 1758) (Hymenoptera: Formicidae), eine weitere mit *Polyommatus coridon* (Poda, 1761) (Lepidoptera: Lycaenidae) vergesellschaftete Ameisenart. – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 8 (2): 60.
- [5] Fiedler, K. (1988): Die Beziehungen von Bläulingspuppen (Lepidoptera: Lycaenidae) zu Ameisen (Hymenoptera: Formicidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 9: 33-58.
- [6] Fiedler, K. (1989): Differences in the behaviour of ants towards two larval instars of *Lycaena tityrus* (Lepidoptera: Lycaenidae). – Dtsch. entomol. Z., N. F. 36 (4/5): 267-271. doi: 10.1002/mmnd.19890360406.
- [7] Fiedler, K., & MASCHWITZ, U. (1989): Functional analysis of the myrmecophilous relationships between ants (Hymenoptera: Formicidae) and lycaenids (Lepidoptera: Lycaenidae). I. Release of food recruitment in ants by lycaenid larvae and pupae. – Ethology 80: 71-80.
- [8] Fiedler, K., & MASCHWITZ, U. (1988): ... II. Lycaenid larvae as trophobiotic partners of ants – a quantitative approach. – Oecologia 75: 204-206.
- [9] Fiedler, K., & MASCHWITZ, U. (1987 [1988]): ... III. New aspects of the function of the retractile tentacular organs of lycaenid larvae. – Zool. Beitr., N. F. 31 (3): 409-416.
- [10] Fiedler, K., & NÄSSIG, W.A. (1985): *Adscita* (= *Procris*) *statives* L. und *heuseri* Reichl – zwei getrennte Arten? (Lep., Zygaenidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 6 (4): 161-179.
- [11] Maschwitz, U., & FIEDLER, K. (1988): Koexistenz, Symbiose, Parasitismus: Erfolgsstrategien der Bläulinge. – Spektr. Wissensch. 1988 (5): 56-66.
- [12] Nässig, W. A., & FIEDLER, K. (1986): Bemerkungen zum Artenkomplex *Horisme tersata* Den. & Schiff./*radicaria* de la Harpe im Nordwesten der Stadt Frankfurt am Main (Lepidoptera: Geometridae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 7 (1): 29-38.
- [13] Fiedler, K. (1988): The preimaginal epidermal organs of *Lycaena tityrus* (Poda, 1761) and *Polyommatus coridon* (Poda, 1761) (Lepidoptera: Lycaenidae) – a comparison. – Nota lepid. 11 (2): 100-116.
- [14] Maschwitz, U., Nässig, W. A., Dumpert, K., & Fiedler, K. (1988): Larval carnivory and myrmecoxeny, and imaginal myrmecophily in miletine lycaenids (Lepidoptera: Lycaenidae) on the Malay Peninsula. – Tyô to Ga 39 (3): 167-181.
- [15] Fiedler, K. (1989): Rüsseltrillern: eine neue Form taktiler Kommunikation zwischen Bläulingen und Ameisen. – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 10 (2): 125-132.
- [16] Fiedler, K., & MASCHWITZ, U. (1989): The symbiosis between the weaver ant *Oecophylla smaragdina* and *Anthene emolus*, an obligate myrmecophilous lycaenid butterfly. – J. nat. Hist. 23: 833-846. doi: 10.1080/00222938900770441.
- [17] Fiedler, K. (1989): Die Zucht von *Lycaena phlaeas* und *L. tityrus* (Lep., Lycaenidae) auf verschiedenen Polygonaceen. – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 10 (4): 299-300.
- [18] Fiedler, K. (1990): New information on the biology of *Maculinea nausithous* and *M. teleius* (Lepidoptera: Lycaenidae). – Nota lepid. 12 (4): 246-256.
- [19] Fiedler, K., & MASCHWITZ, U. (1989): Adult myrmecophily in butterflies: the role of the ant *Anoplolepis longipes* in the feeding and oviposition behaviour of *Allotinus unicolor*. – Tyô to Ga 40 (4): 241-251.
- [20] Fiedler, K. (1990): Beobachtungen bei der Zucht von *Polyommatus icarus* Rott. (Lepidoptera: Lycaenidae) auf verschiedenen Futterpflanzen. – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 11 (1): 1-9.

- [21] Fiedler, K. (1990): Effects of larval diet on myrmecophilous qualities of *Polyommatus icarus* caterpillars (Lepidoptera: Lycaenidae). – *Oecologia* 83: 284-287.
- [22] Fiedler, K. (1990): Bemerkungen zur Larvalbiologie von *Callophrys rubi* L. (Lepidoptera: Lycaenidae). – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 11 (3): 121-141.
- [23] Fiedler, K., & ROSCISZEWSKI, K. (1990): *Myrmica sabuleti* (Hym., Formicidae) – ein weiterer Symbiosepartner von *Polyommatus coridon* (Lep., Lycaenidae). – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 11 (3): 141-142.
- [24] Fiedler, K. (1990): Untersuchungen zur Funktion, Ökologie und Evolution der Myrmekophilie der Bläulinge (Lepidoptera: Lycaenidae). – Dissertation Univ. Frankfurt, iv + 219 S.
- [25] Fiedler, K., & MASCHWITZ, U. (1990): Quantitative Analyse myrmekophiler Parameter in Assoziationen von Bläulingsraupen (Lepidoptera: Lycaenidae) und Ameisen. – *Verh. dtsh. zool. Ges.* 83: 614-615.
- [26] Fiedler, K. (1991): European and North West African Lycaenidae and their associations with ants. – *J. Res. Lepid.* 28 (4): 239-257.
- [27] Fiedler, K. (1991): Systematic, evolutionary, and ecological implications of myrmecophily within the Lycaenidae (Insecta: Lepidoptera: Papilionoidea). – *Bonner zool. Monogr.* 31: 1-210.
- [28] Fiedler, K. (1991): Zoogeographische Aspekte der myrmekophilen Beziehungen der Bläulinge (Lepidoptera: Lycaenidae). – *Verh. dtsh. zool. Ges.* 84: 502.
- [29] Fiedler, K., & NÄSSIG, W.A. (1991): Nochmals zum Thema "*Adscita stives* und *A. heuseri* – zwei getrennte Arten?" (Lep., Zygaenidae, Procridinae). – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 12 (2): 137-138.
- [30] Schurian, K.G., & Fiedler, K. (1991): Einfache Methoden zur Schallwahrnehmung bei Bläulingslarven (Lepidoptera, Lycaenidae). – *Entomol. Z.* 101 (21): 393-398.
- [31] Fiedler, K. (1991): Weitere Beobachtungen bei Fütterungsversuchen mit Feuerfalterraupen (Lep., Lycaenidae). – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 12 (3): 214-216.
- [32] Fiedler, K., Schurian, K. G., & Seufert, P. (1992): Neue Beobachtungen zu Ameisenassoziationen europäischer Bläulingsraupen (Lepidoptera: Lycaenidae). – *Mitt. intern. entomol. Ver. Frankfurt*. 17 (3): 121-130.
- [33] Balint, Z.S., & Fiedler, K. (1992): *Plebeius sephirus* (Frivaldszky, 1835) in Pannonia, with special reference to its status and ecology in Hungary. – *Oedipus* 4: 1-24.
- [34] Fiedler, K., & Hölldobler, B. (1992): Ants and *Polyommatus icarus* immatures (Lycaenidae) – sex-related developmental benefits and costs of ant-attendance. – *Oecologia* 91: 468-473.
- [35] Fiedler, K. (1992): Notes on the biology of *Hypolycaena othona* (Lepidoptera: Lycaenidae) in West Malaysia. – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 13 (2): 65-92.
- [36] Fiedler, K. (1992): The life-history of *Surendra florimel* DOHERTY 1889 (Lepidoptera: Lycaenidae) in West Malaysia. – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 13 (2): 107-135.
- [37] Fiedler, K. (1992): *Rhodometra sacraria* L. (Lepidoptera, Geometridae) in Mainfranken beobachtet. – *Nachr. entomol. Ver. Apollo Frankfurt*, N. F. 13 (2a): 212.
- [38] Tautz, J., & Fiedler, K. (1992): Mechanoreceptive properties of caterpillar hairs involved in mediation of butterfly-ant symbioses. – *Naturwissenschaften* 79: 561-563.
- [39] Fiedler, K. (1992): Recent contributions to the behavioural ecology and evolution of lycaenid-ant interactions. – *Nota lepid.* 14, Suppl. 4: 11-13.
- [40] Schurian, K.G., Fiedler, K., & Maschwitz, U. (1993): Parasitoids exploit myrmecophilous secretions of lycaenid butterfly caterpillars. – *J. Lepid. Soc.* 47 (2): 150-154.
- [41] Fiedler, K., Krug, E., & Proksch, P. (1993): Complete elimination of host-plant quinolizidine alkaloids by larvae of a polyphagous lycaenid butterfly, *Callophrys rubi*. – *Oecologia* 94: 441-445. doi: 10.1007/BF00317121.
- [42] Fiedler, K. (1993): The remarkable life-history of two Malaysian lycaenid butterflies. – *Nature Malaysiana* 18 (2): 35-43.
- [43] Mattoni, R.H.T., & Fiedler, K. (1993): On the functional foreleg tarsus in *Caerulea* males (Lepidoptera: Lycaenidae: Polyommataini). – *J. Res. Lepid.* 30: 289-296.

- [44] Fiedler, K. (1994): Observations on the biology of *Eooxylides tharis* (Lepidoptera: Lycaenidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 14: 325-337.
- [45] Fiedler, K. (1994): The life-history of *Caleta roxus* (Lepidoptera: Lycaenidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 14: 371-384.
- [46] Schurian, K. G., & Fiedler, K. (1994): Zur Biologie von *Polyommatus (Lysandra) dezinus* (de Freina & Witt) (Lepidoptera: Lycaenidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 14: 339-353.
- [47] Fiedler, K. (1994): Lycaenid butterflies and plants: is myrmecophily associated with amplified hostplant diversity? – Ecol. Entomol. 19: 79-82. doi: 10.1111/j.1365-2311.1994.tb00393.x.
- [48] Fiedler, K., Schurian, K.G., & Hahn, M. (1994): The life-history and myrmecophily of *Polyommatus candalus* (Herrich-Schäffer) from Turkey (Lep., Lycaenidae). – Linneana Belg. 14: 315-332.
- [49] Fiedler, K., & Saam, C. (1994): Does ant-attendance influence development in 5 European Lycaenidae species? – Nota lepid. 17 (1/2): 5-24.
- [50] Fiedler, K., & Schurian, K.G. (1994): Oviposition behaviour in *Lycaena thetis* Klug (Lycaenidae). – Nota lepid. 17 (1/2): 25-29.
- [51] Baumgarten, H.-Th., & Fiedler, K. (1994): Zum Abwehrverhalten der Raupen von *Saturnia (Eudia) pavonia* (Lepidoptera: Saturniidae) gegenüber Ameisen. – Nachr. entomol. Ver. Apollo Frankfurt, N. F. 15 (3): 315-322.
- [52] Wiesen, B., Krug, E., Fiedler, K., Wray, V., & Proksch, P. (1994): Sequestration of hostplant-derived flavonoids by lycaenid butterfly *Polyommatus icarus*. – J. Chem. Ecol. 20: 2523-2538. doi: 10.1007/BF02036189.
- [53] Fiedler, K., Seufert, P., Maschwitz, U., & Azarae, I. (1995): Notes on larval biology and pupal morphology of Malaysian *Curetis* butterflies (Lepidoptera: Lycaenidae). – Tyô to Ga 45 (4): 287-299.
- [54] Fiedler, K., & Saam, C. (1995): A "microbial cost" of butterfly-ant mutualisms (Lycaenidae). – J. Lepid. Soc. 49: 80-84.
- [55] Fiedler, K. (1995): Bläulinge in Südostasien: Diversität, Spezifität und Wirtspflanzenbeziehungen im Überblick. – Verh. dt. zool. Ges. 88.1: 257.
- [56] Burghardt, F., Fiedler, K., & Proksch, P. (1995): Wirtspflanzenabhängige Flavonoidmuster im Bläuling *Polyommatus icarus* (Lycaenidae). – Verh. dt. zool. Ges. 88.1: 256.
- [57] Seufert, P., & Fiedler, K. (1995): Diversität und Strukturierung einer Bläulingsgilde im tropischen Regenwald Malaysias. – Verh. dt. zool. Ges. 88.1: 265.
- [58] Fiedler, K. (1995): Lycaenid butterflies and plants: is myrmecophily associated with particular hostplant preferences? – Ethol. Ecol. Evol. 7(2): 107-132. doi: 0.1080/08927014.1995.9522960.
- [59] Fiedler, K. (1995): Associations of lycaenid butterflies with ants in Turkey. – S. 437-450 in: HESSELBARTH, G., VAN OORSCHOT, H., & Wagener, S.: Die Tagfalter der Türkei unter Berücksichtigung der angrenzenden Länder, Vol. 1. S. Wagener Verlag, Bocholt.
- [60] Fiedler, K., & Seufert, P. (1995): The mature larva and pupa of *Semanga superba* (Lepidoptera: Lycaenidae). – Nachr. entomol. Ver. Apollo, Frankfurt, N. F. 16: 1-12.
- [61] Fiedler, K. (1995): Lycaenid butterflies and plants: Hostplant relationships, tropical versus temperate. – Ecotropica 1: 51-58.
- [62] Fiedler, K. (1995): Lebenszyklen tropischer Bläulinge – von Interaktionen mit Ameisen geprägt. – Rundgespr. Kommiss. Ökol. Bayr. Akad. Wiss. 10: 199-214.
- [63] Fiedler, K., & Saam, C. (1995): Ants benefit from attending facultatively myrmecophilous Lycaenidae caterpillars: evidence from a survival study. – Oecologia 104: 316-322.
- [64] Fiedler, K., Seufert, P., Pierce, N.E., Pearson, J.G., & Baumgarten, T. (1995): Exploitation of lycaenid-ant mutualisms by braconid parasitoids. – J. Res. Lepid. 31 (3/4): 153-168.
- [65] Fiedler, K., & Hagemann, D. (1995): The influence of larval age and ant number on myrmecophilous interactions of the African Grass Blue butterfly, *Zizeeria knysna*. – J. Res. Lepid. 31 (3/4): 213-232.
- [66] Fiedler, K., Hölldobler, B., & Seufert, P. (1996): Butterflies and ants: the communicative domain. – Experientia 52: 14-24. doi: 10.1007/BF01922410.

- [67] Burghardt, F., & Fiedler, K. (1996): The influence of diet on growth and secretion behaviour of myrmecophilous *Polyommatus icarus* caterpillars (Lepidoptera: Lycaenidae). – Ecol. Entomol. 21: 1-8. doi: 10.1111/j.1365-2311.1996.tb00259.x.
- [68] Sanetra, M., & Fiedler, K. (1996): Behaviour and morphology of an aphytophagous lycaenid caterpillar: *Cigaritis (Apharitis) acamas* Klug 1834 (Lepidoptera: Lycaenidae). – Nota lepid. 18 (1): 57-76.
- [69] Hesbacher, S., Giez, I., Embacher, G., Fiedler, K., Max, W., Trawöger, A., Türk, R., Lange, O.L., & Proksch, P. (1995): Sequestration of lichen compounds by lichen-feeding members of the Arctiidae (Lepidoptera). – J. Chem. Ecol. 21: 2079-2089. doi: 10.1007/BF02033864.
- [70] Schurian, K.G., & Fiedler, K. (1996): Adult behaviour and early stages of *Lycaena ochimus* (Herrich-Schäffer [1851]) (Lepidoptera: Lycaenidae). – Nachr. entomol. Ver. Apollo Frankfurt, N. F.16 (4): 329-343.
- [71] Fiedler, K., & Hummel, V. (1996): Myrmecophily in the brown argus butterfly, *Aricia agestis* (Lepidoptera: Lycaenidae): effects of larval age, ant number and persistence of contacts with ants. – Zoology 99: 128-137.
- [72] Seufert, P., & Fiedler, K. (1996): The influence of ants on patterns of colonization and establishment within a set of coexisting lycaenid butterflies in a South-East Asian tropical rain forest. – Oecologia 106: 127-136. doi: 10.1007/BF00334414.
- [73] Fiedler, K. (1996): Interactions between lycaenid butterflies and ants in Peninsular Malaysia. – In: EDWARDS, D.S. et al. (eds.): Tropical rainforest research – current issues, pp. 291-296. Kluwer Acad. Publ., Dordrecht. doi: 10.1007/978-94-009-1685-2_28.
- [74] Fiedler, K. (1996): Host-plant relationships of lycaenid butterflies: large-scale patterns, interactions with plant chemistry, and mutualism with ants. – Entomol. exp. appl. 80: 259-267. doi: 10.1007/978-94-009-1720-0_59.
- [75] Seufert, P., & Fiedler, K. (1996): Life-history diversity and local co-existence of three closely related lycaenid butterflies (Lepidoptera: Lycaenidae) in Malaysian rainforests. – Zool. Anz. 234 (4): 229-239.
- [76] Burghardt, F., & Fiedler, K. (1996): Myrmecophilous behaviours in caterpillars of the butterfly, *Polyommatus icarus* (Rottemburg, 1775): temporal patterns and age dependency (Lepidoptera: Lycaenidae). – Nota lepid. 19 (3/4): 212-229.
- [77] Schulze, C.H., & Fiedler, K. (1996): First record of the family Dudgeoneidae (Lepidoptera, Ditrysia) for Borneo. – Tinea 15 (1): 74-77.
- [78] Horstmann, K., Fiedler, K., & Baumgarten, H.-T. (1997): Zur Taxonomie und Bionomie einiger Ichneumonidae (Hymenoptera) als Parasitoide westpaläarktischer Lycaenidae (Lepidoptera). – Nachrbl. Bayr. Entomol. 46 (1/2): 2-7.
- [79] Schulze, C.H., & Fiedler, K. (1997): Notes on the biology of *Dervas gobrias* (Hewitson, 1864) (Lepidoptera: Pieridae, Coliadinae). – Trans. Lepid. Soc. Japan 48 (1): 25-30.
- [80] Zub, P., Fiedler, K., & Nässig, W.A. (1997): Zur Artenschutz-Problematik bei Insekten. – Natur & Museum 127 (5): 147-152.
- [81] Burghardt, F., Fiedler, K., & Proksch, P. (1997): Uptake of flavonoids from *Vicia villosa* (Fabaceae) by the lycaenid butterfly, *Polyommatus icarus* (Lepidoptera: Lycaenidae). – Biochem. Syst. Ecol. 25 (6): 527-536. doi: 10.1016/S0305-1978(97)00057-4.
- [82] Geuder, M., Wray, V., Fiedler, K., & Proksch, P. (1997): Sequestration and metabolism of host-plant flavonoids by the lycaenid butterfly *Polyommatus bellargus*. – J. Chem. Ecol. 23: 1361-1372. doi: 10.1023/B:JOEC.0000006469.00557.69.
- [83] Fiedler, K. (1997): Life-history patterns of myrmecophilous butterflies and other insects: their implications on tropical species diversity. – Pp. 71-92 in: ULRICH, H. (ed.): Tropical biodiversity and systematics. Proceedings of the International Symposium on Biodiversity & Systematics in Tropical Ecosystems, Museum Alexander Koenig, Bonn.
- [84] Häuser, C.L., Schulze, C.H., & Fiedler, K. (1997): The butterfly species (Insecta: Lepidoptera: Rhopalocera) of Kinabalu Park, Sabah. – Raffles Bull. Zool. 45 (2): 281-304.

- [85] Burghardt, F., Fiedler, K., & Proksch, P. (1997): Flavonoidaufnahme im Bläuling *Polyommatus icarus*: der Einfluß von Nahrungspflanze, Größe und Geschlecht. – Mitt. Dt. Ges. allg. angew. Entomol. 11: 533-536.
- [86] Schulze, C.H., & Fiedler, K. (1997): Patterns of diversity and stratification in hawkmoths (Lepidoptera: Sphingidae) of a Bornean rain forest. – Mitt. Dt. Ges. allg. angew. Entomol. 11: 767-770.
- [87] Fiedler, K. (1998): Lycaenid-ant interactions of the *Maculinea* type: tracing their historical roots in a comparative framework. – J. Insect Conserv. 2: 3-14. doi: 10.1023/A:1009618721239.
- [88] Baumgarten, H.-T., & Fiedler, K. (1998): Parasitoids of lycaenid butterfly caterpillars: different patterns in resource use and their impact on the hosts' symbiosis with ants. – Zool. Anz. 236: 167-180.
- [89] Fiedler, K. (1998): Geographical patterns in life-history traits of lycaenid butterflies – ecological and evolutionary implications. – Zoology (Analysis of Complex Systems) 100: 336-347.
- [90] Fiedler, K. (1998): Diet breadth and host plant diversity of tropical- versus temperate-zone herbivores: South East Asian and West Palaearctic butterflies as a case study. – Ecol. Entomol. 23: 285-297. doi: 10.1046/j.1365-2311.1998.00132.x
- [91] Beck, J., Mühlenberg, E., & Fiedler, K. (1999): Mud-puddling behavior in tropical butterflies: in search of proteins or minerals? – Oecologia 119: 140-148. doi: 10.1007/s004420050770.
- [92] Schittko, U., Burghardt, F., Fiedler, K., Wray, V., & Proksch, P. (1999): Sequestration and distribution of flavonoids in the common blue butterfly *Polyommatus icarus* reared on *Trifolium repens*. – Phytochemistry 51 (5): 609-614. doi: 10.1016/S0031-9422(98)00746-8.
- [93] Süssenbach, D., & Fiedler, K. (1999): Noctuid moths attracted to fruit baits: testing models and methods of estimating species diversity. – Nota lepid. 22 (2): 115-154.
- [94] Fiedler, K. (1998): Bläulinge in Südostasien: Vielfalt der Lebenszyklen und Artendiversität. – Bielefelder ökol. Beitr. 12: 29.
- [95] Knüttel, H., & Fiedler, K. (1999): Flavonoids from larval food plants determine UV wing patterns in *Polyommatus icarus* (Lepidoptera: Lycaenidae). – Zoology 102, Suppl. II (DZG 92.1): 83.
- [96] Schulze, C.H., & Fiedler, K. (1998 [1999]): Habitat preferences and flight activity of Morphinae butterflies in a Bornean rain forest, with a note on sound production by adult *Zeuxidia* (Lepidoptera: Nymphalidae). – Mal. Nat. J. 52 (3/4): 163-176.
- [97] Seufert, P., & FIEDLER, K. (1999): Myrmecophily and parasitoid infestation of South-East Asian lycaenid butterfly larvae. – Ecotropica 5: 59-64.
- [98] Schulze, C.H., & Fiedler, K. (1999): Species richness of South East Asian butterflies – how can it be estimated using faunal lists at different geographic scales. – Pp. 63-70 in: Mohamed, M., & Bernard, H. (eds.): Tropical ecosystem research in Sabah – for whom and for what? Universiti Malaysia, Sabah, Kota Kinabalu.
- [99] Ruf, C., & Fiedler, K. (2000): Thermal gains through collective metabolic heat production in social caterpillars of *Eriogaster lanestris*. – Naturwissenschaften 87: 193-196. doi: 10.1007/s001140050702
- [100] Fischer, K., & Fiedler, K. (2000): On the costs of being a male: case studies on protandrous copper butterflies (Lepidoptera: Lycaenidae). – Zoology 103, Suppl. III (DZG 93.1): 41.
- [101] Ruf, C., & Fiedler, K. (2000): Communal foraging behaviour and communication in social caterpillars of *Eriogaster lanestris* (Lepidoptera: Lasiocampidae). – Zoology 103, Suppl. III (DZG 93.1): 110.
- [102] Fischer, K., & Fiedler, K. (2000): Response of the copper butterfly *Lycaena tityrus* to increased leaf nitrogen in natural food-plants: evidence against the nitrogen limitation hypothesis. – Oecologia 124: 235-241. doi: 10.1007/s004420000365.
- [103] Fischer, K., & Fiedler, K. (2000): Sex-related differences in reaction norms in the butterfly *Lycaena tityrus*. – Oikos 90: 372-380. doi: 10.1034/j.1600-0706.2000.900218.x.
- [104] Burghardt, F., Knüttel, H., Becker, M., & Fiedler, K. (2000): Flavonoid wing pigments increase attractiveness of female common blue (*Polyommatus icarus*) butterflies to mate-searching males. – Naturwissenschaften 87: 304-307. doi: 10.1007/s001140050726.
- [105] Ruf, C., & Fiedler, K. (2000): Trail following as a rare phenomenon among non-social lappet moth larvae (Lasiocampidae). – Entomol. gener. 25: 17-25.

- [106] Süssenbach, D., & Fiedler, K. (2000): Faunistische Ergebnisse einer Köderfangstudie an Eulenfaltern (Noctuidae) im Obermain-Hügelland. – Ber. Naturwiss. Ges. Bayreuth 24: 255-272.
- [107] Stadler, B., Fiedler, K., Kawecki, T., & Weisser, W. (2001): Costs and benefits for phytophagous myrmecophiles: when ants are not always available. – Oikos 92: 467-478. doi: 10.1034/j.1600-0706.2001.920308.x.
- [108] Fischer, K., & Fiedler, K. (2001): Resource based territoriality in the butterfly *Lycaena hippothoe* and environmentally introduced behavioural shifts. – Anim. Behav. 61: 723-732. doi: 10.1006/anbe.2000.1662.
- [109] Fischer, K., & Fiedler, K. (2001): Dimorphic growth patterns and sex-specific reaction norms in the butterfly *Lycaena hippothoe sumadiensis*. – J. Evol. Biol. 14: 210-218. doi: 10.1046/j.1420-9101.2001.00280.x.
- [110] Ruf, C., Costa, J. C., & Fiedler, K. (2001): Trail-based communication in social caterpillars of *Eriogaster lanestris* (Lepidoptera: Lasiocampidae). – J. Insect Behav. 14: 231-245. doi: 10.1023/A:1007841828892.
- [111] Fischer, K., & Fiedler, K. (2001): Egg weight variation in relation to offspring fitness in the butterfly *Lycaena hippothoe*: more small or fewer large eggs? – Popul. Ecol. 43: 105-109. doi: 10.1007/PL00012009.
- [114] Fiedler, K. (2001): Ants that associate with lycaenid butterfly larvae: diversity, ecology and biogeography. – Diversity & Distributions 7: 45-60. doi: 10.1046/j.1472-4642.2001.00096.x.
- [115] Burghardt, F., Proksch, P., & Fiedler, K. (2001): Flavonoid sequestration by the common blue butterfly *Polyommatus icarus*: quantitative intraspecific variation in relation to larval hostplant, sex and body size. – Biochem. Syst. Ecol. 29: 875-889. doi: 10.1016/S0305-1978(01)00036-9.
- [116] Knüttel, H., & Fiedler, K. (2001): Hostplant derived variation in ultraviolet wing patterns influences mate selection of male butterflies. – J. Exp. Biol. 204: 2447-2459.
- [117] Elmes, G.W., Thomas, J.A., Munguira, M.L., Fiedler, K. (2001): Larvae of lycaenid butterflies that parasitise ant colonies provide exceptions to normal insect growth rules. – Biol. J. Linn. Soc. 73: 259-278. doi: 10.1111/j.1095-8312.2001.tb01362.x
- [118] Burghardt, F., Proksch, P., & Fiedler, K. (2001): Loss of flavonoid pigments with ageing in male *Polyommatus icarus* butterflies (Lepidoptera: Lycaenidae). – Nota lepid. 24 (1/2): 77-84.
- [119] Schulze, C.H., Linsenmair, K.E., & Fiedler, K. (2001): Understorey versus canopy – patterns of vertical stratification and diversity among Lepidoptera in a Bornean rainforest. – Plant Ecol. 153: 133-152. doi: 10.1023/A:1017589711553.
- [120] Fischer, K., & Fiedler, K. (2000 [2001]): Methodische Aspekte von Fang-Wiederfangstudien am Beispiel der Feuerfalter *Lycaena belle* und *L. hippothoe*. – Beitr. Ökol. 4 (2): 157-172.
- [121] Fischer, K., & Fiedler, K. (2001): Partial biennialism in alpine *Lycaena hippothoe* (Lycaenidae: Lycaenini)? – Nota lepid. 24 (1/2): 73-76.
- [122] Fischer, K., & Fiedler, K. (2001): Effects of larval starvation on adult life-history traits in the butterfly *Lycaena tityrus*. – Entomol. Gener. 25 (4): 249-254.
- [123] Fischer, K., & Fiedler, K. (2001): Effects of adult feeding and temperature regime on fecundity and longevity in the butterfly *Lycaena hippothoe*. – J. Lepid. Soc. 54: 91-95.
- [124] Fischer, K., & Fiedler, K. (2001): Sexual differences in life-history traits in the butterfly *Lycaena tityrus*: a comparison between diapause and direct development. – Entomol. Exp. Appl. 100: 325-330. doi: 10.1046/j.1570-7458.2001.00879.x.
- [125] Beck, J., Schulze, C.H., Linsenmair, K.E. & Fiedler, K. (2002): From forest to farmland: diversity of geometrid moths along two habitat gradients on Borneo. – J. Trop. Ecol. 18: 33-51. doi: 10.1017/S026646740200202X.
- [126] Ruf, C., Kornmaier, B., & Fiedler, K. (2001): Continuous long-term monitoring of daily foraging patterns in three species of lappet moth caterpillars (Lepidoptera: Lasiocampidae). – Nota lepid. 24 (3): 87-99.
- [127] Fischer, K., & Fiedler, K. (2001): Spatio-temporal dynamics in a population of the copper butterfly *Lycaena hippothoe* (Lycaenidae). – Nota lepid. 24 (3): 77-86.

- [128] Knüttel, H., & Fiedler, K. (2001): On the use of UV photography and UV wing patterns in butterfly morphology and taxonomy. – J. Lepid. Soc. 54: 137-144.
- [129] Fischer, K., & Fiedler, K. (2002): Life-history plasticity in the butterfly *Lycaena hippothoe*: local adaptations and trade-offs. – Biol. J. Linn. Soc. 75: 173-185. doi: 10.1046/j.1095-8312.2002.00014.x
- [130] Ruf, C., Schreier, H.-P., & Fiedler, K. (2002): Zur Entwicklung und Morphologie der Prä-imaginalstadien und Falter von *Bufoidia pittawayi* Wiltshire & Legrain, 1997 (Lepidoptera, Lasiocampidae). – Entomol. Z. 112: 114-119.
- [131] Ruf, C., & Fiedler, K. (2002): Plasticity in foraging patterns of larval colonies of the small egg moth, *Eriogaster lanestris* (Lepidoptera: Lasiocampidae). – Oecologia 131: 626-634. doi: 10.1007/s00442-002-0914-8.
- [132] Freese, A., & Fiedler, K. (2002): Experimental evidence for specific distinctness of the two wood white butterfly taxa, *Leptidea sinapis* and *L. reali* (Lepidoptera: Pieridae). – Nota lepid. 25(1): 39-59.
- [133] Fischer, K., & Fiedler, K. (2002): Reaction norms for age and size at maturity in response to temperature: a test of the compound interest hypothesis. – Evol. Ecol. 16: 333-349. doi: 10.1023/A:1020271600025.
- [134] Blüthgen, N., & Fiedler, K. (2002): Interactions between weaver ants (*Oecophylla smaragdina*), homopterans, trees and lianas in an Australian rainforest canopy. – J. Anim. Ecol. 71: 793-801. doi: 10.1046/j.1365-2656.2002.00647.x.
- [135] Ruf, C., & Fiedler, K. (2002): Tent-based thermoregulation in social caterpillars: behavioural mechanisms and physical features of the tent. – J. therm. Biol. 27: 493-501. doi: 10.1016/S0306-4565(02)00022-0.
- [136] Brehm, G., & Fiedler, K. (2003): Faunal composition of geometrid moths changes with altitude in an Andean montane rainforest. – J. Biogeogr. 30: 431-440. doi: 10.1046/j.1365-2699.2003.00832.x.
- [137] Schulze, C.H., & Fiedler, K. (2003): Vertical and temporal diversity of a species-rich moth taxon in Borneo. – Pp. 69-85 in: Y. BASSET, V. NOVOTNY, S. MILLER & R. KITCHING (eds.): *Arthropods of tropical forests: spatio-temporal dynamics and resource use in the canopy*. Cambridge Univ. Press.
- [138] Stadler, B., Kindlmann, P., Šmilauer, P., & Fiedler, K. (2003): A comparative analysis of morphological and ecological characters of European aphids and lycaenids in relation to ant attendance. – Oecologia 135: 422-430. doi: 10.1007/s00442-003-1193-8.
- [139] Brehm, G., Süßenbach, D., & Fiedler, K. (2003): Unique elevational diversity patterns of geometrid moths in an Andean montane rainforest. – Ecography 26: 356-366. doi: 10.1034/j.1600-0587.2003.03498.x.
- [140] Brehm, G., Homeier, J., & Fiedler, K. (2003): Beta-diversity of geometrid moths in an Andean montane rainforest. – Diversity & Distributions 9: 351-366. doi: 10.1046/j.1472-4642.2003.00023.x.
- [141] Blüthgen, N., Gebauer, G., & Fiedler, K. (2003): Disentangling a rainforest food web using stable isotopes: dietary diversity in a species-rich ant community. – Oecologia 137: 426-435. doi: 10.1007/s00442-003-1347-8.
- [142] Ruf, C., Freese, A., & Fiedler, K. (2003): Larval sociality in three species of central-place foraging lappet moths (Lasiocampidae): a comparative survey. – Zool. Anz. 242: 209-222. doi: 10.1078/0044-5231-00099.
- [143] Schulze, C.H., & Fiedler, K. (2003): Hawkmoth diversity in northern Borneo does not reflect the influence of anthropogenic habitat disturbance. – Ecotropica 9: 99-102.
- [144] Blüthgen, N., & FIEDLER, K. (2004): Preferences for sugars and amino acids and their conditionality in a diverse nectar-feeding ant community. – J. Anim. Ecol. 73: 155-166. doi: 10.1111/j.1365-2656.2004.00789.x.
- [145] Brehm, G., & Fiedler, K. (2004): Bergmann's rule does not apply to geometrid moths along an elevational gradient in an Andean montane rain forest. – Global Ecol. Biogeogr. 13: 7-14. doi: 10.1111/j.1466-882X.2004.00069.x.
- [146] Fiedler, K., & Ruf, C. (2003): *Araschnia levana* larvae (Nymphalidae) do not accept *Humulus lupulus* (Cannabaceae) as food plant. – Nota lepid. 25 (4): 265-266.

- [147] Brehm, G., & Fiedler, K. (2004): Ordinating tropical moth samples from an elevational gradient: a comparison of common methods. – *J. Trop. Ecol.* 20: 165-172. doi: 10.1017/S0266467403001184.
- [148] Axmacher, J.C., Tünte, H., Schrupf, M., Müller-Hohenstein, K., Lyaruu, H.V.M., & Fiedler, K. (2004): Diverging diversity dynamics of vascular plants and moths during forest regeneration on Mt. Kilimanjaro, Tanzania. – *J. Biogeogr.* 31: 895-904. doi: 10.1111/j.1365-2699.2004.00995.x.
- [149] Axmacher, J.C., Holtmann, G., Scheuermann, L., Brehm, G., Müller-Hohenstein, K., & Fiedler, K. (2004): Diversity of geometrid moths (Lepidoptera: Geometridae) along an Afrotropical elevational rainforest transect. – *Diversity & Distributions* 10: 293-302. doi: 10.1111/j.1366-9516.2004.00101.x.
- [150] Blüthgen, N., Stork, N., & Fiedler, K. (2004): Bottom-up control and co-occurrence in complex communities: honeydew and nectar determine a rainforest ant mosaic. – *Oikos* 106: 344-358. doi: 10.1111/j.0030-1299.2004.12687.x.
- [151] Freese, A., & Fiedler, K. (2004): Unterscheidungsmerkmale von *Leptidea sinapis* (Linnaeus, 1758) und *Leptidea reali* (Reissinger, 1989) (Lepidoptera, Pieridae) und ihre Eignung zur Artbestimmung. – *Nachr. entomol. Ver. Apollo Frankfurt, N. F.*: 25: 65-77.
- [152] Blüthgen, N., & FIEDLER, K. (2004): Competition for composition: lessons from nectar-feeding ant communities. – *Ecology* 85: 1479-1485. doi: <http://dx.doi.org/10.1890/03-0430>.
- [153] Blüthgen, N., GOTTSBERGER, G., & FIEDLER, K. (2004): Sugar and amino acid composition of ant-attended nectar and honeydew sources from an Australian rainforest. – *Austral Ecol.* 29: 418-429. doi: 10.1111/j.1442-9993.2004.01380.x.
- [154] Schulze, C.H., & Fiedler, K. (2004): *Orthonama obstipata* Fabricius, 1794 (Geometridae: Larentiinae) – new for the fauna of Borneo. – *Nachr. entomol. Ver. Apollo Frankfurt, N. F.* 25: 153-154.
- [155] Fiedler, K., & Schulze, C. H. (2004): Forest modification affects diversity, but not dynamics of speciose tropical pyraloid moth communities. – *Biotropica* 36: 615-627. doi: 10.1111/j.1744-7429.2004.tb00355.x.
- [156] Axmacher, J.C., & Fiedler, K. (2004): Manual versus automatic moth sampling at equal light sources – a comparison of catches from Mt. Kilimanjaro. – *J. Lepid. Soc.* 58: 196-202.
- [157] Megens, H.-J., de Jong, R., & Fiedler, K. (2005): Phylogenetic patterns in larval host plant and ant association of Indo-Australian Arhopalini butterflies (Lycaenidae: Theclinae). – *Biol. J. Linn. Soc.* 84: 225-242. doi: 10.1111/j.1095-8312.2005.00426.x.
- [158] Fischer, K., Zeilstra, I., Hetz, S. K., & Fiedler, K. (2004): Physiological costs of growing fast: Does accelerated growth reduce pay-off in adult fitness? – *Evol. Ecol.* 18: 343-353. doi: 10.1007/s10682-004-2004-3.
- [159] Brehm, G., & Fiedler, K. (2005): Diversity and community structure of geometrid moths of disturbed habitat in a montane area in the Ecuadorian Andes. – *J. Res. Lepid.* 38: 1-14. <http://lepidopteraresearchfoundation.org/pdf/pdf38/38-001.pdf>
- [160] Ruf, C., & Fiedler, K. (2005): Colony survivorship of social caterpillars in the field: a case study of the small eggar moth (Lep., Lasiocampidae). – *J. Res. Lepid.* 38: 15-25. <http://lepidopteraresearchfoundation.org/pdf/pdf38/38-015.pdf>
- [161] Hilt, N., & Fiedler, K. (2005): Diversity and composition of Arctiidae moth ensembles along a successional gradient in the Ecuadorian Andes. – *Diversity & Distributions* 11: 387-398. doi: 10.1111/j.1366-9516.2005.00167.x.
- [162] Brehm, G., Pitkin, L.M., Hilt, N., & Fiedler, K. (2005): Montane Andean rain forests are a global diversity hotspot of geometrid moths. – *J. Biogeogr.* 32: 1621-1627. doi: 10.1111/j.1365-2699.2005.01304.x.
- [163] Hilt, N., & Fiedler, K. (2006): Arctiid moth ensembles along a successional gradient in the Ecuadorian montane rain forest zone: how different are subfamilies and tribes? – *J. Biogeogr.* 33: 108-120. doi: 10.1111/j.1365-2699.2005.01360.x.
- [164] Hilt, N., Brehm, G., & Fiedler, K. (2006): Diversity and ensemble composition of geometrid moths along a successional gradient in the Ecuadorian Andes. – *J. Trop. Ecol.* 22: 1-12. doi: 10.1017/S0266467405003056.

- [165] Daniels, H., Gottsberger, G., & Fiedler, K. (2005): Nutrient composition of larval nectar secretions from three species of myrmecophilous butterflies. – *J. Chem. Ecol.* 31: 2805-2821. doi: 10.1007/s10886-005-8395-y.
- [166] Axmacher, J.C., Scheuermann, L., Schruppf, M., Lyaruu, H., Fiedler, K., & Müller-Hohenstein, K. (2006): Effects of fire on the diversity of geometrid moths on Mt. Kilimanjaro, Tanzania. – Pp. 69-75 in: SPEHN, E., LIEBERMAN, M., & KÖRNER, C. (eds.), *Land use change and mountain diversity*. CRC Press/Taylor and Francis.
- [167] Schlick-Steiner, B.C., Steiner, F.M., Moder, K., Bruckner, A., Fiedler, K., & Christian, E. (2006): Assessing ant assemblages: pitfall trapping versus nest counting (Hymenoptera, Formicidae). – *Insect. Soc.* 53: 274-281. doi: 10.1007/s00040-006-0869-6.
- [168] Hilt, N., Brehm, G., & Fiedler, K. (2007): Temporal dynamics of rich moth ensembles in the montane forest zone in southern Ecuador. – *Biotropica* 39: 94-104. doi: 10.1111/j.1744-7429.2006.00219.x.
- [169] Fiedler, K. (2006): Ant-associates of Palearctic lycaenid butterfly larvae (Hymenoptera: Formicidae; Lepidoptera: Lycaenidae) – a review. – *Myrmecologische Nachrichten* 9: 77-87.
- [170] Fiedler, K., Hilt, N., Brehm, G., & Schulze, C.H. (2007): Moths at tropical forest margins – how mega-diverse insect assemblages respond to forest disturbance and recovery. – Pp. 39-60 in: Tscharnatke, T., Leuschner, C., Zeller, M., Guhardja, E. & Bidin, A. (eds.), *The stability of tropical rainforest margins: Linking ecological, economic and social constraints of land use and conservation*. Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-30290-2_3.
- [171] Wiemers, M., & Fiedler, K. (2007): Does the DNA barcoding gap exist? A case study in blue butterflies (Lepidoptera: Lycaenidae). – *Frontiers Zool.* 4 (8): 1-16. doi: 10.1186/1742-9994-4-8.
- [172] Fiedler, K., F. Kuhlmann, B.C. Schlick-Steiner, F.M. Steiner & Gebauer, G. (2007): Stable N-isotope signatures of central European ants – assessing positions in a trophic gradient. – *Ins. Soc.* 54: 393-402. doi: 10.1007/s00040-007-0959-0.
- [173] Nöske, N. M., Hilt, N., Werner, F. A., Brehm, G., Fiedler, K., Sipman, H. J. M., & Gradstein, S. R. (2008): Disturbance effects on diversity of epiphytes and moths in a montane forest in Ecuador. – *Basic Appl. Ecol.* 9: 4-12. doi: 10.1016/j.baae.2007.06.014.
- [174] Brehm, G., Fiedler, K., Häuser, C. L. & Dalitz, H. (2008): Methodological challenges of a megadiverse mountain rain forest ecosystem. – Pp. 41-47 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_5.
- [175] Brehm, G., Homeier, J., Fiedler, K., Kottke, I., Illig, J., Nöske, N. M., Werner, F. & Breckle, S.-W. (2008): Mountain rain forests in southern Ecuador as a hotspot of biodiversity – Limited knowledge and diverging patterns. – Pp. 15-23 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_2
- [176] Fiedler, K., & Beck, E. (2008): Investigating gradients in ecosystem analysis. – Pp. 49-54 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_6
- [177] Fiedler, K., Brehm, G., Hilt, N., Süßenbach, D., & Häuser, C.L. (2008): Variation of diversity patterns across moth families along a tropical elevational gradient. – Pp. 167-179 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_17.
- [178] Günter, S., Cabrera, O., Weber, M., Stimm, B., Zimmermann, M., Fiedler, K., Knuth, J., Boy, J., Wilcke, W., Iost, S., Makeschin, F., Werner, F., Gradstein, R., & Mosandl, R. (2008): Natural forest management in Neotropical mountain rain forests – an ecological experiment. – Pp. 347-359 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_33.
- [179] Hilt, N., & Fiedler, K. (2008): Succession stages of faunal regeneration – a case study on megadiverse moths. – Pp. 443-449 in: Beck, E., Bendix, J., Kottke, I., Makeschin, F., & Mosandl, R. (eds.), *Gradients*

- in a tropical mountain ecosystem of Ecuador*. Ecol. Studies 198, Springer, Berlin, Heidelberg. doi: 10.1007/978-3-540-73526-7_42.
- [180] Häuser C.L., Fiedler, K., Bartsch, D., Brehm, G., Kling, M., Süßenbach, D., & Onore, G. (2008): Butterflies (Lepidoptera: Papilionoidea). Checklist Reserva Biológica San Francisco (Prov. Zamora-Chinchi, S. Ecuador). – Ecotropical Monographs 4: 145-154.
- [181] Fiedler, K., Brehm, G., Hilt, N., Süßenbach, D., Onore, G., Bartsch, D., Racheli, L. & Häuser, C.L. (2008): Moths (Lepidoptera: Arctiidae, Geometridae, Hedyliidae, Pyraloidea, Sphingidae, Saturniidae, and Uraniidae). Checklist Reserva Biológica San Francisco (Prov. Zamora-Chinchi, S. Ecuador). – Ecotropical Monographs 4: 155-214.
- [182] Axmacher, J.C., & Fiedler, K. (2008): Habitat type modifies geometry of elevational diversity gradients in geometrid moths (Lepidoptera: Geometridae) on Mt. Kilimanjaro, Tanzania. – Trop. Zool. 21: 243-251.
- [183] Wiemers, M., & Fiedler, K. (2008): Butterfly diversity of the Piedras Blancas National Park and its vicinity – a preliminary assessment (Lepidoptera: Papilionoidea & Hesperioidea). – Stapfia 88: 277-294. http://www.zobodat.at/pdf/STAPFIA_0088_0277-0294.pdf
- [184] Beck, J., & Fiedler K. (2009): Adult life spans of butterflies (Lepidoptera: Papilionoidea + Hesperioidea): broadscale contingencies with adult and larval traits in multi-species comparisons. – Biological Journal of the Linnean Society 96: 166-184. doi: 10.1111/j.1095-8312.2008.01102.x
- [185] Axmacher, J.C., Brehm, G., Hemp, A., Tünte, H., Lyaruu, H.V.M., Müller-Hohenstein, K., & Fiedler, K. (2009): Determinants of diversity in afrotropical herbivorous insects (Lepidoptera: Geometridae): plant diversity, vegetation structure or abiotic factors? – J. Biogeography 36: 337-349. doi: 10.1111/j.1365-2699.2008.01997.x.
- [186] Fellner, T., BOROVSKY, V., & FIEDLER, K. (2009): First records of the dacetine ant species *Pyramica argiola* (Emery, 1869) (Hymenoptera: Formicidae) from Austria. – Myrmecological News 12: 167-169.
- [187] Strutzenberger, P., Brehm, G., Bodner, F., Zimmermann, M., Wiemers, M. & Fiedler, K. (2009): DNA barcoding and molecular phylogeny of *Eois* moths (Geometridae) from southern Ecuador. In: HAUSMANN, A. (ed.): *Proceedings of the fifth Forum Herbulot 2008. Global strategies for plotting geometrid biodiversity in web-based databases* (Munich, ZSM, 24-28 June 2008). – Spixiana 32 (1): 131.
- [188] Schulze, C.H., Schneeweih, S., & Fiedler, K. (2010): The potential of land-use systems for maintaining tropical forest butterfly diversity. – Pp. 74-96 in: TSCHARNTKE, T., LEUSCHNER, C., VELDKAMP, E., FAUST, H., GUHARDJA, E., & BIDIN, A. (eds.): *Tropical rainforests and agroforests under global change – ecological and socio-economic valuations*. Springer, Berlin. doi: 10.1007/978-3-642-00493-3_3.
- [189] Bodner, F., Brehm, G., Homeier, J., Strutzenberger, P. & Fiedler, K. (2010): Caterpillars and host plant records for 59 species of Geometridae (Lepidoptera) from a montane rainforest in southern Ecuador. – J. Insect Sci. 10 (67): 1-22. www.insectscience.org/10.67. doi: 10.1673/031.010.6701.
- [190] Fiedler, K. (2010): Commentary: The coming and going of Batesian mimicry in a Holarctic butterfly clade. – BMC Biology 8 (122): 1-3. <http://www.biomedcentral.com/1741-7007/8/122>. doi: 10.1186/1741-7007-8-122.
- [191] Strutzenberger, P., Brehm, G., Bodner, F., & Fiedler, K. (2010): Molecular phylogeny of *Eois* (Lepidoptera, Geometridae): evolution of wing patterns and host plant use in a species rich group of Neotropical moths. – Zool. Scripta 39 (6): 603-620. doi: 10.1111/j.1463-6409.2010.00440.x.
- [192] Bodner, F., Mahal, S., Reuter, M., & Fiedler, K. (2010): Feasibility of a combined sampling approach for studying caterpillar assemblages – a case study from shrubs in the Andean montane forest zone. – J. Res. Lepid. 43: 27-35. www.lepidopteraresearchfoundation.org/journals/43/jrl_43_27_35.pdf
- [193] Laurance, W.F., Useche, D.C., Shoo, L.P., Herzog, S.K., Kessler, M., Escobar, F., Brehm, G., Axmacher, J.C., Chen, I.C., Arellano Gámez, L., Hietz, P., Fiedler, K., Pyrcz, T., Wolf, J., Merkord, C.L., Cardelus, C., Marshall, A.R., Ah-Peng, C., Aplet, G.H., del Coro Arizmendi, M., Baker, W.J., Barone, J., Brühl, C.A., Bussmann, R.W., Cicuzza, D., Eilu, G., Favila, M.E., Hemp, A., Hemp, C., Homeier, J., Huey, R.B., Hurtado, J., Jankowski, J., Kattán, G., Kluge, J., Krömer, T., Lees, D., Lehnert, M., Longino, J.T., Lovett, J., Martin, P.H., Patterson, B.D., Pearson, R.G., Peh, K.S.-H.,

- Richardson, B., Richardson, M., Samways, M., Senbeta, F., Smith, T.B., Utteridge, T., Watkins, J.E., Wilson, R., Williams, S.E., & Thomas, C.D. (2011): Global warming, elevational ranges and the vulnerability of tropical biota. – *Biol. Conserv.* 144: 548-557. doi: 10.1016/j.biocon.2010.10.010.
- [194] Beck, J., Brehm, G., & Fiedler, K. (2011): Links between the environment, abundance and diversity of Andean moths. – *Biotropica* 43: 208-217. doi: 10.1111/j.1744-7429.2010.00689.x.
- [195] Strutzenberger, P., & Fiedler, K. (2011): Temporal patterns of diversification in Andean *Eois*, a species-rich clade of moths (Lepidoptera, Geometridae). – *J. Evol. Biol.* 24: 919-925. doi: 10.1111/j.1420-9101.2010.02216.x
- [196] Strutzenberger, P., Brehm, G., & Fiedler, K. (2011): DNA barcoding based species delimitation increases species count of *Eois* (Geometridae) moths in a well-studied tropical mountain forest by up to 50%. – *Insect Science* 18: 349-362. doi: 10.1111/j.1744-7917.2010.01366.x.
- [197] Tista, M., & Fiedler, K. (2011): How to evaluate and reduce sampling effort for ants. – *J. Insect Conserv.* 15: 547-559. doi: 10.1007/s10841-010-9350-y.
- [198] Brehm, G., Bodner, F., Strutzenberger, P., Hünefeld, F., & Fiedler, K. (2011): Neotropical *Eois* (Lepidoptera: Geometridae): Checklist, biogeography, diversity, and description patterns. – *Ann. Entomol. Soc. Amer.* 104: 1091-1107. doi: 10.1603/AN10050.
- [199] Truxa, C., & Fiedler, K. (2012): Attraction to light – from how far do moths (Lepidoptera) return to weak artificial sources of light? – *Eur. J. Ent.* 109: 77-84.
- [200] Truxa, C., & Fiedler, K. (2012): Down in the flood? How moth communities are shaped in temperate floodplain forests. – *Insect Conserv. Divers.* 5: 389-397. doi: 10.1111/j.1752-4598.2011.00177.x
- [201] Plank, M., Fiedler, K., & Reiter, G. (2012): Use of forest strata by bats in temperate forests. – *J. Zool.* 286: 254-162. doi: 10.1111/j.1469-7998.2011.00859.x.
- [202] Haschek, C., Drapela, T., Schuller, N., Fiedler, K., & Frank, T. (2012): Carabid beetle condition, reproduction and density in winter oilseed rape affected by field and landscape parameters. – *J. Appl. Ent.* 136: 665-674. doi: 10.1111/j.1439-0418.2011.01694.x.
- [203] Fiedler, K. (2012): The host genera of ant-parasitic Lycaenidae butterflies – a review. – *Psyche* 2012 (ID 153975): 1-10. doi: 10.1155/2012/153975.
- [204] Strausz, M., Fiedler, K., Franzén, M., & Wiemers, M. (2012): Habitat and host plant use of the Large Copper Butterfly *Lycaena dispar* in an urban environment. – *J. Insect Conserv.* 16: 709-721. doi: 10.1007/s10841-012-9456-5.
- [205] Pellissier, L., Litsios, G., Fiedler, K., Pottier, J., Dubuis, A., Pradervand, J.-N., Salamin, N., & Guisan, A. (2012): Loss of interactions with ants under cold climate in a regional myrmecophilous butterfly fauna. – *J. Biogeogr.* 39: 1782–1790. doi: 10.1111/j.1365-2699.2012.02743.x
- [206] Pellissier, L., Rasmann, S., Litsios, G., Fiedler, K., Dubuis, A., Pottier, J., & Guisan, A. (2012): High host-plant nitrogen content: a prerequisite for the evolution of ant-caterpillar mutualism? – *J. evol. Biol.* 25: 1658-1666. doi: 10.1111/j.1420-9101.2012.02555.x.
- [207] Pellissier, L., Fiedler, K., Ndribe, C., Dubuis, A., Pradervand, J.-N., Guisan, A., & Rasmann, S. (2012): Shifts in species richness, herbivore specialisation and plant resistance along elevation gradients. – *Ecology & Evolution* 2: 1818-1825. doi: 10.1002/ece3.296.
- [208] Fiedler, K., & Truxa, C. (2012): Species richness measures fail in resolving diversity patterns of speciose forest moth assemblages. – *Biodiversity & Conservation* 21: 2499-2508. doi: 10.1007/s10531-012-0311-5.
- [209] Bodner, F., Strutzenberger, P., Brehm, G., & Fiedler, K. (2012): Species richness and host specificity among caterpillar ensembles on shrubs in the Andes of southern Ecuador. – *Neotrop. Ent.* 41: 375-385. doi: 10.1007/s13744-012-0066-4.
- [210] Strutzenberger, P., Brehm, G., & Fiedler, K. (2012): DNA barcode sequencing from old type specimens as a tool in taxonomy: A case study in the diverse genus *Eois* (Lepidoptera: Geometridae). – *PLoS ONE* 7 (11): e49710. doi: 10.1371/journal.pone.0049710.
- [211] Grill, A., Cerny, A., & Fiedler, K. (2013): Hot summers, long life: egg laying strategies of *Maniola* butterflies are affected by geographic provenance rather than adult diet. – *Contrib. Zool.* 82 (1) 27-36.

- [212] Brehm, G., Strutzenberger, P., & Fiedler, K. (2013): Phylogenetic diversity of geometrid moths decreases with elevation in the tropical Andes. – *Ecography* 36: 1247-1253. doi: 10.1111/j.1600-0587.2013.00030.x
- [213] Fiedler, K., & Strutzenberger, P. (2013): Past dynamics of speciation in Andean mountain rainforests. – Pp. 67-79 in: BENDIX, J., BECK, E., BRÄUNING, A., MAKESCHIN, F., MOSANDL, R., SCHEU, S. & WILCKE, W. (eds.), *Ecosystem services, biodiversity and environmental change in a tropical mountain ecosystem of South Ecuador*. Ecol. Studies 221, Springer, Berlin, Heidelberg, New York. doi: 10.1007/978-3-642-38137-9_6.
- [214] Fiedler, K. (2013): Die Nachtfalterfauna im Gebiet des tropischen Bergregenwaldes in Ecuador – der Einfluss des Menschen. – Pp. 85-96 in: Bayer. Akademie der Wissenschaften (eds.), *Schutz und Nutzung von Tropenwäldern*. Rundgespräche der Kommission für Ökologie, vol. 42. Friedrich Pfeil, München.
- [215] Carneiro, E., Mielke, O.H.H., Casagrande, M.M., & Fiedler K. (2014): Skipper richness (Hesperiidae) along elevational gradients in Brazilian Atlantic forest. – *Neotropical Entomology* 43: 27-38. doi: 10.1007/s13744-013-0175-8.
- [216] Milaković, I., Karrer, G., & Fiedler, K. (2014): Management of naturalized roadside populations of invasive *Ambrosia artemisiifolia* by mowing. – *Weed Research* 54: 256-264. doi: 10.1111/wre.12074.
- [217] Carneiro, E., Mielke, O.H.H., Casagrande, M.M., & Fiedler, K. (2014): Community structure of skipper butterflies (Lepidoptera, Hesperiidae) along elevational gradients in Brazilian Atlantic Forest reflects vegetation type rather than altitude. – *PLoS ONE* 9(10): e108207. doi: 10.1371/journal.pone.0108207.
- [218] Haeler, E., Fiedler, K., & Grill, A. (2014): What prolongs a butterfly's life? Trade-offs between dormancy, fecundity and body size. – *PLoS ONE* 9(11): e111955. doi: 10.1371/journal.pone.0111955.
- [219] Polic, D., Fiedler, K., Nell, C., & Grill, A. (2014): Mobility of ringlet butterflies in high-elevation alpine grassland: Effects of habitat barriers, resources and age. – *J. Insect Conserv.* 18: 1153-1161. doi: 10.1007/s10841-014-9726-5.
- [220] Milaković, I., Karrer, G., & Fiedler, K. (2014): Fine tuning of mowing regime, a method for the management of the invasive *Ambrosia artemisiifolia* L. at different population densities. – *Weed Biology and Management* 14: 232–241. doi:10.1111/wbm.1.
- [221] Kreuzinger, A., Fiedler, K., Letsch, H. & Grill, A. (2015): Tracing the radiation of *Maniola* (Nymphalidae) butterflies: new insights from phylogeography hint at one single incompletely differentiated species complex. – *Ecology and Evolution* 5: 46-58. doi: 10.1002/ece3.1338.
- [222] Zöchling, A., & Fiedler, K. (2015): Einfluss der Nutzungsintensität von Almen auf Tagfalter-Gemeinschaften (Lepidoptera: Papilionoidea) im Nationalpark Gesäuse (Österreich). – *Entomologica Austriaca* 22: 7-26. http://www.zobodat.at/pdf/ENTAU_0022_0007-0026.pdf
- [223] Hülber, K., Haider, J.A., Hager, T.E., Dullinger, S., & Fiedler, K. (2015): Insect herbivory in alpine grasslands is constrained by community and host traits. – *J. Veget. Sci.* 26: 663–673 doi: 10.1111/jvs.12265.
- [224] Seifert, C.L., Lehner, L., Adams, M.-O., & Fiedler, K. (2015): Predation on artificial caterpillars is higher in countryside than near-natural forest habitat in lowland south-western Costa Rica. – *J. Trop. Ecol.* 31 (3): 281-284. doi: 10.1017/S0266467415000012.
- [225] Bodner, F., Brehm, G., & Fiedler, K. (2015): Many caterpillars in a montane rainforest in Ecuador are not classical herbivores. – *J. Trop. Ecol.* 31 (5): 473-476. doi: 10.1017/S0266467415000243.
- [226] Adams, M.-O. & Fiedler, K. (2015): The value of targeted reforestations for local insect diversity – A case study from the Ecuadorian Andes. – *Biodivers. Conserv.* 24 (11): 2709-2734. doi: 10.1007/s10531-015-0958-9.
- [227] Seifert, C.L., Bodner, F., Brehm, G., & Fiedler, K. (2015): Host plant associations and parasitism of South Ecuadorian *Eois* species (Lepidoptera: Geometridae) feeding on *Peperomia* (Piperaceae). – *J. Insect Sci.* 15 (119): 1-7. doi: 10.1093/jisesa/iev098.
- [228] Todisco, V., Fiedler, K., Puliga, E., Dinca, V., Voda, R., & Grill, A. (2015): The origin of the Sardinian Blue, *Pseudophilotes barbagiae* (Lepidoptera: Lycaenidae): Out-of-Europe or out-of-Africa? – *Genome* 58: 288-289. doi: 10.1139/gen-2015-0087.

- [229] Seifert, C.L., Schulze, C. H., Dreschke, T., Frötscher, H., & Fiedler, K. (2016): Day vs. night predation on artificial caterpillars in primary rainforest habitats – an experimental approach. – *Ent. exp. appl.* 158: 54-59. doi: 10.1111/eea.12379.
- [230] Truxa, C., & Fiedler, K. (2016): Massive structural redundancies in species composition patterns of floodplain forest moths. – *Ecography* 39: 253-260. doi: 10.1111/ecog.01430.
- [231] Uhl, B., Wölfling, M., Fiala, B., & Fiedler, K. (2016): Micro-moth communities mirror environmental stress gradients within a Mediterranean nature reserve. – *Basic Appl. Ecol.* 17: 273-281. doi: 10.1016/j.baae.2015.10.002.
- [232] Rodríguez-Castañeda, G., Brehm, G., Fiedler, K. & Dyer, L.A. (2016): Ant predation on herbivores through a multitrophic lens: how effects of ants on plant herbivore defense and parasitoid populations vary along elevational gradients. – *Current Opinion Ins. Sci.* 14: 73-80. doi: 10.1016/j.cois.2016.02.001
- [233] Brehm, G., Hebert, P.D.N., Colwell, R.K., Adams, M.-O., Bodner, F., Friedemann, K., Möckel, L., & Fiedler, K. (2016): Turning up the heat at a hotspot: DNA barcodes reveal 80% more species of geometrid moths along an Andean elevational gradient. – *PLoS ONE* 11(3): e0150327. doi: 10.1371/journal.pone.0150327.
- [234] Adams, M.-O., & Fiedler, K. (2016): Low herbivory in targeted reforestation in the Andean highlands of southern Ecuador. – *PLoS ONE* 11(3): e0151277. doi: 10.1371/journal.pone.0151277.
- [235] Knoll, T., Fiedler, K., & Reiter, G. (2016): Fledermausgemeinschaften und Rufaktivität im Wildnisgebiet Dürrenstein und in umliegenden Wirtschaftswäldern. – *Silva Fera* 5: 23-34.
- [236] Fies, R., Rabl, D., Schulze, C.H., & Fiedler, K. (2016): Summer floods shape meadow butterfly communities in a floodplain nature reserve in Central Europe. – *J. Insect Conserv.* 20: 433-445. doi: 10.1007/s10841-016-9876-8.
- [237] Colwell, R.K., Gotelli, N.J., Ashton, L.A., Beck, J., Brehm, G., Fayle, T.M., Fiedler, K., Forister, M.L., Kessler, M., Kitching, R.L., Klimes, P., Kluge, J., Longino, J.T., Maunsell, S.C., McCain, C.M., Moses, J., Noben, S., Sam, K., Sam, L., Shapiro, A.M., Wang, X., & Novotny, V. (2016): Midpoint attractors and species richness: Modeling the interaction between environmental drivers and geometric constraints. – *Ecol. Letts.* 19(9): 1009-1022. doi: 10.1111/ele.12640.
- [238] Seifert, C.L., Lehner, L., Bodner, F. & Fiedler, K. (2016): Caterpillar assemblages on *Chusquea* bamboos in southern Ecuador: abundance, guild structure, and the influence of host plant quality. – *Ecol. Ent.* 41(6): 698-706. doi: 10.1111/een.12345.
- [239] Adams, M.-O., Seifert, C.L., Lehner, L., Truxa, C., Wanek, W. & Fiedler, K. (2016): Stable isotope signatures reflect dietary diversity in European forest moths. – *Frontiers Zool.* 13: 37. doi: 10.1186/s12983-016-0170-0. <http://rdcu.be/noQE>
<http://frontiersinzoology.biomedcentral.com/articles/10.1186/s12983-016-0170-0>
- [240] Nepel, M., Voglmayr, H., Blatrix, R., Longino, J.T., Fiedler, K., Schönenberger, J., & MAYER, V.E. (2016): Ant-cultivated Chaetothyrals in hollow stems of myrmecophytic *Cecropia* sp. trees – diversity and patterns. – *Fungal Ecol.* 23: 131-140. doi: 10.1016/j.funeco.2016.07.007.
- [241] Beck, J., Liedtke, H.C., Widler, S., Altermatt, F., Loader, S., Hagmann, R., Lang, S., & Fiedler, K. (2016): Patterns or mechanisms? Bergmann's and Rapoport's rule in moths along an elevational gradient. – *Community Ecol.* 17 (2): 137-148. doi: 10.1556/168.2016.17.2.2.
- [242] Brehm, G., Fiedler, K. & Gaona, F.P. (2016): Andean rainforests: studying the global hotspot of Lepidopteran diversity. – Pp. 110-115 in: F.X. BOGNER, J. BENDIX & E. BECK (eds.): Biodiversity hotspot tropical mountain rainforest. *Naturaleza y Cultura, Loja* (Ecuador).
- [243] Wölfling, M., Becker, M.C., Uhl, B., Traub, A., & Fiedler, K. (2016): How differences in the settling behaviour of moths (Lepidoptera) may contribute to sampling bias when using automated light traps. – *Eur. J. Ent.* 113: 502-506. doi: 10.14411/eje.2016.066.
- [244] Brehm, G. & Fiedler, K. (2016): Ordnung: Lepidoptera, Schmetterlinge. Pp. 464-496 in: Schaefer, M. (ed.), *Brohmer – Fauna von Deutschland*. 24th ed., Quelle & Meyer, Wiebelsheim.
- [245] Beck, J., McCain, C.M., Axmacher, J.C., Ashton, L.A., Bärtschi, F., Brehm, G., Choi, S.-W., Cizek, O., Colwell, R.K., Fiedler, K., Francois, C.L., Highland, S., Holloway, J.D., Intachat, J., Kadlec, T., Kitching, R.L., Maunsell, S.C., Merckx, T., Nakamura, A., Odell, E., Sang, W., Toko, P.S., Zamecnik, J.,

Zou, Y. & Novotny, V. (2017): Elevational species richness gradients in a hyperdiverse insect taxon: a global meta-study on geometrid moths. – *Global Ecol. Biogeogr.* 26 (in press). doi: 10.1111/geb.12548.