	<b>ABHANDLUNGEN DER GEOLOGISCHEN BUNDESANSTALT</b>					
	Abh. Geol. B.-A.	ISSN 0378-0864	ISBN 3-85316-007-7	Band 56/2	S. 465-474	Wien, Dezember 1999
<b>Geologie ohne Grenzen</b> Festschrift 150 Jahre Geologische Bundesanstalt			<b>Redaktion:</b> Harald Lobitzer & Pavol Grecula			

## Biostratigraphy of Fissure Fillings in the Ernstbrunn Limestone of the Waschberg Zone (Lower Austria)

THOMAS HOFMANN\*), LILIAN ŠVÁBENICKÁ\*\*) & LENKA HRADECKÁ\*\*)

2 Text-Figures and 3 Plates

*Outer Western Carpathians*  
*Waschberg Zone*  
*Tithonian*  
*Upper Cretaceous*  
*Biostratigraphy*  
*Foraminifera*  
*Calcareous Nannofossils*

Österreichische Karte 1: 50.000  
Blatt 24

### Inhalt

Abstract	465
Zusammenfassung	465
1. Geological setting	466
2. Fissure fillings in the Ernstbrunn Limestone of Lower Austria – an overview	466
3. Description of the localities	466
3.1. Ernstbrunn Quarry	466
3.2. Staatz Klippe	467
4. Material and Methods	467
5. Results	467
5.1. Ernstbrunn Quarry	467
5.1.1. Foraminifera	467
5.1.2. Calcareous Nannofossils	468
5.2. Staatz Klippe	470
5.2.1. Calcareous Nannofossils	470
6. Conclusions	470
Acknowledgement	472
References	472
Appendix	474

### Stratigraphie von Spaltenfüllungen im Ernstbrunner Kalk der Waschbergzone (Niederösterreich)

#### Zusammenfassung

Die biostratigraphische Auswertung von Foraminiferen und Nannofossilien aus mergeligen Spaltenfüllungen im Ernstbrunner Kalk (Oberjura) der Waschberg Zone brachte folgende Ergebnisse: Im Steinbruch Ernstbrunn konnten Sedimente aus dem Tithon, als auch aus der Oberkreide (Santon und Oberes Campan) gefunden werden. Das tithone Alter der Spaltenfüllungen konnte mit einer Nannofloraassoziation, bei der *Watznaueria* dominiert, belegt werden. Santon ist durch *Rucinolithus wissei* und *Orastrum campanensis* des UC11c-UC12 Zonenintervalls nachgewiesen. Diese Daten sind auch durch Foraminiferen aus dem Oberen Santon bis Unteren Campan untermauert. Oberes Campan ist durch die Globotruncanita calcarata Zone, bzw. durch UC15c-d<sup>TP</sup> Zone mittels Nannofossilien nachgewiesen. In der Staatzer Klippe belegen Nannofossilien-Vergesellschaftungen mit *Rucinolithus* spp. und *Reinhardtites anthophorus* die UC11c Subzone des Unteren Santons. Das gemeinsame Auftreten von Nannoplanktonarten hoher und niedriger Breiten deutet auf einen Ablagerungsraum im Übergangsbereich der tethyalen-mediterranen Bioprovinz mit starkem borealen Einfluß hin.

#### Abstract

Biostratigraphic study (foraminifers and calcareous nannofossils) of marly sediments collected from the dissolution-dilated fractures in the Ernstbrunn Limestone (Upper Jurassic) of the Waschberg Zone provided the following information: In the Ernstbrunn Quarry, both Tithonian

\*) Geological Survey of Austria, Rasumofskygasse 23, A-1030 Wien, Austria (email: thofmann@cc.geolba.ac.at)

\*\*) Czech Geological Survey, Klárov 131/3, P.O.Box 85, CZ-118 21 Praha 1, Czech Republic (email: svab@cgu.cz, hradecka@cgu.cz)

and Upper Cretaceous (Santonian and Upper Campanian) sediments were found. Tithonian deposits provided nannofossil association dominated by the genus *Watznaueria*. The Santonian is documented by nannofossils with *Rucinolithus wissei* and *Orastrum campanensis* of the UC11c-UC12 zone interval and by foraminifers of the Upper Santonian – Lower Campanian age. The Upper Campanian is confirmed by the *Globotruncanita calcarata* Zone and nannofossils of the UC15c-d<sup>1F</sup> Zone. In the Staatz Klippe, nannofossil associations with *Rucinolithus* spp. and *Reinhardtites anthophorus* evidence the UC11c Subzone of the Lower Santonian age. The common occurrence of high- and low-latitude nannoplankton species indicates that the depositional area was situated within the Tethyan/Mediterranean province with a strong influence from the Boreal area.

## 1. Geological setting

The Waschberg Zone (Text-Fig. 1) represents the southern part of the Waschberg-Ždánice Unit of the Outer Western Carpathians. It is a highly tectonized zone in the north eastern part of Lower Austria and extends from the Waschberg (338 m) near Stockerau in northeastern direction to Southern Moravia. So, the Leiser Berge, the dominant Klippe of Staatz as well as the Mountains of Falkenstein are part of the Waschberg Zone. The zone also includes the mountains around Mikulov (Pavlovské vrchy Hills) N of the Austrian/Czech border. Due to the extension to the Czech Republic the Waschberg Zone is called Waschberg-Ždánice Unit, which expresses in better way its common features. This was made evident by ELIÁŠ et al. (1990, page 44): "The Waschberg Zone (GRILL 1953, 1968) is the only zone where the junction of the Eastern Alps with the Carpathians can be traced on surface. A relationship of the Waschberg Zone with the Ždánice Unit is evident by their lithostratigraphic and facies development."


According to STEININGER et al. (1986) the Waschberg Zone is a part of the Molassezone. In detail the authors described it to the "allochthonous Molassezone" which was scraped from its footwall during late Alpine orogenic movements. As a result of this, Waschberg Zone is situated between the Molassezone in the west – where it is overthrust – and the Vienna Basin in the east. In the southern part the Flysch Zone is overthrust upon the Waschberg Zone. These overthrusts can be dated to the Lower Miocene (Karpatian). According to FUCHS (1980) various phases were determined within the period of the Eggenburgian and Karpatian in the Lower Miocene.

The most significant elements of the Waschberg Zone are the different isolated klippen composed of Jurassic sediments (Ernstbrunn Limestone and Klentnice Member). The Ernstbrunn Limestone represents a part of shallow Upper Jurassic carbonate platform fringing the southeastern part of the Bohemian Massif. After the sedimentation stopped in the Upper Jurassic-Lower Cretaceous (ELIÁŠ, 1992) intensive karstification took place until a new transgression of the Upper Cretaceous Klement and Pálava formations (STRÁNIK et al., 1996; SUMMESBERGER et al., 1999). As a result, Cretaceous and even Tertiary sediments have been known from fissures in the Ernstbrunn Limestone since the 19<sup>th</sup> century.

## 2. Fissure fillings in the Ernstbrunn Limestone of Lower Austria – an overview

BACHMAYER (1964) described a fissure filling (width 5.6 m) of Quaternary age with redeposited nannoflora (determined by E. KAMPTNER) from the Cretaceous and Tertiary, with some molluscs indicating the loess environment of Staatz.

As a result of an UNESCO-Course held at the Geological Survey of Austria, AL-SHAIBANI (1971) described Upper Cretaceous Foraminifera from "an old Quarry in Dörfles" (from which of the five quarries in Dörfles the material came is not clear). He presented a list with the following taxa indi-



Text-Fig. 1.  
Location of working area in Lower Austria.

cating Maastrichtian age: *Globotruncana gansseri* BOLLI (= *Gansserina gansseri*), *Bolivinoidea draco draco* MARSSON etc. Nevertheless, the presence of species such as *Reussella szajnochae* (GRYZBOWSKI) and *Globotruncana arca* (CUSHMAN) indicate rather Lower Maastrichtian age of sediments from Dörfles.

HOFMANN (1990) described two fissures with Tertiary sediments (marls of different ages) in the Dörfles region – abandoned quarries of Dörfles III and V. In the northwestern part of the Dörfles III quarry R. BRAUNSTEIN (in: HOFMANN, 1990, Fig. 9) determined rich, moderately preserved nannoflora with *Cruciplacolithus tenuis* and *Chiasmolithus danicus* (= *Sullivania danica*) of the Lower Paleocene age. In the nearby quarry, Dörfles V, (see Fig. 12 in: HOFMANN, 1990), an equally rich but poorly preserved floral association with *Isthmolithus recurvus* and *Dictyococcites bisectus* of the Upper Eocene age was found.

## 3. Description of the localities

### 3.1. Ernstbrunn Quarry

The quarry Ernstbrunn, which is known as "Kalkwerk II" or "Werk II" (BACHMAYER, 1940) is owned by the "Ernstbrunner Kalktechnik – Kalkgewerkschaft in Ernstbrunn Ges.m.b.H". The white limestone is used in the industry (BULLINGER, 1997). Five quarry levels are worked at present. At level 4

and 5 an approximately 2 m intercalation of black (sample No. A/1998; very dark - grey to black claystone with silty admixture) to greenish, soft marly sediments with blocks of the Ernstbrunn Limestone striking from south to north dipping at 70° to 80° is found. This feature was documented by SAUER et al. (1992) in Figures 99, 103 and 104.

### 3.2. Staatz Klippe

A thin (approx. 50 cm) steeply inclined fissure of greenish marls and claystones (sample No. 3R/98) with breccia of Ernstbrunn Limestone within the Ernstbrunn Limestone is exposed in the middle part of northern wall in the abandoned quarry of Staatz, which is used as an open air theatre now (Text-Fig. 2).

In addition, some samples (3/1994, 1R/1998) were taken in the northern part where rock (limestones) is stabilized by shotcrete.

A large fissure with brownish marls and breccias of limestones is located in its southern part. Because of the large dimension, this could correspond with the fissure described by BACHMAYER (1964).

## 4. Material and Methods

Material was collected within the framework of the bilateral cooperation between the Czech and Austrian Geological surveys.

Samples for foraminiferal study were disintegrated at the laboratory of the Geological Survey in Vienna using standard washing method. Foraminifers were separated under binocular microscope and photographs of species were taken using scanning electron microscope in the Laboratory of Czech Geological Survey in Prague. Planktonic zonation of ROBASYNSKI & CARON (1995) was used for correlation of Cretaceous sediments.

Samples for nannofossil research were processed in the Laboratory of Czech Geological Survey, Prague. Smear slides were prepared using standard method of decantation, samples were inspected under light microscope Nikon at 1,000x magnification. Cretaceous biostratigraphic data were correlated with the UC zones sensu BURNETT (1998). Data concerning province appurtenance of Cretaceous nannofossil species are interpreted mainly according to WIND (1979), WATKINS (1992), WATKINS et al. (1996) and BURNETT (1998).

## 5. Results

Samples were collected from marly sediments that were available in narrow dissolution fractures in Ernstbrunn Limestone of localities of Ernstbrunn, Staatz, Südmährenkreuz and Falkenstein. Sediments of Ernstbrunn Quarry provided both foraminifers and nannofossils, Staatz Klippe yielded only calcareous nannofossils and Falkenstein and Südmährenkreuz were free of both foraminifers and calcareous nannofossils.

### 5.1. Ernstbrunn Quarry

#### 5.1.1. Foraminifera

Upper Cretaceous sediments, which were deposited in fissures in the Ernstbrunn Limestone (Ernstbrunn-Kalkwerk II, samples Nos. 1, 1A, 3, 4 and 5) contained foraminiferal assemblages with low diversity of species but with relatively well-preserved foraminiferal tests.


Ernstbrunn-Kalkwerk II, sample No. 1A:

Planktonic species were more abundant in comparison with other samples from this locality. Benthos was represented by important species such as *Bolivina incrassata* REUSS, *Gavelinella monterelensis* (MARIE), *G. stelligera* (MARIE), *G. pertusa* (MARSSON), *G. clementiana costata* (MARIE), *Bolivinooides delicatus* CUSHMAN, *B. decoratus* (JONES), *B. granulatus* HOFKER, *Stensioeina pommerana* BROTZEN, *Vaginulina trilobata* (D'ORBIGNY) and *Reussella szajnochae* (GRZYBOWSKI) etc. Especially three species of genus *Bolivinooides* are cosmopolitan in the Campanian sediments (HANZLIKOVÁ 1972). Among planktonic foraminifers the following taxa dominated in the foraminiferal assemblage: *Globotruncana linneiana* (D'ORBIGNY), *Archaeoglobigerina cretacea* (D'ORBIGNY), *Rugoglobigerina rugosa* (PLUMMER), *Rosita fornicata* (PLUMMER) *Globotruncanita calcarata* CUSHMAN, *Heterohelix striata* (EHRENBERG), *H. navarroensis* LOEBLICH, *Globigerinelloides ultramica* (SUBBOTINA) and *G. aspera* (EHRENBERG).

The foraminiferal assemblage from Ernstbrunn-Kalkwerk II, sample No. 1A belongs to the *Globotruncanita calcarata* total range zone sensu ROBASYNSKI & CARON (1995) of the Upper Campanian age on the basis of the presence of *G. calcarata* CUSHMAN.

Ernstbrunn-Kalkwerk II, samples Nos. 1, 3, 4 and 5

The studied samples contained poorly preserved tests of foraminifers (rolled, dissolved, mechanically damaged), fragments of echinodermata and sponges.



Text-Fig. 2.  
Position of samples at the abandoned quarry of Staatz.

The foraminiferal assemblages had a similar but not identical character and the age of sediments from these samples is probably Campanian. Planktonic species were not present in high quantities as it was the case in Ernstbrunn-Kalkwerk II, sample No. 1A. Several species found in the foraminiferal assemblage from Ernstbrunn-Kalkwerk II, sample No. 1A such as *Gavelinella clementiana costata* (MARIE), *G. monterelensis* (MARIE), *Bolivina incrassata* REUSS, *Bandyella greatvalleyensis* (TRUJILLO), *Rosita fornicata* (PLUMMER) etc. were found in these four samples too.

Nevertheless, stratigraphic important taxa such as *G. calcarata* CUSHMAN and benthic species such as *Reussella szajnochae* (GRZYBOWSKI) and *Stensioeina pommerana* BROTZEN (GASINSKI et al., 1999) were missing. Representatives of the family Nodosariidae (*Fronicularia*, *Nodosaria*) were more frequent. On the basis of the occurrence of *Arenobulimina d'orbigny* (REUSS) and *Stensioeina exsculpta* (REUSS) in Ernstbrunn-Kalkwerk II, samples Nos. 1 and 5, an older age (Upper Santonian – Lower Campanian) is supposed for these samples (HANZLÍKOVÁ 1972) than for sample Ernstbrunn-Kalkwerk II, sample No. 1A.

Unfortunately, stratigraphically important planktonic species could not be identified. Hence, it was not possible to define an exact stratigraphic range of these samples.

### 5.1.2. Calcareous Nannofossils

Samples obtained from marly sediments transgressing the Ernstbrunn Limestone provided the following nannofossil assemblages:

1. Monogeneric association of Tithonian age dominated by *Watznaueria barnesae* and also containing *W. biporta*, *W. britannica*, *W. manivitae* and *Cyclagelosphaera margerelii* was found in grey to black claystones with silty admixture (Kalkwerk II, level No. 5, sample No. A/1998).

2. Association with abundant *Watznaueria barnesae* (> 10 %) and relative high numbers of *Rucinolithus wissei*, *R. hayi*, *Orastrum campanensis* and *Micula decussata* (1–5 %) which is associated by *Kamptnerius magnificus*, *Gartnerago obliquum*, *Reinhardtites anthophorus*, *Cylindralithus nudus*, *Grantarhabdus coronadventis*, *Arkhangelskiella specillata*

and others gives evidence for the Lower Santonian, the UC11c – UC12 (lower part) zones interval (Kalkwerk II, level No. 5, sample No. 1). Nannofossil preservation (overgrowth) is very similar to another one observed in the Santonian sediments of the locality of Staatz Klippe.

3. Rich and well preserved nannofossils of the Upper Campanian age. Assemblages of high species diversity with *Broinsonia parca constricta*, *Arkhangelskiella cymbiformis*, *Stauroolithites aachenus*, *Angulofenestrellithus snyderi*, *Reinhardtites anthophorus*, *R. levis*, *Eiffellithus eximius*, *Micula swastica*, *Prediscosphaera grandis*, *Petrarhabdus copulatus*, *Markalius inversus* and others were found in the fissure fillings in Kalkwerk II, samples Nos. 1A and 4. High number of low-latitude species, such as *Uniplanarius sissinghii*, *Ceratolithoides aculeus*, *C. quasiarcuatus* and *C. sesquipedalis* indicates that the depositional area was situated within the Tethyan/Mediterranean bioprovince. Nevertheless, common occurrence of high-latitude species, among others *Orastrum campanensis*, *Biscutum magnum*, *Prediscosphaera stoveri*, *Neocrepidolithus watkinsii* or *Monomarginatus quaternarius* gives evidence for strong influence from the Boreal bioprovince. By virtue of the common occurrence of high- and low-latitude species, both Boreal and Tethyan zonation was used for biostratigraphical correlations: the UC15d<sup>BP</sup> Zone and UC15c<sup>TP</sup> Zone, respectively, which points to the lower part of Upper Campanian. The above mentioned nannofossil association is similar in its character to those of the same age from the type locality of the Pálava Formation, Zďánice Unit, Outer Western Carpathians (ŠVÁBENICKÁ in STRÁNIK et al., 1996). Furthermore, the assemblage of Kalkwerk II, sample No. 3 yielded species *Uniplanarius trifidus* which gives evidence for the younger UC15d<sup>TP</sup> Zone within the lower part of Upper Campanian.


The occurrence of species *Petrarhabdus copulatus*, which is only rarely mentioned from the northern hemisphere, was reported by ŠVÁBENICKÁ (1998). This species was observed in low numbers in the Outer Western Carpathians, forming characteristic component of assemblages where mixing of low- and high-latitude nannofossils is evident. Here, it can be even used for biostratigraphic conclusions in the Upper Campanian.

4. Finally, intercalations of grey-green claystones (Kalkwerk II, level No. 5, sample No. B/1998) which were found

## Plate 1: Foraminifers from Ernstbrunn

- Fig. 1: *Gavelinella monterelensis* (MARIE); dorsal view, Ernstbrunn-Kalkwerk II, sample No. 1A, x60.  
 Figs. 2, 3: *Gavelinella stelligera* (MARIE); 2 – ventral view, 3 – dorsal view, Ernstbrunn-Kalkwerk II, sample No. 1A, x120.  
 Fig. 4: *Bandyella greatvalleyensis* (TRUJILLO); sample Ernstbrunn-Kalkwerk II, sample No. 1A, x80.  
 Fig. 5: *Vaginulina trilobata* (D'ORBIGNY); Ernstbrunn-Kalkwerk II, sample No. 1A, x40.  
 Fig. 6: *Gavelinopsis bembix* (MARSSON); Ernstbrunn-Kalkwerk II, sample No. 1A, x70.  
 Fig. 7: *Stensioeina pommerana* BROTZEN; lateral view, Ernstbrunn-Kalkwerk II, sample No. 1A, x120.  
 Fig. 8: *Astacolus cretaceus* (CUSHMANN); Ernstbrunn-Kalkwerk II, sample No. 1A, x90.  
 Fig. 9: *Patelina subcretacea* CUSHMANN-ALEXANDER; Ernstbrunn-Kalkwerk II, sample No. 5, x120.  
 Fig. 10: *Bolivina incrassata* REUSS; Ernstbrunn-Kalkwerk II, sample No. 3, x70.  
 Fig. 11: *Bolivinoidea delicatulus* CUSHMANN; Ernstbrunn-Kalkwerk II, sample No. 1A, x120.  
 Fig. 12: *Bolivinoidea granulatus* HOFKER; Ernstbrunn-Kalkwerk II, sample No. 1A, x110.  
 Fig. 13: *Heterohelix striata* (EHRENBERG); Ernstbrunn-Kalkwerk II, sample No. 1A, x110.  
 Fig. 14: *Stensioeina exsculpta* (REUSS); Ernstbrunn-Kalkwerk II, sample No. 1, x120.  
 Fig. 15: *Gavelinella clementiana costata* (MARIE); Ernstbrunn-Kalkwerk II, sample No. 4, x100.  
 Fig. 16: *Bolivinoidea decoratus* (JONES); Ernstbrunn-Kalkwerk II, sample No. 1A, x120.  
 Fig. 17: *Pullenia cretacea* CUSHMANN; Ernstbrunn-Kalkwerk II, sample No. 4, x170.  
 Fig. 18: *Gavelinella thalmani* (BROTZEN); Ernstbrunn-Kalkwerk II, sample No. 5, x140.  
 Fig. 19: *Globotruncanella calcarata* CUSHMANN; Ernstbrunn-Kalkwerk II, sample No. 1A, x100.  
 Fig. 20: *Globigerinelloides prairiehillensis* PESSAGNO; Ernstbrunn-Kalkwerk II, sample No. 5, x100.  
 Fig. 21: *Globotruncanella subspinosa* (PESSAGNO); Ernstbrunn-Kalkwerk II, sample No. 1A, x100.  
 Fig. 22: *Archaeoglobigerina cretacea* (D'ORBIGNY); Ernstbrunn-Kalkwerk II, sample No. 1A, x100.  
 Fig. 23: *Rosita fornicata* (PLUMMER); Ernstbrunn-Kalkwerk II, sample No. 4., x120.  
 Fig. 24: *Globotruncana linneiana* (D'ORBIGNY); Ernstbrunn-Kalkwerk II, sample No. 1A, x120.  
 Fig. 25: *Nodosaria* sp.; Ernstbrunn-Kalkwerk II, sample No. 4, x60.

Microphotographs by Ananda Gabašová, Laboratory of the Czech Geological Survey.



nearby sample No. A/1998 did not provide any calcareous nanofossils.

Species *Biantolithus sparsus* was recorded as an accessory in Kalkwerk II, sample No. 1A (see Plate 3, Figs. 37, 38). The presence of *B. sparsus* in an assemblage which otherwise is of the Upper Campanian age and where no contamination by the Upper Maastrichtian or Tertiary nanofossils was observed remains a mystery. *B. sparsus* is usually supposed to be a marker species for the base of the Tertiary. Even though this species was observed in the uppermost Maastrichtian sediments of the Outer Western Carpathians (ŠVÁBENICKÁ in BUBÍK et al., 1999 and SUMMESBERGER et al., 1999), no indications of its occurrence are known from stratigraphically older sediments by now.

## 5.2. Staatz Klippe

### 5.2.1. Calcareous Nanofossils

Samples provided mostly poorly preserved and markedly overgrown and/or dissolved calcareous nanofossils. Nanofossil associations are characterized by the dominance of species *Watznaueria barnesae* and relatively high abundance of *Rucinolithus* spp. and *Gartnerago obliquum* where overgrowth is evident. In contrast, specimens of genera *Prediscosphaera* and *Cribrosphaerella* occur rarely here and show strong dissolution. Furthermore, many nanofossil species which usually form assemblages of Coniacian and Santonian age were not recorded probably due to the specific paleoenvironmental conditions in the depositional area.

Marly sediments obtained from narrow dissolution fractures in the Ernstbrunn Limestone (samples Nos. 3/1994 and 1R/1998) yielded poorly preserved and markedly overgrown nanofossil association represented by species *Watznaueria barnesae* (quantitatively prevailing), *W. biporta*, *W. britannica*, *Rucinolithus hayi*, *R. wissei*, *Gartnerago obliquum*, rare *Kamptnerius magnificus*, *Reinhardtites anthophorus*, *Quadrum gartneri*, *Q. gartneri* ssp. 2 (sensu CRUX, 1982), *Eprolithus floralis*, *Eiffellithus eximius*, *E. gorkae*, *Retacapsa angustiforata*, *Chiastozygus litterarius*, *Prediscosphaera cretacea*, *P. columnata*, *Manivitella pemmatoidea*, *Staurolithites* cf. *imbricatus* which gives evidence for the Lower Santonian age, the upper part of the UC11c Subzone. It is interesting

that the solution-resistant species *Micula staurophora* was absent from both the above mentioned samples.

Sediments obtained from an about 0.5 m wide fissure (middle part of northern wall) in the limestones provided the following nanofossil associations (sample No. 3R/1998):

1. Markedly overgrown nanofossils of Santonian age where preservation of coccoliths is comparable with that mentioned in sample No. 3/1994): *Micula staurophora*, *Gartnerago obliquum*, *Reinhardtites anthophorus*, *Kamptnerius magnificus*, *Rucinolithus wissei*, *R. hayi*, *Cylindralithus biarcus*, *C. nudus* and others.

2. Calcareous nanofossils without marked overgrowths evidencing the following stratigraphical horizons:

- Association with species *Arkhangelskiella cymbiformis*, *Broinsonia parca constricta*, *Biscutum magnum*, *Calculites obscurus*, *Reinhardtites levis*, *Lithraphidites praequadratus* evidencing the Upper Campanian.

- Association with *Markalius apertus*, *M. astroporus*, *Cruciplacolithus primus*, *C. tenuis* and *Biantolithus hughesii* evidencing the Lower Danian.

- Association with *Helicosphaera carteri*, *Pontosphaera multipora*, *Reticulofenestra coenura*, *R. hillae*, *Dictyococcites bisectus*, *Coccolithus pelagicus*, *C. formosus* etc. evidencing the Oligocene – Lower Miocene age.

Although the above mentioned species form a negligible part of the nanofossil association, their presence needs further discussion: younger nanofossils here indicate either the Miocene age of sediments or the contamination of Cretaceous sediments by younger ones caused e. g. by solifluction. The Miocene age can be supported by the youngest species found here: *Helicosphaera carteri*. High proportion (over 90 %) of Upper Cretaceous reworked nanofossils is a typical phenomenon of the Miocene thanatocoenoses in the Alpine-Carpathian Foredeep (ŠVÁBENICKÁ & ČTYROKÁ, 1998). This observation is similar to BACHMAYER (1964) who described Cretaceous and Tertiary nanofossils from fissure filling which otherwise are Quaternary.

## 6. Conclusions

### Ernstbrunn Quarry

- Grey to black clay (Kalkwerk II, level 5, sample No. A/1998) provided the monogeneric nanofossil association of

## Plate 2: Calcareous Nanofossils from Ernstbrunn and Staatz


- Figs. 1, 2: *Broinsonia parca constricta* HATTNER et al.; Ernstbrunn-Kalkwerk II, sample No. 3.  
 Figs. 3, 4: *Arkhangelskiella specillata* VEKSHINA; Ernstbrunn-Kalkwerk II, sample No. 1.  
 Figs. 5, 6: *Arkhangelskiella cymbiformis* VEKSHINA; Ernstbrunn-Kalkwerk II, sample No. 1A.  
 Figs. 7, 8: *Dodekapodorhabdus noeliae* PERCH-NIELSEN; Ernstbrunn-Kalkwerk II, sample No. 3.  
 Figs. 9, 10: *Petrarhabdus copulatus* (DEFLANDRE) WIND and WISE; Ernstbrunn-Kalkwerk II, sample No. 3.  
 Figs. 11, 12: *Cylindralithus serratus* BRAMLETTE & MARTINI; Ernstbrunn-Kalkwerk II, sample No. 4.  
 Figs. 13, 14: *Eiffellithus eximius* (STOVER) PERCH-NIELSEN; Staatz Klippe, sample No. 1R/1998.  
 Figs. 15, 16: *Eiffellithus turriseiffelii* (DEFLANDRE) REINHARDT; Staatz Klippe, sample No. 1R/1998.  
 Figs. 17, 18: *Cylindralithus* cf. *nudus* BUKRY; Staatz Klippe, sample No. 3R/1998.  
 Figs. 19, 20: *Prediscosphaera columnata* (STOVER) PERCH-NIELSEN; Staatz Klippe, sample No. 1R/1998.  
 Figs. 21, 22: *Helicolithus trabeculatus* (GÓRKA) VERBEEK; Staatz Klippe, sample No. 3R/1998.  
 Figs. 23, 24: *Gartnerago obliquum* (STRADNER) NOËL; Staatz Klippe, sample No. 1R/1998.  
 Figs. 25, 26: *Zeugrhabdodus embergeri* (NOËL) PERCH-NIELSEN; Staatz Klippe, sample No. 1R/1998.  
 Figs. 27, 28: *Rucinolithus* cf. *wisei* THIERSTEIN; Staatz Klippe, sample No. 1R/1998.  
 Figs. 29, 30: *Rucinolithus wisei* THIERSTEIN; Staatz Klippe, sample No. 1R/1998.  
 Figs. 31, 32: *Rucinolithus* cf. *hayi* STOVER; Staatz Klippe, sample No. 1R/1998.  
 Figs. 33, 34: *Rucinolithus hayi* STOVER; Staatz Klippe, sample No. 1R/1998.  
 Figs. 35, 36: *Micula staurophora* (GARDET) STRADNER; Staatz Klippe, sample No. 3R/1998.  
 Figs. 37, 38: *Kamptnerius magnificus* DEFLANDRE; Staatz Klippe, sample No. 3R/1998.  
 Figs. 39, 40: *Quadrum gartneri* ssp. 2 sensu CRUX (1982); Staatz Klippe, sample No. 1R/1998.  
 Fig. 41: *Micula-Quadrum*; Staatz Klippe, sample No. 3R/1998.  
 Microphotographs by L. Švábenická, magnification 1,000x.



1




2



3




4



5



6



7



8



9




10



11




12



13



14



15




16



17




18



19




20



21



22



23



24




25




26



27




28




29




30




31




32



33



34



35



36



37



38



39



40



41

Tithonian age which is dominated by *Watznaueria barnesae*.

- Santonian is proved by nannofossil assemblage with *Rucinolithus* spp. and *Orastum campanensis* of the UC11c-UC12 zone interval which is supported by poorly preserved foraminifers of the Upper Santonian-Lower Campanian age with *Arenobulimina d'orbignyi* (REUSS) and *Stensioeina exsculpta* (REUSS).

- Upper Campanian is documented by foraminiferal assemblage of the Globotruncanita calcarata total range zone and by nannofossils of the UC15c-d<sup>TP</sup> Zone.

- Common occurrence of high- and low-latitude nannofossil species in the Upper Campanian sediments indicates that the depositional area was probably situated within the Tethyan/Mediterranean province with a strong influence from the Boreal area.

- Upper Campanian nannofossil associations are similar in their character to those of the same age from the Pálava Formation in the Ždánice Unit, Outer Western Carpathians.

#### Staatz Klippe

Two different nannofossil associations were observed:

1. Association with *Rucinolithus hayi*, *R. wissei* and *Reinhardtites anthophorus* which gives evidence for the UC11c Subzone of Lower Santonian age.

2. Mixture of nannofossil species of the Oligocene-Lower Miocene, Lower Danian, Upper Campanian and Santonian age which can be explained either by contamination of Cretaceous sediments by younger ones or by reworking older nannofossils into the Miocene sediments.

To conclude, nannofossil preservation (overgrowth) in sediments of locality Staatz Klippe is very similar to another one observed in the Santonian sediments of the Ernstbrunn Quarry.

### Acknowledgement

Thanks are due to Mr. Manfred Schiller from the Geological Survey of Austria for processing the foraminiferal samples, to Mrs. Alena Tichá for nannofossil samples, to Mrs. Ananda Gabašová for scanning microphotographs and to Mrs. Nadežda Hrdličková for taking the photographs (all the ladies from the Czech Geological Survey). For drawings thanks are due to Leo Leitner (Krems/Lower Austria) and Jacek Ruthner (Geological Survey of Austria). For help with the field work we want to thank Dr. Herbert Stradner (Klosterneuburg/Lower Austria) and for permission to work in Ernstbrunn quarry we want to

thank Ing. Josef Gayer from the "Ernstbrunner Kalktechnik – Kalkgewerkschaft in Ernstbrunn Ges.m.b.H". For helping with the English language: K. Histon (Geological Survey of Austria).

Nannofossil study is a contribution to the project No. 205/97/0687 "Low- and high-latitude nannofossils and their biostratigraphic significance (Campanian and Maastrichtian, Western Carpathians)" supported by the Grant Agency of the Czech Republic.


### References

- AL-SHAIBANI, S. K. (1971): An Upper Maastrichtian Foraminiferal Fauna from Dörfles, Lower Austria. – Jahrb. Geol.-B.A., Sdbd. 17, 105–119, Wien.
- BACHMAYER, F. (1940): Beiträge zur Kenntnis der Tithonfauna aus dem Raume von Ernstbrunn (Niederdonau). – Unpublished PhD. Thesis, Univ. Vienna, 73 p., Wien.
- BACHMAYER, F. (1964): Untersuchung einer Kluffüllung im Steinbruch Staatz (Kautendorf), nördliches Niederösterreich. – Ann. Naturhistor. Mus. Wien, 67, 181–187, Wien.
- BUBIK, M., BAK, M. & ŠVÁBENICKÁ, L. (1999): Biostratigraphy of the Maastrichtian to Paleocene distal flysch sediments of the Rača Unit in the Uzgruň section (Magura Group of nappes, Czech Republic). – Geologica Carpathica, 50, 1, 33–48, Bratislava.
- BULLINGER, E. (1997): Die Bedeutung des Ernstbrunner Kalkes für die Industrie. – ÖGG Exk. Führer 17, 37–38, Wien.
- BURNETT, J. (1998): Upper Cretaceous. In: BOWN, P. R. (Ed.): Calcareous Nannofossil Biostratigraphy. – Cambridge University Press, 132–199, Cambridge.
- CRUX, J. A. (1982): Upper Cretaceous (Cenomanian to Campanian) calcareous nannofossils. In: LORD, A. R. (Ed.): A Stratigraphical Index of Calcareous Nannofossils. – British Micropalaeontological Society Series, Ellis Horwood Limited, 81–135, Chichester.
- ELIÁŠ, M. (1992): Sedimentology of the Kientnice Formation and the Ernstbrunn Limestone (Ždánice–Subsilesian unit of the Outer West Carpathians). – Věst. Čes. geol. Úst., 67, 3, 179–193, 6 figs., 2 pls., Praha.
- ELIÁŠ, M., SCHNABEL, W. & STRÁNIK, Z. (1990): Comparison of the Flysch Zone of the Eastern Alps and the Western Carpathians based on recent observations. – In: MINAŘIKOVÁ & LOBITZER (Eds.): Thirty Years of Geological Cooperation between Austria and Czechoslovakia, 37–46, Vienna/Praha.
- FUCHS, W. (1980): Das Jungalpidikum (Die geologische Entwicklung vom Obereozän bis in die Gegenwart), 49–55, Abb.14.– In: OBERHAUSER, R. (Ed.): Der geologische Aufbau Österreichs, 700 S., Wien – New York.
- GASINSKI, M., JUGOWIEC, M. & ŚLACZKA, A. (1999): Late Cretaceous foraminiferids and calcareous nannoplankton from the Węglówka Marls (Subsilesian Unit, Outer Carpathians, Poland). – Geologica Carpathica, 50, 1, 63–73, Bratislava.
- GRILL, R. (1953): Der Flysch, die Waschbergzone und das

### Plate 3: Calcareous Nannofossils from Ernstbrunn and Staatz

- Figs. 1, 2: *Ahmuellerella regularis* (GÓRKA) REINHARDT; Ernstbrunn-Kalkwerk II, sample No. 3.
- Fig. 3: *Ahmuellerella octoradiata* (GÓRKA) REINHARDT; Ernstbrunn-Kalkwerk II, sample No. 3.
- Fig. 4: *Corollithion exiguum* STRADNER; Ernstbrunn-Kalkwerk II, sample No. 4.
- Figs. 5, 6: *Orastrum campanensis* (CEPEK) WIND & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 7, 8: *Reinhardtites anthophorus* (DEFLANDRE) PERCH-NIELSEN; Ernstbrunn-Kalkwerk II, sample No. 4.
- Figs. 9, 10: *Reinhardtites levis* PRINS & SISSINGH; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 11, 12: *Cretarhabdus conicus* BRAMLETTE & MARTINI; Ernstbrunn-Kalkwerk II, sample No. 4.
- Figs. 13, 14: *Biscutum magnum* WIND & WISE; Ernstbrunn-Kalkwerk II, sample No. 4.
- Figs. 15, 16: *Biscutum coronum* WIND & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 17, 18: *Biscutum ellipticum* (GÓRKA) GRÜN; Staatz Klippe, sample No. 3R/1998.
- Figs. 19, 20: *Angulofenestrelithus snyderi* BUKRY; Ernstbrunn-Kalkwerk II, sample No. 1A.
- Figs. 21, 22: *Staurolithes imbricatus* (GARTNER) BURNETT; Staatz Klippe, sample No. 3R/1998.
- Figs. 23, 24: *Tortolithus cf. hallii* (BUKRY) CRUX; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 25, 26: *Uniplanarius sissinghii* PERCH-NIELSEN; Ernstbrunn-Kalkwerk II, sample No. 1A.
- Figs. 27, 28: *Uniplanarius trifidus* (STRADNER) HATTNER & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 29, 30: *Ceratolithoides sesquipedalis* BURNETT; Ernstbrunn-Kalkwerk II, sample No. 1A.
- Figs. 31, 32: *Monomarginatus quaternarius* WIND & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 33, 34: *Neocrepidolithus watkinsii* POSPICAL & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 35, 36: *Neocrepidolithus watkinsii* POSPICAL & WISE; Ernstbrunn-Kalkwerk II, sample No. 3.
- Figs. 37, 38: *Biantholithus sparsus* BRAMLETTE & MARTINI (?contamination from Tertiary deposits or the autochthonous component of the Upper Campanian association); Ernstbrunn-Kalkwerk II, sample No. 1A.
- Figs. 39, 40: *Kamptnerius magnificus* DEFLANDRE; Ernstbrunn-Kalkwerk II, sample No. 1A.
- Microphotographs by L. Švábenická, magnification 1,000x.





- Jungtertiär um Ernstbrunn (Niederösterreich). – Jahrb. Geol. B.-A., 96, 65–116, Wien.
- GRILL, R. (1968): Erläuterungen zur Geologischen Karte des nordöstlichen Weinviertels und zu Blatt Gänserndorf. – 155 p., Geol. B.-A., Wien.
- HANZLÍKOVÁ, E. (1972): Carpathian Upper Cretaceous Foraminifera of Moravia (Turonian-Maastrichtian). – Rozpravy Ústř. Úst. geol. 39, 1–160, Praha.
- HOFMANN, T. (1990): Der Ernstbrunner Kalk im Raum Dörfles (Niederösterreich) Mikrofazies und Kalkalgen. – Unpubl. Diplom, Univ. Vienna, 164 p., Wien.
- ROBASZYNSKI, F. & CARON, M. (1995): Foraminifères planctoniques du Crétacé: commentaire de la zonation Europe – Méditerranée. – Bull. Soc. Géol. France, 166, 6, 681–692, Orleans.
- SAUER, R., SEIFERT, P. & WESSELY, G. (1992) Guidebook to Excursions in the Vienna Basin and the Adjacent Alpine – Carpathian Thrustbelt in Austria. – Mitt. Österr. Geol. Ges., 85, 239 p., 200 figs., Wien.
- STEININGER, F. F., WESSELY, G., RÖGL, F. & WAGNER, L. (1986): Tertiary sedimentary history and tectonic evolution of the eastern Alpine Foredeep. – Giorn. di Geol., 3, 48/1–2, 285–297, Bologna.
- STRÁNÍK, Z., BUBÍK, M., ČECH, S. & ŠVÁBENICKÁ, L. (1996): The Upper Cretaceous in South Moravia. – Věst. Čes. geol. Úst., 71, 1, 1–30, Praha.
- SUMMESBERGER, H., ŠVÁBENICKÁ, L., ČECH, S., HRADECKÁ, L. & HOFMANN, T. (1999): New palaeontological Data on the Klement and Pálava Formations (Late Cretaceous) in Austria (Waschberg-Zdánice Unit). – Ann. Naturhistor. Mus. Wien, 100, 39–79, 2 figs., 3 tab., 6 pls., Wien.
- ŠVÁBENICKÁ, L. (1998): Occurrence of *Petrarhabdus copulatus* (Deflandre) Wind and Wise in the Outer Western Carpathians. – Věst. Čes. geol. Úst., 73, 2, 89–97, Praha.
- ŠVÁBENICKÁ, L. & ČTYROKÁ, J. (1998): Foraminifera and nanofossil biostratigraphic correlation of the Karpatian and Lower Badenian sediments in the Carpathian Foredeep of Moravia (Preliminary results). – Zprávy o geologických výzkumech v roce 1997, 166–170, Praha.
- WATKINS, D. K. (1992): Upper Cretaceous nanofossils from Leg 120, Kerguelen Plateau, Southern Ocean. – Proceedings of the ODP, Scientific Results, 120, 343–370, Washington.
- WATKINS, D. K., WISE, S. W., POSPÍCHAL, J. J. & CRUX, J. (1996): Upper Cretaceous calcareous nanofossil biostratigraphy and paleoceanography of the Southern Ocean. In: MOGUILVSKY, A. & WHATLEY, R.: Microfossils and Oceanic Environments. – University of Wales, Aberystwyth Press, 355–381, Aberystwyth.
- WIND, F. H. (1979): Maestrichtian-Campanian nanofloral provinces of the southern Atlantic and Indian Oceans. In: TALWANI, M. HAY, W. W. & RYAN, W. B. F. (Eds.): Deep drilling results in the Atlantic Ocean: Continental Margins and Paleoenvironment. – Am. Geophys. Union, 123–137, Washington.
- Ceratolithoides sesquipedalis* BURNETT, 1998  
*Chiastozygus litterarius* (GÓRKA, 1957) MANIVIT, 1971  
*Corollithion exiguum* STRADNER, 1961  
*Creterhabdus conicus* BRAMLETTE & MARTINI, 1964  
*Cyclagelosphaera magerelii* NOËL, 1965  
*Cylindralithus biarcus* BUKRY, 1969  
*Cylindralithus nudus* BUKRY, 1969  
*Cylindralithus serratus* BRAMLETTE & MARTINI, 1964  
*Dodekapodorhabdus noeliae* PERCH-NIELSEN, 1968  
*Eiffellithus eximius* (STOVER, 1966) PERCH-NIELSEN, 1968  
*Eiffellithus gorkae* REINHARDT, 1965  
*Eiffellithus turiseiffelii* (DEFLANDRE in DEFLANDRE & FERT, 1954) REINHARDT, 1965  
*Eprolithus floralis* (STRADNER, 1962) STOVER, 1966  
*Gartnerago obliquum* (STRADNER, 1963) NOËL, 1970  
*Grantarhabdus coronadventis* (REINHARDT, 1966) GRÜN in GRÜN & ALLEMANN, 1975  
*Helicolithus trabeculatus* (GÓRKA, 1957) VERBEEK, 1977  
*Kamptnerius magnificus* DEFLANDRE, 1959  
*Lithraphidites praequadratus* ROTH, 1978  
*Manivittella pemmatoidea* (DEFLANDRE in MANIVIT, 1965) THIERSTEIN, 1971  
*Markalius inversus* (DEFLANDRE in DEFLANDRE & FERT, 1954) BRAMLETTE & MARTINI, 1964  
*Micula staurophora* (GARDET, 1955) STRADNER, 1963  
*Micula swastica* STRADNER & STEINMETZ, 1984  
*Monomarginatus quaternarius* WIND & WISE in WISE & WIND, 1977  
*Neocrepidolithus watkinsii* POSPÍCHAL & WISE, 1990  
*Orastrum campanensis* (CEPEK, 1970) WIND & WISE in WISE & WIND, 1977  
*Petrarhabdus copulatus* (DEFLANDRE, 1959) WIND & WISE in WISE, 1983  
*Prediscosphaera columnata* (STOVER, 1966) PERCH-NIELSEN, 1984  
*Prediscosphaera cretacea* (ARKHANGELSKY, 1912) GARTNER, 1968  
*Prediscosphaera grandis* PERCH-NIELSEN, 1979  
*Prediscosphaera stoveri* (PERCH-NIELSEN, 1968) SHAFIK & STRADNER, 1971  
*Quadrum gartneri* PRINS & PERCH-NIELSEN in MANIVIT et al., 1977  
*Reinhardtites anthophorus* (DEFLANDRE, 1959) PERCH-NIELSEN, 1968  
*Reinhardtites levis* PRINS & SISSINGH in SISSINGH, 1977  
*Retacapsa angustiforata* BLACK, 1971  
*Rucinolithus hayi* STOVER, 1966  
*Rucinolithus wissei* THIERSTEIN, 1971  
*Staurolithites achenus* (BUKRY, 1969) BURNETT, 1998  
*Staurolithites imbricatus* (GARTNER, 1968) BURNETT, 1998  
*Tortololithus hallii* (BUKRY, 1969) CRUX in CRUX et al., 1982  
*Uniplanarius sissinghii* PERCH-NIELSEN, 1986  
*Uniplanarius trifidus* (STRADNER in STRADNER & PAPP, 1961)  
*Watznaueria barnesae* (BLACK, 1959) PERCH-NIELSEN, 1968  
*Watznaueria biporta* BUKRY, 1969  
*Watznaueria britannica* (STRADNER, 1963) REINHARDT, 1964  
*Watznaueria manivittae* BUKRY, 1963  
*Zeughrabdotos embergeri* (NOËL, 1958) PERCH-NIELSEN, 1984

## Appendix

### List of calcareous nanofossil taxa mentioned in this study, in the alphabetical order of generic epithets:

#### Cretaceous taxa

- Ahmullerella octoradiata* (GÓRKA, 1957) REINHARDT, 1966  
*Ahmullerella regularis* (GÓRKA, 1957) REINHARDT & GÓRKA, 1967  
*Angulofenestrellithus snyderi* BUKRY, 1969  
*Arkhangelskiella cymbiformis* VEKSHINA, 1959  
*Arkhangelskiella specillata* VEKSHINA, 1959  
*Biscutum coronum* WIND & WISE in WISE & WIND, 1977  
*Biscutum ellipticum* (GÓRKA, 1957) GRÜN in GRÜN & ALLEMANN, 1975  
*Biscutum magnum* WIND & WISE in WISE & WIND, 1977  
*Broinsonia parca constricta* HATTNER et al., 1980  
*Calculites obscurus* (DEFLANDRE, 1959) PRINS & SISSINGH in SISSINGH, 1977  
*Ceratolithoides aculeus* (STRADNER, 1961) PRINS & SISSINGH in SISSINGH, 1977  
*Ceratolithoides quasiarquatus* BURNETT, 1998

#### Tertiary taxa

- Biantholithus hughesii* VAROL, 1989  
*Biantholithus sparsus* BRAMLETTE & MARTINI, 1964  
*Coccolithus formosus* (KAMPTNER, 1963) WISE, 1973  
*Coccolithus pelagicus* (WALLICH, 1871) SCHILLER, 1930  
*Cruciplacolithus primus* PERCH-NIELSEN, 1977  
*Cruciplacolithus tenuis* (STRADNER, 1961) HAY & MOHLER in HAY et al., 1967  
*Dictyococcites bisectus* (HAY et al., 1966) BUKRY & PERCIVAL, 1971  
*Helicosphaera carteri* (WALLICH, 1877) KAMPTNER, 1954  
*Isthmolithus recurvus* DEFLANDRE in DEFLANDRE & FERT, 1954  
*Markalius apertus* PERCH-NIELSEN, 1979  
*Markalius astroporus* (STRADNER, 1963) HAY & NOHLER in HAY et al., 1967  
*Pontosphaera multipora* (KAMPTNER, 1948) ROTH, 1970  
*Reticulofenestra coenura* (REINHARDT, 1966) ROTH, 1970  
*Reticulofenestra hillae* BUKRY & PERCIVAL, 1971  
*Sullivania danica* (BROTZEN, 1959) VAROL, 1992

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Abhandlungen der Geologischen Bundesanstalt in Wien](#)

Jahr/Year: 1999

Band/Volume: [56\\_2](#)

Autor(en)/Author(s): Hofmann Thomas, Svábenická Lilian, Hradecka Lenka

Artikel/Article: [Biostratigraphy of Fissure Fillings in the Ernstbrunn Limestone of the Waschberg Zone \(Lower Austria\) 465-474](#)