



**PRELIMINARY REPORT ON THE AUCHENORRHYNCHA
(HEMIPTERA) FAUNA OF KAZDAĞI NATIONAL PARK WITH TWO
NEW RECORDS FOR TURKEY***

Emine DEMİR

Gazi University, Faculty of Science & Arts, Department of Biology,
06500 Teknikokullar Ankara, Turkey. E-mail: eminedemird@yahoo.com

Abstract - Auchenorrhyncha specimens collected in the years 2004 and 2005 in the Kazdağı (Ida Mountain) National Park in north-west Turkey were examined. 40 species belonging to 9 families were found: 1 species belongs to the family Cixiidae, 2 to Delphacidae, 2 to Tettigometridae, 1 to Meenoplidae, 3 to Issidae, 1 to Flatidae, 7 to Cercopidae, 2 to Membracidae, and 21 species to Cicadellidae. Distribution of these species in Turkey and their host plants are given along with their locality records. 31 species records are new to Balıkesir, and *Bubastia (s.str.) ludviki* Dlabola, 1979 and *Iassus mirabilis* Orosz, 1979 are new records for Turkish fauna.

KEY WORDS: Hemiptera, Auchenorrhyncha, fauna, Kazdağı National Park, Turkey.

Izveček – PREDHODNO POROČILO O FAVNI ŠKRŽATKOV (HEMIPTERA: AUCHENORRHYNCHA) V NARODNEM PARKU KAZDAĞI Z DVEMA NOVIMA NAJDBAMA ZA TURČIJO

Pregledali smo primerke škržatkov, zbrane v letih 2004 in 2005 v Narodnem parku Kazdağı (gora Ida) v severozahodni Turčiji. Našli smo 40 vrst, pripadajočih 9 družinam: 1 vrsta pripada družini Cixiidae, 2 družini Delphacidae, 2 družini Tettigometridae, 1 družini Meenoplidae, 3 družini Issidae, 1 družini Flatidae, 7 družini Cercopidae, 2 družini Membracidae in 21 vrst družini Cicadellidae. Razširjenost teh vrst v Turčiji in njihove hranilne rastline so navedeni poleg podatkov o najdiščih. Podatki o 31 vrstah so novi za Balıkesir, vrsti *Bubastia (s.str.) ludviki* Dlabola, 1979, in *Iassus mirabilis* Orosz, 1979, pa sta prvič zabeleženi v turški favni.

KLJUČNE BESEDE: Hemiptera, Auchenorrhyncha, favna, Narodni park Kazdağı, Turčija.

* This study was supported by TUBITAK (The Project number: TB YDBAG-1034110). The present work includes only Auchenorrhyncha part of the project.

Introduction

Turkey's first faunistic records of Auchenorrhyncha were given by Fahringer (1922), Haupt (1930), Kerville (1939), Zachvatkin (1937, 1946), Dlabola (1957, 1971a, 1971b, 1974, 1981, 1982, 1983, 1985a, 1985b, 1986), Linnavuori (1965), Asche (1982), Remane and Hoch (1985), Dworakowska (1970, 1972, 1982) Kalkandelen (1974, 1980, 1985, 1987, 1988, 1989a, 1989b, 1990, 1993, 1994, 2000) and Kartal (1981, 1983, 1985a, 1985b, 1986, 1987, 1988). Lodos and Kalkandelen (1980-1988) compiled the faunistic records of the studies until 1988 and listed 725 species in a preliminary checklist. Later, the studies of Abdul-Nour (1988), Meyer-Arndt (1988, 1991), Kartal and Zeybekoğlu (1992, 1994a, 1994b, 1997), Zeybekoğlu (1996, 1998), Zeybekoğlu and Bulut (2000), Kartal, Zeybekoğlu and Dursun (2001), Başpınar and Uygun (1991a, 1991b, 1991c, 1992a, 1992b), Boulard (1979, 1993), Güçlü and Özbek (1995, 1996), Gnezdilov (2002), Emeljanov (2003), Demir (1998, 2004a, 2004b, 2005a, 2005b, 2006) presented records of 48 additional species, raising the total species number for the country to 773.

Province of Kazdağı National Park (in Balıkesir country), which was selected as the study area, is located in southwest Turkey, between 26° 30' - 27° 00' eastern longitude and 39° 30' - 39° 45' northern latitude. The elevation ranges between 100 m and 1774 m. 26 Auchenorrhyncha species were known in Balıkesir area prior to our study, a compilation of records in the studies of the authors mentioned above.

Material and methods

Field studies were carried out in the years 2004 and 2005. 257 adult Auchenorrhyncha specimens were collected and analyzed. They were collected by sweeping the herbaceous plants with a wood shaft net. The host plants of species collected from only one plant species by wood shaft net were determined. The material is kept dry in the author's collection.

Results

Altogether 40 species were found. Along with their distribution in Turkey and the plants found to be their hosts, the examined material of these species is listed below.

Family: CIXIIDAE SPINOLA, 1839

Hyalesthes obsoletus Signoret, 1866

Material examined: Balıkesir, Kazdağı National Park, Zeytinli, 150 m, 06.06.2005 7♂9♀. On *Vitex agnus-castus*. **Distribution in Turkey:** Adıyaman, Afyon, Ağrı, Ankara, Antalya, Aydın, Balıkesir, Bolu, Burdur, Bursa, Çankırı, Çanakkale, Çorum, Diyarbakır, Düzce, Elazığ, Erzincan, Erzurum, Eskişehir, Gaziantep, Giresun, Hakkari, Iğdır, Isparta, İçel, İstanbul, Kahramanmaraş, Konya, Mardin, Muğla, Rize, Sakarya, Sinop, Sivas, Tokat, Trabzon, Urfa, Van (Fahringer, 1922; Lodos & Kalkandelen, 1980; Hoch & Remane, 1985; Kalkandelen, 2000).

Family: DELPHACIDAE LEACH, 1815

***Kelisia ribauti* Wagner, 1938**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 9♂3♀. On *Juncus*. **Distribution in Turkey:** Ankara, Bitlis, Diyarbakır, Erzurum, Hakkari, Siirt, Van (Lodos & Kalkandelen, 1980b; Güçlü, 1996).

***Conomelus odryssi* Dlabola, 1965**

Material examined: Balıkesir, Kazdağı National Park, Tozlu place, 1200 m, 28.08.2004, 3♂29♀; Tozlu place, 1200 m, 05.06.2005, 2♂2♀. On *Juncus*. **Distribution in Turkey:** Konya (Remane & Asche, 1979).

Family: MEENOPLIDAE FIEBER, 1872

***Meenoplus albosignatus* Fieber, 1866**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♂. On *Quercus*. **Distribution in Turkey:** Adıyaman, Ankara, Bolu, Hakkari, Mardin, Muş (Linnavuori, 1965; Lodos & Kalkandelen, 1980c; Demir, 2006).

Family: TETTIGOMETRIDAE GERMAR, 1821

***Tettigometra (s.str.) sulphurea* Mulsant et Rey, 1855**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 1♀. It has been collected from weeds in a *Pinus brutia* forest. **Distribution in Turkey:** Artvin, Aydın, Bilecik, Bursa, Diyarbakır, Elazığ, İzmir, Kütahya, Nevşehir, Sakarya, Urfa, Uşak, Van (Lodos & Kalkandelen, 1980c; Demir, 2006).

***Tettigometra (Metroplaca) baranii* (Signoret, 1866)**

Material examined: Balıkesir, Kazdağı National Park, Tozlu place, 1100 m, 24.07.2004, 1♂3♀ YD leg.; Top of Kazdağı, 1700 m, 28.08.2004, 1♂1♀. **Distribution in Turkey:** Ankara, Mardin (Lodos & Kalkandelen, 1980c; Demir, 2006).

Family: ISSIDAE SPINOLA, 1839

***Peltonotellus punctifrons* Horváth 1895**

Material examined: Balıkesir, Kazdağı National Park, Top of Kazdağı, 1700 m, 28.08.2004, 1♂. **Distribution in Turkey:** Ankara, Kırşehir, Yozgat (Dlabola, 1957; Kartal, 1985).

***Bubastia (s.str.) ludviki* Dlabola, 1979**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♂. **Distribution in Turkey:** This is the first record of this species for Turkey.

***Agalmatium bilobum* (Fieber, 1877)**

Material examined: Balıkesir, Kazdağı National Park, Zeytinli, 150 m, 06.06.2005, 1♂1♀. **Distribution in Turkey:** Adana, Adıyaman Afyon, Ankara,

Aydın, Balıkesir, Bilecik, Burdur, Bursa, Çanakkale, Çorum, Denizli, Eskişehir, Gaziantep, Hatay, İzmir, Gümüşhane, Kırklareli, Kırşehir, Kütahya, Malatya, Manisa, Muğla, Sakarya, Sivas, Tekirdağ, Tokat, Uşak, Yozgat (Linnvuori, 1965; Lodos & Kalkandelen, 1981a; Kartal, 1985).

Family: FLATIDAE SPINOLA, 1839

***Phantia subquadrata* (Herrich-Schaeffer, 1838)**

Material examined: Balıkesir, Kazdağı National Park, Zeytinli stream, 150 m, 28.08.2004, 1♀. On *Genista*. **Distribution in Turkey:** Ankara, Aydın, Bilecik, Denizli, Diyarbakır, Edirne, Isparta, İzmir, Manisa, Mardin, Muğla (Lodos & Kalkandelen, 1981b; Demir, 2004b).

Family: CERCOPIDAE LEACH, 1815

***Cercopis intermedia* Kirschbaum, 1868**

Material examined: Balıkesir, Kazdağı National Park, South side, 1200 m, 03.07.2004, 1♀; Tozlu place, 1480 m, 03.07.2004, 1♂ Y.Durmuş leg. **Distribution in Turkey:** Adıyaman, Amasya, Artvin, Ankara, Bitlis, Çanakkale, Diyarbakır, Elazığ, Eskişehir, Gaziantep, Giresun, Hakkari, Hatay, Isparta, İzmir, Kahramanmaraş, Kayseri, Kırklareli, Konya, Kütahya, Mardin, Niğde, Rize, Samsun, Siirt, Trabzon, Urfa, Uşak (Lodos & Kalkandelen, 1981c; Kartal et. al., 1994; Demir, 2006).

***Cercopis vulnerata* Rossi, 1807**

Material examined: Balıkesir, Kazdağı National Park, Tozlu place, 24.07.2004, 2♂1♀; Tozlu place, 03.07.2004, 2♂1♀ Y.Durmuş leg. **Distribution in Turkey:** Bursa, Kocaeli, Sakarya (Lodos & Kalkandelen, 1981c).

***Triecphorella geniculata* (Horvath, 1881)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 1♀. **Distribution in Turkey:** Adana, Kahramanmaraş, Konya, Samsun (Fahringer, 1922; Kartal et. al., 1994).

***Neophilaenus campestris* (Fallén, 1805)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 1♂. **Distribution in Turkey:** Adana Hatay, Adıyaman Ankara, Artvin, Bitlis, Bursa, Çanakkale, Eskişehir, Giresun, İzmir, Kırklareli, Kütahya, Manisa, Muğla, Nevşehir, Samsun, Sinop, Siirt, Trabzon, Van (Linnvuori, 1965; Lodos & Kalkandelen, 1981c; Demir, 2006).

***Neophilaenus minor* (Kirschbaum, 1868)**

Material examined: Balıkesir, Kazdağı National Park, Top of Kazdağı, 1700 m, 28.08.2004, 3♂5♀. **Distribution in Turkey:** Ankara, Konya (Dlabola, 1957; Demir, 2004b, 2006).

***Aphrophora alni* (Fallén, 1805)**

Material examined: Balıkesir, Kazdaği National Park, Tozlu place, 24.07.2004, 1♂; Tozlu place, 23.07.2004, 1♀ Y.Durmuş leg. **Distribution in Turkey:** Afyon, Ankara, Artvin, Aydın, Balıkesir, Bitlis, Bolu, Çanakkale, Çorum, Diyarbakır, Erzincan, Erzurum, Giresun, İstanbul, İzmir, Kayseri, Kırklareli, Konya, Kütahya, Mardin, Muğla, Manisa, Ordu, Rize, Samsun, Sinop, Tekirdağ, Trabzon, Yozga (Linnavuori, 1965; Lodos & Kalkandelen, 1981c; Kartal et. al., 1994; Demir, 2004b, 2006).

***Philaenus spumarius* (Linnaeus, 1758)**

Material examined: Balıkesir, Kazdaği National Park, Tozlu place, 1480 m, 03.07.2004, 15♂20♀ Y.Durmuş leg.; Camp field, 700 m, 06.06.2005, 1♀; Tozlu place, 23.07.2004, 2♂2♀; Tozlu place, 24.07.2004, 3♂6♀; Tozlu place, 1480 m, 03.07.2004, 6♂4♀. **Distribution in Turkey:** Ağrı, Amasya, Ankara, Artvin, Aydın, Balıkesir, Bilecik, Bitlis, Bolu, Bursa, Çanakkale, Elazığ, Erzincan, Erzurum, Giresun, Gümüşhane, Hakkari, İzmir, Kars, Kırklareli, Kocaeli, Kütahya, Malatya, Manisa, Mardin, Muğla, Ordu, Rize, Samsun, Siirt, Sinop, Tekirdağ, Trabzon, Tokat, Van (Linnavuori, 1965; Lodos & Kalkandelen, 1981c; Kartal et. al., 1994; Demir, 2004b, 2006).

Family: MEMBRACIDAE RAFINESQUE, 1815***Gargara genistae* (Fabricius, 1775)**

Material examined: Balıkesir, Kazdaği National Park, Top of Kazdaği, 1700 m, 28.08.2004, 1♀. **Distribution in Turkey:** Ankara, Aydın, Balıkesir, Çorum, Erzurum, İzmir, Muğla (Lodos & Kalkandelen, 1981c; Demir, 2006).

***Centrotus cornutus* (Linnaeus, 1758)**

Material examined: Balıkesir, Kazdaği National Park, Tozlu place, 1200 m, 28.08.2004, 1♀; Tozlu place, 1200 m, 03.07.2004, 1♂ Y.Durmuş leg. **Distribution in Turkey:** Artvin, Aydın, Bilecik, Bursa, Çorum, Denizli, Erzincan, Edirne, Giresun, İzmir, Kocaeli, Kütahya, Kırklareli, Ordu, Sakarya, Tekirdağ (Lodos & Kalkandelen, 1981c; Demir, 2006).

Family: CICADELLIDAE LATREILLE, 1825***Iassus mirabilis* Orosz, 1979**

Material examined: Balıkesir, Kazdaği National Park, Camp field, 800 m, 29.08.2004, 1♂1♀. On *Quercus*. **Distribution in Turkey:** This is the first record of this species for Turkey.

***Eupelix cuspidata* (Fabricius, 1779)**

Material examined: Balıkesir, Kazdaği National Park, Camp field, 1000 m, 04.06.2005, 1♂. **Distribution in Turkey:** Adıyaman, Afyon, Ankara, Artvin, Çanakkale, Diyarbakır, Konya, Malatya, Mardin, Niğde, Urfa (Lodos & Kalkandelen, 1982b; Demir, 2006).

***Aphrodes makarovi* Zachvatkin, 1948**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 700 m, 06.06.2005, 3♂. **Distribution in Turkey:** Adana, Ankara (Başpınar & Uygun, 1991; Demir, 2006).

***Alebra albostriella* (Fallén, 1826)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♂1♀. On *Quercus*. **Distribution in Turkey:** Ankara, Artvin, Bolu, Çorum, Hatay, İzmir, Kayseri, Kocaeli, Ordu (Linnavuori, 1965; Lodos & Kalkandelen, 1983b; Demir, 2006).

***Micantulina (Mulsantina) stigmatipennis* (Mulsant et Rey, 1855)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 1♂2♀. On *Verbascum*. **Distribution in Turkey:** Ankara, Burdur, Isparta, İstanbul, İzmir, Manisa (Lodos & Kalkandelen, 1983b; Demir, 2006).

***Eupteryx gyaurdagicus* Dlabola, 1957**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 16♂14♀. On grass with *Mentha*. **Distribution in Turkey:** Ankara, Bolu, Isparta, İzmir, Manisa, Nevşehir, Sinop (Lodos & Kalkandelen, 1984b; Kartal, 1983; Demir, 2006).

***Eupteryx melissae* Curtis, 1837**

Material examined: Balıkesir, Kazdağı National Park, Zeytinli stream, 150 m, 28.08.2004, 4♂. On grass with *Mentha*. **Distribution in Turkey:** Ankara, Kastamonu, Sinop (Lodos & Kalkandelen, 1984b).

***Grypotes puncticollis* (Herrich-Schaeffer, 1834)**

Material examined: Balıkesir, Kazdağı National Park, Top of Kazdağı, 1700 m, 28.08.2004, 1♂; Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♀. On *Pinus nigra*. **Distribution in Turkey:** Amasya, Ankara, Çankırı, Ordu, Samsun, Sinop (Kalkandelen, 1974; Lodos & Kalkandelen, 1985; Zeybekoğlu, 1994; Zeybekoğlu, 1998).

***Balclutha rhenana* Wagner, 1939**

Material examined: Balıkesir, Kazdağı National Park, Zeytinli, 150 m, 06.06.2005, 2♂1♀. **Distribution in Turkey:** Erzurum (Lodos & Kalkandelen, 1985b).

***Platymetopius henribauti* Dlabola, 1961**

Material examined: Balıkesir, Kazdağı National Park, Zeytinli, 150 m, 06.06.2005, 1♂; Balıkesir, Kazdağı National Park, Zeytinli stream, 150 m, 28.08.2004, 1♀. **Distribution in Turkey:** Ankara, Konya (Kalkandelen, 1974; Lodos & Kalkandelen, 1986b; Demir, 2006).

***Anoplotettix fuscovenosus* (Ferrari, 1882)**

Material examined: Balıkesir, Kazdağı National Park , Camp field, 1000 m, 04.06.2005, 1♂. **Distribution in Turkey:** Ankara, Bursa, Çorum, Edirne, İzmir, Kırklareli, Manisa, Rize, Sakarya, Samsun, Sinop, Tokat (Lodos & Kalkandelen, 1986b; Kartal & Zeybekoğlu, 1991; Zeybekoğlu, 1998; Demir, 2006).

***Selenocephalus pallidus* Kirschbaum, 1868**

Material examined: Balıkesir, Kazdağı National Park , Zeytinli stream, 150 m, 28.08.2004, 1♀. On *Cistus creticus*. **Distribution in Turkey:** Adana, Ankara, Antalya, Aydın, Bolu, Burdur, Bursa, Çanakkale, Çankırı, Çorum, Denizli, Isparta, İzmir, Kayseri, Kırşehir, Konya, Manisa, Samsun, Sinop, Tokat, Yozgat, (Linnavuori, 1965; Kalkandelen, 1974; Lodos & Kalkandelen, 1986c; Kartal & Zeybekoğlu, 1991; Demir, 2006).

***Hardya anatolica* Zachvatkin, 1946**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 1♀. **Distribution in Turkey:** Adıyaman, Ankara, Antalya, Bolu, Elazığ, Erzincan, Isparta, İzmir, Karaman, Kayseri, Kırşehir, Nevşehir, Ordu (Zachvatkin, 1946; Linnavuori, 1965; Kalkandelen, 1974; Lodos & Kalkandelen, 1987; Demir, 2004b, 2006).

***Cicadula (Henriana) lineatopunctata* (Matsumura, 1908)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 3♂. **Distribution in Turkey:** Ankara, Aydın, Bolu, Bursa, Çankırı, Diyarbakır, Düzce, Erzincan, İzmir, Mardin, Sakarya, Sinop, Urfa (Kalkandelen, 1974; Lodos & Kalkandelen, 1987).

***Handianus procerus* (Herrich-Schaeffer, 1835)**

Material examined: Balıkesir, Kazdağı National Park , Top of Kazdağı, 1700 m, 28.08.2004, 1♂3♀. **Distribution in Turkey:** Afyon, Ağrı, Ankara, Antalya, Bitlis, Erzurum, Isparta, İzmir, Kırıkkale, Konya, Malatya, Nevşehir, Niğde, Urfa, Van, Zonguldak (Kalkandelen, 1974; Lodos & Kalkandelen, 1987b; Demir, 2006).

***Euscelis incisus* (Kirschbaum, 1868)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♂. **Distribution in Turkey:** Ankara, Burdur, Isparta, Eskişehir, Kayseri, Kırşehir (Kalkandelen, 1974; Demir, 2004b, 2006).

***Artianus manderstjernii* (Kirschbaum, 1868)**

Material examined: Balıkesir, Kazdağı National Park, Camp field, 800 m, 29.08.2004, 1♂. **Distribution in Turkey:** Ağrı, Ankara, Balıkesir, Çankırı, Diyarbakır, Gaziantep, İzmir, Kırklareli, Urfa, Van, Zonguldak (Kalkandelen, 1974; Lodos & Kalkandelen, 1987b; Demir, 2004b, 2006).

Arocephalus (s.str.) longiceps (Kirschbaum, 1968)

Material examined: Balıkesir, Kazdağı National Park, Camp field, 700 m, 06.06.2005, 5♂5♀; Camp field, 800 m, 05.06.2005, 2♂4♀; Camp field, 1000 m, 04.06.2005, 1♂5♀. **Distribution in Turkey:** Ankara, Bolu, Erzincan, Hakkari, İzmir, Konya (Lodos & Kalkandelen, 1987c; Kartal & Zeybekoğlu et. al., 2001; Demir, 2006).

Psammotettix alienus (Dahlbom, 1850)

Material examined: Balıkesir, Kazdağı National Park, Zeytinli, 150 m, 06.06.2005, 1♂. **Distribution in Turkey:** Adana, Amasya, Ankara, Balıkesir, Bursa, Diyarbakır, Erzincan, Hakkari, Iğdır, Kastamonu, Konya, Mardin, Muş, Nevşehir, Sakarya, Samsun, Siirt, Sinop, Uşak, Van, Yozgat (Kalkandelen, 1974; Lodos & Kalkandelen, 1987c; Kartal ve Zeybekoğlu, 1991; Kartal & Zeybekoğlu et. al., 2001; Demir, 2004b, 2006).

Psammotettix confinis (Dahlbom, 1850)

Material examined: Balıkesir, Kazdağı National Park, Camp field, 1000 m, 04.06.2005, 5♂3♀. **Distribution in Turkey:** Ağrı, Ankara, Balıkesir, Erzincan, Erzurum, Giresun, Hakkari, Iğdır, Isparta, Ordu, Samsun, Sivas, Van (Kalkandelen, 1974; Lodos & Kalkandelen, 1987c; Kartal ve Zeybekoğlu, 1991; Kartal & Zeybekoğlu et. al., 2001).

Jassargus (Aurkius) repletus (Fieber, 1869)

Material examined: Balıkesir, Kazdağı National Park, Top of Kazdağı, 1700 m, 28.08.2004, 4♂1♀. **Distribution in Turkey:** Amasya, Ordu, Samsun, Sinop, Tokat (Kartal & Zeybekoğlu, 1992).

Discussion

Our study is first contribution to the knowledge of the Auchenorrhyncha fauna of the Kazdağı National Park. Until now, twenty six species have been known from the Balıkesir province including the National Park. Thirty one of forty determined species are new to the Balıkesir province and the number of species known from the research area increased to fifty seven. *Bubastia (s.str.) ludviki* Dlabola, 1979 and *Iassus mirabilis* Orosz, 1979 are first records for the Turkish fauna. Before this study, *Conomelus odryssi* Dlabola, 1965 has been known only from Konya and *Balclutha rhenana* Wagner, 1939 only from Erzurum. Up to now, *Neophilaenus minor* (Kirschbaum, 1868) has been known only from Central Anatolia. We determined its presence also in the Marmara region.

Acknowledgements

I thank Prof. Dr. Ali Demirsoy as administrator of project and Msc. Yusuf Durmuş for their help during different phases of this research.

References

- Asche M.**, 1982: Beiträge zur Delphaciden-Fauna der Türkei (Anatolien) (Homoptera Cicadina Delphacidae). *Marburger Entomologische Publikation*, 1(7): 71-98.
- Başpınar H., Uygun U.**, 1991a-1992b: Doğu Akdeniz Bölgesi turuncğil bahçelerindeki Cicadellidae türleri üzerinde faunistik ve sistematik çalışmalar. *Türkiye entomoloji dergisi*, 15(2-4): 89-106, 157-172, 203-222; 16(1-2): 47-64, 99-114.
- Boulard M.**, 1979: Cigales du genre *Cicada* Linne, originaires de Turquie (Hom. Cicadidae). *Türkiye Bitki Koruma Dergisi*, 3(2): 67-74.
- Boulard M.**, 1993: *Pagiphora yanni*, Cigale anatolienne inédite. Description et premières informations biologiques (cartes d'identité et d'ethologie sonores) (Homoptera, Cicadoidea, Tibicinidae). *Türkiye entomoloji dergisi*, 17(1): 1-9.
- Demir E.**, 1998: Türkiye Homoptera Faunası İçin Yeni Kayıtlar (Homoptera, Auchenorrhyncha, Tropiduchidae). *Centr. ent. Studt., Miscellenaus Papers*, 55/56: 1-3.
- Demir E.**, 2004: *Goniagnathus guttulinervis* (Kirschbaum, 1868) new to Turkey, with data on ditribution of the Genus in Antalya (Homoptera: Auchenorrhyncha: Cicadellidae). *Acta entomologica Slovenica*, 12 (2): 255-257.
- Demir E.**, 2005a: Review of *Paradorydium* Kirkaldy (Homoptera, Auchenorrhyncha, Cicadellidae) from Turkey, with the description of a new species. *Entomological News*, 116(2): 75-82.
- Demir E.**, 2005b: Two new records of Cicadellidae fauna in Turkey (Homoptera, Auchenorrhyncha). *Zoology in the Middle East*, 35: 111.
- Demir E.**, 2006: Contributions to the knowledge of Turkish Auchenorrhyncha (Homoptera) with a new record, *Pentastiridius nanus* (Ivanov, 1885). *Munis entomology and zoology*, 1(1): 97-122.
- Dlabola J.**, 1957: Results of the Zoological Expedition of the National Museum in Prague to Turkey 20. Homoptera, Auchenorrhyncha. *Acta entomologica Musei Nationalis Pragae*, 31(469): 19-68.
- Dlabola J.**, 1971: Taxonomische und chorologische Ergänzungen der Zikadenfauna von Anatolien, Iran, Afghanistan und Pakistan (Homoptera, Auchenorrhyncha). *Acta entomologica bohemoslovaca*, 68: 377-396.
- Dlabola J.**, 1971: Taxonomische und chorologische Ergänzungen zur türkischen und iranischen Zikadenfauna (Homoptera, Auchenorrhyncha). *Acta faunistica entomologica Musei Nationalis Pragae*, 14(163): 115-138.
- Dlabola J.**, 1974: Zur Taxonomie und Chorologie einiger mediterraner Zikaden (Homoptera, Auchenorrhyncha). *Acta Zoologica Academiae Scientiarum Hungaricae*, 20(3-4): 289-308.
- Dlabola J.**, 1979: Neue Zikaden aus Anatolien, Iran und aus südeuropäischen Ländern (Homoptera, Auchenorrhyncha). *Acta Zoologica Academiae Scientiarum Hungaricae*, 25(3-4): 235-257.

- Dlabola J.**, 1981: Ergebnisse der Tschechoslowakisch-Iranischen Entomologischen Expeditionen nach dem Iran (1970 und 1973). *Acta entomologica Musei Nationalis Pragae*, 40: 127-311.
- Dlabola J.**, 1982: Fortsetzung der Ergänzungen zur Issiden-Taxonomie von Anatolien, Iran und Griechenland (Homoptera, Auchenorrhyncha). *Sbornik Narodniho Muzea V Praze*, 38(3): 113-169.
- Dlabola J.**, 1983: Neue mediterrane, meistens anatolische Issiden (Homoptera, Auchenorrhyncha). *Acta entomologica bohemoslovaca*, 80:114-136.
- Dlabola J.**, 1985a: Neue Cixiide vom Iran, Nachbarländern und anderen Mediterrangebieten (Homoptera, Auchenorrhyncha). *Acta entomologica bohemoslovaca*, 82(2): 95-128.
- Dlabola J.**, 1985b: Zwei neue *Fieberiella*-Arten aus der Türkei und Spanien (Homoptera, Cicadellidae). *Türkiye Bitki Koruma Dergisi*, 9: 75-78.
- Dlabola J.**, 1986: Neue Arten der Fulgoromorphen Zikaden-Familien vom Mittelmeergebiet und nahen Osten. *Sbornik Narodniho Muzea V Praze*, 42(3-4): 169-196.
- Dworakowska I.**, 1970: On the Genus *Arboridia* Zachv. (Auchenorrhyncha, Cicadellidae, Typhlocybinae). *Bulletin de L'Academie Polonaise des Sciences*, 18(10): 607-615.
- Dworakowska I.**, 1972: On some species of the Genus *Eupteryx* Curt. (Auchenorrhyncha, Cicadellidae, Typhlocybinae). *Bulletin de L'Academie Polonaise des Sciences*, 20(10): 727-734.
- Dworakowska I.**, 1982: Typhlocybini of Asia. *Entomologische Abhandlungen*, 45(6): 99-181.
- Emeljanov A. F.**, 2003: New taxa and new data on distribution of the subfamily Orgeriinae in the Mediterranean (Homoptera: Dictyopharidae). *Zoosystematica Rossica*, 11 (2): 311-319.
- Fahringer J.**, 1922: Eine Rhynchotenausbeute aus der Türkei, Kleinasien und den benachbarten Gebieten. *Konowia*, 1: 296-307.
- Gadeau de Kerville H.**, 1939: Recit Sommaire du Voyage en Liste Methodique des Invertebres et des Vertebres Recoltes en Asie-Mineure. Voyage Zoologique D'Henri Gadeau de Kerville en Asie-Mineure (Avril-Mai, 1912). *Paul Le Chevalier, Paris*, 148 pp.
- Gnezdilov V. M.**, 2002: New species of the genus *Tshurtshurnella* Kusnezov, 1927 (Homoptera: Cicadina: Issidae) from Turkey and Lebanon. *Russian Entomological Journal*, 11(3): 233-240.
- Güçlü Ş., Özbek H.**, 1995: Türkiye Cicadellidae (Homoptera) Faunası İçin Yeni Kayıtlar. *Türkiye entomoloji dergisi*, 19(4): 305-308.
- Güçlü Ş.**, 1996: Studies on Delphacidae (Hom., Auchenorrhyncha) from Turkey. *Turkish Journal of Zoology*, 20: 407-411.
- Haupt H.**, 1930: Ein neuer *Paralimnus* Mats. Aus Kleinasien (Homopt. Cicad.). *Deutsche Entomologische Zeitschrift*: 207-208.

- Hoch H., Remane R.,** 1985: Evolution und Speziation der Zikaden-Gattung *Hyalesthes* Signoret, 1865 (Homoptera, Auchenorrhyncha, Fulgoroidea, Cixiidae). *Marburger Entomologische Publikation*, 2(2): 1-427.
- Kalkandelen A.,** 1974: Orta Anadolu'da Homoptera: Cicadellidae familyası türlerinin taksonomileri üzerinde arařtırmalar. *Zirai Mücadele ve Karantina Genel Müdürlüğü, Ankara*, 1-220.
- Kalkandelen A.,** 1980: Contributions to the families Delphacidae and Cicadellidae (Homoptera) from Turkey. *Türkiye Bitki koruma Dergisi*, 4(3): 147-154.
- Kalkandelen A.,** 1985: Four new species of genus *Zyginidia* (*Zyginidia*) Haupt (Homoptera: Cicadellidae) and with notes on the taxonomy and distributions of the species of this genus in Turkey. *Türkiye Bitki koruma Dergisi*, 9: 13-25.
- Kalkandelen A.,** 1987-2000: Türkiye Cixiidae (Homoptera) türleri üzerine taksonomik çalışmalar. *Bitki Koruma Bülteni*, 27(3-4): 119-146; 28(3-4), 113-140; 29(1-4): 1-17, 117-132; 30(1-4): 3-27; 33(3-4): 65-82; 34(1-2): 1-21; 40(3-4): 91-123.
- Kartal V.,** 1980: Neue und wenig bekannte Arten der Gattungen *Cicadetta* und *Cicadatra* aus dem Irak und der Türkei (Homoptera, Auchenorrhyncha, Cicadidae). *Entomologische Zeitschrift*, 17: 185-192.
- Kartal V.,** 1981: Neue Homopteren aus der Türkei-I. *Priamus*, 1(1): 24-30.
- Kartal V.,** 1983: Neue Homopteren aus der Türkei II (Homoptera Auchenorrhyncha). *Marburger Entomologische Publikation*, 1(8): 235-248.
- Kartal V.,** 1985: Neue und wenig bekannte *Tshurtshurnella*-Arten (Homoptera Auchenorrhyncha Issidae) aus der Türkei. *Marburger Entomologische Publikation*, 1(10): 191-218.
- Kartal V.,** 1985: Türkiye Yukarı Kızılırmak Havzasındaki Issidae (Homoptera, Auchenorrhyncha) Familyası Türlerinin Taksonomik Yönden İncelenmesi. *Doğa*, A2: 64-77.
- Kartal V.,** 1986: Wenig bekannte und für die Türkei neue *Nymphorgerius*-Arten (Homoptera, Auchenorrhyncha, Dictyopharidae). *Türkiye Bitki koruma Dergisi*, 10(2): 99-103.
- Kartal V.,** 1987: Eine neue und wenig bekannte Arten der Gattung *Ranissus* aus der Türkei (Homoptera, Auchenorrhyncha, Dictyopharidae). *Türkiye Entomoloji Dergisi*, 11(3): 145-150.
- Kartal V.,** 1988: Türkiye Cicadellid Faunası İçin Yeni Bir Cins *Kazachtanicus* Dlab.,1961 (Homoptera, Auchenorrhyncha, Cicadellidae). *Doğa Zooloji*, D. : 68-69.
- Kartal V., Zeybekođlu Ü.,** 1992: Türkiye Cicadellidae (Hom., Auchenorrhyncha) Faunası İçin Yeni Türler. *Turkish Journal of Zoology*, 16: 349-352.
- Kartal V., Zeybekođlu Ü.,** 1994: A New Species *Allygidius nihati* n. sp. (Hom., Auch., Cicadellidae) from Turkey. *Turkish Journal of Zoology*, 18: 259-261.
- Kartal V., Zeybekođlu Ü.,** 1994: Türkiye Cicadellidae (Hom., Auchenorrhyncha) Faunası İçin Yeni Bir Tür Kaydı. *Turkish Journal of Zoology*, 18: 37-39.

- Kartal V., Zeybekoğlu Ü.,** 1997: Türkiye Faunası İçin Yeni Üç *Diplocolenus* Rib. (Hom., Auch., Cicadellidae, Deltocephalinae) Türü Üzerine Taksonomik Bir Araştırma. *Turkish Journal of Zoology*, 21: 291-294.
- Kartal V., Zeybekoğlu Ü., Dursun A.,** 2001: A faunistic study on Deltocephalinae (Homoptera: Auchenorrhyncha: Cicadellidae) in Yüksekova (Hakkari). *Türkiye Entomoloji Dergisi*, 25(2): 83-92.
- Linnavuori R.,** 1965: Studies on the South and East-Mediterranean Hemipterous fauna. *Acta Entomologica Fennica, Helsinki*, 21: 1-70.
- Lodos N., Kalkandelen A.,** 1980a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey I. Family Cixiidae Spinola. *Türkiye Bitki koruma Dergisi*, 4(1): 15-27.
- Lodos N., Kalkandelen A.,** 1980b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey II. Family Delpacidae Leach. *Türkiye Bitki koruma Dergisi*, 4(2): 103-117.
- Lodos N., Kalkandelen A.,** 1980c: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey III. Families Meenoplidae, Derbidae, Achilidae, Dictyopharidae and Tettigometridae. *Türkiye Bitki koruma Dergisi*, 4(3): 161-178.
- Lodos N., Kalkandelen A.,** 1981a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey IV. Family Issidae Spinola. *Türkiye Bitki koruma Dergisi*, 5(1): 5-21.
- Lodos N., Kalkandelen A.,** 1981b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey V. Families Flatidae, Ricaniidae and Cicadidae. *Türkiye Bitki koruma Dergisi*, 5(2): 67-82.
- Lodos N., Kalkandelen A.,** 1981c: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey VI. Families Cercopidae and Membracidae. *Türkiye Bitki koruma Dergisi*, 5(3): 133-149.
- Lodos N., Kalkandelen A.,** 1981d: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey VII. Family Cicadellidae: Ulopiniae, Megophthalminae, Ledrinae, Macropsinae and Agallinae. *Türkiye Bitki koruma Dergisi*, 5(4): 215-230.
- Lodos N., Kalkandelen A.,** 1982a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey VIII. Family Cicadellidae: Idiocerinae. *Türkiye Bitki koruma Dergisi*, 6(1): 15-28.
- Lodos N., Kalkandelen A.,** 1982b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey IX. Family Cicadellidae: Iassinae, Penthiminae, Dorycephalinae, Hecalinae and Aphrodinae. *Türkiye Bitki koruma Dergisi*, 6: 147-159.
- Lodos N., Kalkandelen A.,** 1983a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey X. Family Cicadellidae: Xestocephalinae, Stegelytrinae and Cicadellinae. *Türkiye Bitki koruma Dergisi*, 7: 23-28.
- Lodos N., Kalkandelen A.,** 1983b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XI. Family Cicadellidae,

- Typhlocybinæ: Alebrini and Dikraneurini. *Türkiye Bitki koruma Dergisi*, 7: 107-115.
- Lodos N., Kalkandelen A.**, 1983c: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XII. Family Cicadellidae, Typhlocybinæ: Emposcini. *Türkiye Bitki koruma Dergisi*, 7: 153-165.
- Lodos N., Kalkandelen A.**, 1984a-b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XIII. Family Cicadellidae, Typhlocybinæ: Typhlocybini. *Türkiye Bitki koruma Dergisi*, 8: 33-44, 87-97.
- Lodos N., Kalkandelen A.**, 1984c-d: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XV. Family Cicadellidae, Typhlocybinæ: Erythroneurini. *Türkiye Bitki koruma Dergisi*, 8: 159-168, 201-210.
- Lodos N., Kalkandelen A.**, 1985a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XVII. Family Cicadellidae, Deltocephalinae: Grypotini, Goniagnathini and Opsini. *Türkiye Bitki koruma Dergisi*, 9: 79-90.
- Lodos N., Kalkandelen A.**, 1985b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XVIII. Family Cicadellidae, Deltocephalinae: Macrostelini. *Türkiye Bitki koruma Dergisi*, 9(3): 147-161.
- Lodos N., Kalkandelen A.**, 1985c: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XIX. Family Cicadellidae, Deltocephalinae: Deltocephalini, Scaphytopiini, Doraturini. *Türkiye Bitki koruma Dergisi*, 9: 207-215.
- Lodos N., Kalkandelen A.**, 1986a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XX. Family Cicadellidae, Deltocephalinae: Fieberiellini, Stirellini and Tetartostyliini. *Türkiye Bitki koruma Dergisi*, 10(1): 25-32.
- Lodos N., Kalkandelen A.**, 1986b-87b: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XXI. Family Cicadellidae, Deltocephalinae: Athysanini. *Türkiye Bitki koruma Dergisi*, 10(3-4): 131-139, 203-211; 11(1-2): 29-40, 97-109.
- Lodos N., Kalkandelen A.**, 1987c-d: Preliminary list of Auchenorrhyncha with notes on distribution and importance of species in Turkey XXV. Family Cicadellidae, Deltocephalinae: Paralimnini Distans. *Türkiye Entomoloji Dergisi*, 11(3-4): 151-162, 195-202.
- Lodos N., Kalkandelen A.**, 1988a: Preliminary list of Auchenorrhyncha with notes on distribution and importance of Turkey XXVII. (Addenda and Corrigenda). *Türkiye Entomoloji Dergisi*, 12(1): 11-22.
- Meyer-Arndt S.**, 1988: Vier neue Arten der Gattung *Fieberiella* Signoret, 1879 aus Griechenland und der Türkei (Homoptera Auchenorrhyncha Cicadellidae). *Marburger Entomologische Publikation*, 2(4): 179-208.

- Meyer-Arndt S.**, 1991: A new species of *Fieberiella* Signoret, 1879 (Hem., Cicadellidae) from western Turkey and the Aegean Islands with new locality records for the genus. *Entomologist's Monthly Magazine*, 127: 61-72.
- Zachvatkin A. A.**, 1937: Deux *Zyginidia* nouvelles des Turquie (Homoptera, Eupterygidae). *Soc. Nat. Moscou (Sect. Biol.) Bull.*, 46: 317-322.
- Zachvatkin A. A.**, 1946: Studies on the Homoptera of Turkey. *Trans. R. Ent. Soc. London*, 97(6): 149-176.
- Zeybekoğlu Ü., Bulut N.**, 2000: Türkiye Cicadellidae (Homoptera, Auchenorrhyncha) Faunası İçin Yeni İki Tür Kaydı ve Taksonomik Özellikleri. *OMÜ Fen-Edebiyat Fakültesi Fen Dergisi*, 11(1): 85-90.
- Zeybekoğlu Ü.**, 1996: Türkiye'de saptanan Ebarrius Ribaut, 1946 (Homoptera, Auchenorrhyncha, Cicadellidae, Deltocephalinae) cinsine bağlı türler üzerine sistematik bir araştırma. *Turkish Journal of Zoology*, 20: 187-189.
- Zeybekoğlu Ü.**, 1998: The species of Deltocephalinae (Homoptera: Auchenorrhyncha: Cicadellidae) found in the Middle and East Black Sea Regions. *Türkiye Entomoloji Dergisi*, 22(1): 37-45.

Received / Prejeto: 22. 3. 2006

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Acta Entomologica Slovenica](#)

Jahr/Year: 2006

Band/Volume: [14](#)

Autor(en)/Author(s): Demir Emine

Artikel/Article: [PRELIMINARY REPORT ON THE AUCHENORRHYNCHA \(HEMIPTERA\) FAUNA OF KAZDA?I NATIONAL PARK WITH TWO NEW RECORDS FOR TURKEY* 89-102](#)