An illustrated key to the genera of Hebridae (Gerromorpha) in Thailand by Herbert Zettel

Abstract: Five genera of the family Hebridae are known from Thailand. An illustrated key is presented and notes on the habitat preferences are given.

Introduction

Hebrids (Velvet Water Bugs) are very small insects, which live predominantly or obligatorily predacious. Hebridae is a basic family of the suborder Gerromorpha ("Semiaquatic Bugs"), but most species inhabit terrestrial habitats or the extreme edge of the water. Only a few species live steadily at the water surface, and a few are even subaquatic.

The morphology and systematics of Hebridae have been excellently treated by ANDERSEN (1981, 1982). Since then, only one genus, *Nieserius* ZETTEL, 1999, has been added. In Thailand the family is represented by five genera in two subfamilies: *Hyrcanus* DISTANT, 1910, and *Nieserius* of the exclusively Oriental Hyrcaninae, which was recently revised by the author (ZETTEL 1998, 1999a); and *Hebrus* CURTIS, 1833, *Merragata* BUCHANAN-WHITE, 1877, and *Timasius* DISTANT, 1909, of the Hebrinae.

Hebridae can be easily distinguished from other Thai Gerromorpha by using the key by CHEN & ZETTEL (1996).

Key to the genera of Hebridae in Thailand

(modified after ANDERSEN 1981, 1982)

Very small species (body length less than 1.8 mm); head short and broad; eyes pedunculate, situated close to antero-lateral margin of pronotum (Fig. 8); antenna slender (Fig. 1); corium of forewing with white patch; femora slender (Fig. 7).

One species, *Merragata pallescens* DISTANT, 1909, is known from Thailand. It inhabits the edges of large stagnant waters (see ZETTEL 1999b).

3 Antennal segment 1 shorter than segments 2 and 4 (Fig. 2); eye far removed from the pronotum, distance between eye and antero-lateral margin of the pronotum more than 0.4 times width of eye (Fig. 10); head below eyes with-



Figs. 1 - 10: (1 - 5) antenna of (1) *Merragata pallescens*, (2) *Nieserius subaquaticus*, (3) *Hyrcanus draculus*, (4) *Hebrus cruciatus*, (5) *Timasius miyamotoi*; (6, 7) hind leg of (6) *Nieserius subaquaticus* and (7) *Merragata pallescens*; (8 - 10) head, lateral view, of (8) *Merragata pallescens*, (9) *Hyrcanus draculus*, and (10) *Nieserius subaquaticus*.



Figs. 11 12: Meso- and metasternum and base of abdomen, schematically, in (11) *Timasius* and (12) *Hebrus*.

One species is known from North Thailand. The biology of *Nieserius* differs from that of all other Hebridae: *Nieserius* lives subaquatically on the fine sediments of smaller streams and brooks (ZETTEL 1999a).

Four species are so far known from Thailand, two of which seem to be endemic. *Hyrcanus* species inhabit places associated with brooks, streams, and rivers, either hygropetric (especially in mosses) or semiaquatic between leaf litter at the edge or even in the middle of streams. Probably some species go into the water, too (ZETTEL 1998).

4 Paired, longitudinal carinae of thoracic venter converging and meeting before hind margin of metasternum (Fig. 11)
Timasius

Only two species are reported from Thailand, but in fact the number of species is much higher. Two species groups, the *T. livens* group and the *T. chinai* group (see ANDERSEN 1981), form complexes of allopatric species with high radiation in the Indo-Chinese area. *Timasius* species are typically found hygropetric on wet, shaddy rock faces at the edge of running waters.

The Oriental species of the genus *Hebrus* are unrevised. Many, mostly unnamed, species are known from Thailand. Nearly all Oriental *Hebrus* species are hygropetric (often on mosses) or ripicolous, but a few species are also found at the edge of the surface of stagnant waters. Specimens of *Hebrus cruciatus* DISTANT were found under large stones on a river bank in Mindanao, Philippines, where they found shelter from the heat at noon; this species occurs also in Thailand.

Check-list of Hebridae species known from Southeast Asia

NHMW = Natural History Museum Vienna

.

Subtamily Hyrcaninae	
Hyrcanus chenae ZETTEL, 1998	China (Yunnan)
Hyrcanus draculus ZETTEL, 1998	Thailand (Chiang Mai, Mae Hong Son), North Laos, South China
Hyrcanus saxatilis ANDERSEN, 1981	Thailand (Chiang Mai)
Hyrcanus shepardi ZETTEL, 1998	Thailand (Phetchabun)
Hyrcanus varicolor Andersen, 1981	Thailand (widespread), Viet Nam, Indonesia (Sumatra, Java)
Nieserius subaquaticus ZETTEL, 1999	Thailand (Chiang Mai, Mae Hong Son)
Nieserius brachypterus ZETTEL, 1999	North Laos
Subfamily Hebrinae	
<i>Merragata pallescens</i> (DISTANT, 1909)	India, Thailand, Java (see ZETTEL 1999b)
<i>Timasius livens</i> Andersen, 1981	Thailand (Chiang Mai)
<i>Timasius miyamotoi</i> Andersen, 1981	Thailand (Chiang Mai)
Timasius spp. (several undescribed species from Thailand in NHMW)	
Hebrus cruciatus (DISTANT, 1910)	Philippines, Thailand (Chiang Mai, NHMW)
Hebrus nereis Polhemus & Polhemus, 1989	Thailand (Andersen, unpubl. list), Singapore

Hebrus spp. (numerous undescribed species from Thailand in NHMW)

Acknowledgements: My thanks are due to all colleagues who supported this study by sending material for study, especially to N. Nieser (Tiel) and to W.D. Shepard (Sacramento); and to my friends in Thailand, who helped me so much during my field studies.

References

- ANDERSEN, N.M. 1981: Semiaquatic bugs: phylogeny and classification of the Hebridae (Heteroptera: Gerromorpha) with a revision of *Timasius*, *Neotimasius* and *Hyrcanus*. Systematic Entomology 6: 377-412.
- ANDERSEN, N.M. 1982: The Semiaquatic Bugs (Hemiptera, Gerromorpha). Phylogeny, adaptations, Biogeography and Classification - Entomonograph 3: 1-455.
- CHEN, P.P. & ZETTEL, H. 1996: An illustrated key to the families of Gerromorpha in Thailand. Amemboa 1: 10-13.

- ZETTEL, H. 1998: Eine taxonomische Revision der Gattung *Hyrcanus* DISTANT 1910 (Heteroptera: Hebridae) mit Neubeschreibungen von vier Arten aus Indien, Thailand, Laos und China. Stapfia 55: 585-606.
- ZETTEL, H. 1999a: *Nieserius* gen.n., a new genus of the subfamily Hyrcaninae (Heteroptera: Hebridae) from Thailand, Laos, and Nepal, with the first known subaquatic species of Gerromorpha. Aquatic Insects 21(1): 39-52.
- ZETTEL, H. 1999b: The genus *Merragata* BUCHANAN-WHITE, 1877 (Gerromorpha: Hebridae) in Thailand. Amemboa 3: 6-8.

Author's address: Dr. Herbert Zettel, Naturhistorisches Museum Wien, 2. Zoologische Abteilung, Burgring 7, A-1014 Vienna, Austria (e-mail: herbert.zettel@nhm-wien.ac.at)

Opinion:

Owners of the Bugs

Roam a meadow and enjoy nature. Observe the thousands of buzzing, creeping and swarming insects and think: Who is their owner?

This crazy thought has a real background. In our materialistic world everything (and everybody? see discussion on genetic "resources" of ethnic minorities) *has* to be owned. Since the agreement in the "Convention of Biodiversity" wild animals (including protozoes, worms, and insects), plants, and even microbes are owned by the states, who own the rights to self-controlled exploitation. The "common heritage principle" (in my opinion even this term is worth discussion) was replaced by the "common concern principle". Respect for nature is obviously not even considered.

It is not only a Christian attitude, than men subject Mother Earth. In most "civilizations" nature is owned by somebody; those creatures which are not owned by a person or a company are owned by the state. Ecological movements did not reduce this trend, but even enforced it involuntarily. Ecologists said, "be responsible for our nature", but politicians heard only "our nature": they made laws stating that wildlife (more modern: its genetic "resources") belongs to the country, which must be a little responsible for it ("... not too much, please, respect economic needs!"), but can use it as it likes. Biodiversity projects were (and still are) justified by (often not existing) follow-up research on usage of yet undiscovered species (in medical plants, etc.). Politicians heard "usage" and made laws to regulate this possible use in advance.

One of the disastrous consequences of the "Convention of Biodiversity" will be a quickly growing administration and regulation of biodiversity research (especially concerning export regulations), which may result even in a de facto end to international cooperation. Consequently, nature destruction may overtake biodiversity research, especially in those tropical countries where scientific cooperation is essential for well-founded nature protection.

Be sure: It's good for the economy!

Herbert Zettel

Entomological Department Naturhistorisches Museum Wien Burgring 7, A-1014 Vienna, Austria e-mail: herbert.zettel@nhm-wien.ac.at

P.S.: Yesterday, at a newspaper kiosk, I read a headline in a popular journal: "Biodiversity: There are at least 13 million living creatures. - Who really needs them all?"

Imprint: "*Amemboa*" is a newsletter for taxonomical, systematical, and faunistical co-operative works on Thai Heteroptera. Editor: Dr. Herbert Zettel, Naturhistorisches Museum Wien, 2. Zoologische Abteilung, Burgring 7, A-1014 Vienna, Austria. Proof reading: Prof. Dr. C. W. Schaefer, Storrs, USA. Published by Naturhistorisches Museum Wien. © Naturhistorisches Museum Wien, 1999. Authors are fully responsible for the content of their papers.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Amemboa

Jahr/Year: 1999

Band/Volume: 3_1999

Autor(en)/Author(s): Zettel Herbert

Artikel/Article: <u>An illustrated key to the genera of Hebridae (Gerromorpha) in Thailand. 22-</u> 26