

On the distribution of *Astrobus laevipes* CANESTRINI, 1872 (Arachnida: Opiliones) in Central Europe

Anke Maria HÖFER & Jörg SPELDA

Abstract: Published data and unpublished communications show that the range of *Astrobus laevipes* in Central Europe is much larger than previously believed. The present review extends the list of records to the German states of Baden-Württemberg, Hesse, Rhineland-Palatinate, Bavaria, Northrhine-Westphalia, Saxony, and Lower Saxony and provides a map of the present distribution in Germany. Furthermore, it lists new findings of *A. laevipes* in the Czech Republic and Hungary (Rakaca/Serehat Valley). Records of *A. laevipes* in Austria are not included in this review.

INTRODUCTION

Astrobus laevipes was first recorded in Germany in Dresden (HERBST 1799, under *Opilio spinosus*). An additional and isolated finding was reported by Höregott (cited in MARTENS 1978) in Mainz (Rhineland-Palatinate). MARTENS (1978) stated that the distribution of the thermophile harvestman *A. laevipes* in Central Europe is limited to the countries of the Carpathians, the Hungarian lowland including the valleys of the rivers Elbe and Danube, and the plains of the river Po, with an isolated population in Mainz, Germany. However, more recent publications as well as our present data show that the range is much larger than previously believed and that the population around Mainz is not isolated. We present here a summary of all records of *A. laevipes* in Germany, the Czech Republic and Hungary, including a characterisation of the biotopes where we have found the species ourselves. These data can serve as a base for further studies in this species.

RESULTS

The new records of *A. laevipes* are summarised in Table 1 and shown in Figure 1. A colour version of the map indicating topographical features can be found at <http://www.spiderling.de.vu/>. The majority of the new findings

presented here were obtained from pitfall traps. As the records are the result of occasional sampling in studies of numerous investigators no standard methods can be described. The single female obtained at Karlsruhe-Grötzingen was found on April 7th, 1994 under a stone between a hedge and a narrow pass during manual sampling. The locality is situated in a limestone area (Middle Triassic: shell limestone) between vineyards and orchards. Similarly, the findings at the "Drachenfelser Ländchen", near Bonn also were made near vineyards (Trachyt-Rigosol mixed with stony loam), where over 140 individuals were pitfall-trapped in the surrounding undergrowth between March 27th and October 10th, 1996.

DISCUSSION

At present the known distribution of *A. laevipes* suggests an isolated occurrence/population in western Germany. But this might only be an artefact of insufficient knowledge. The map given by MARTENS (1978) as well as our presented data (Tab. 1, Fig. 1) suggests that *A. laevipes* mainly used the valleys of the big streams for spreading to the south. The question is where and when the species was able to overcome the distance between the river systems of the Danube and the rivers flowing into the North Sea. The first step was to cross the Bohemian depression, a very warm area containing many southeast European floristic (e. g. *Iris aphylla*) and faunistic elements. Although our knowledge has increased, it remains uncertain how *A. laevipes* was able to overcome the distance between the river systems of the Elbe and the Rhine. There are at least three possibilities:

1. Along the northern border of the highlands
2. Across Thuringia along Saale-Rodach into the Main valley
3. Along the Danube and later either across Altmühl-Regnitz, Altmühl-Tauber or Wörnitz-Jagst.

The first possibility avoids overcoming highlands, but would be an unusual route for southeast European elements. The second route demands transition of the Thuringian Forest, a rather cold area, but this route is mostly supported by the presently known records. The third possibility is a well-known route for several southeast European elements, which occur either in the Jagst- or the Tauber-valley (e. g. the millipede *Mastigona bosniense*, SPELDA 1999). However we have no records of *A. laevipes* from the Bavarian part of the Danube river system, which might support this route.

Table 1: Records of *Astrobus laevis* CANESTRINI, 1872 in Germany, the Czech Republic and Hungary.

Country	State	Locality	MTB-grid	Reference
Germany	Baden Württemberg	Hoehberg, near Werbach and Tauber-bischofsheim	6323-4	1987, leg. Raqué, det. Harms* vid. Rausch cited in BAUMANN et al. (1992) vid. Spelda
		„Schwabenheimer Insel“ near Heidelberg N of Grötzingen, 7 km E of Karlsruhe	6517-4 6917-3	1987, leg. Raqué, det. Harms, vid. Rausch* 1994, Spelda*
	Hessia	Near Offenbach	5818/5918	LANG et al. (1993)
		Botanical Garden, Frankfurt/Main	5817-4	1991, Malten*
		NSG (Nature Reserve) „Lorcher Werth“ County of Rheingau-Taunus	5912-2	1993, Malten*
		Riverbanks of the Main, W of Kelsterbach	5917-1	1999, 2000, Malten*
		Cityforest Frankfurt/Main	5918-1	
		Cityforest Frankfurt/Main, Monte Scherbelino („alter Müllberg“)	5918-1	2000, Malten*
	Rhineland-Palatinate	Frankfurt/Main E and Riverbanks of the Nidda, Frankfurt/Main-Berkersheim	5818-1	
		„Gonsenheimer Sand“, Forest of Gonsenheim, near Mainz	5915-3	HÖREGOTT (1963), cited in MARTENS (1978)
		NSG „Sandlache“ near Ingelheim, County of Mainz-Bingen	5914-3	1995, Malten*
	Bavaria	„Bopparder Hamm“	5711-2	2000, Jakobs*
		Riverbanks of the Main near Klingenberg	6221-1	BAUMANN et al. (1992)
		Riverbanks of the Main near Klingenberg	6221-1	LANG et al. (1993); same record as previous one
		Riverbanks of the Main in S-Trennfeld and Trennfeld-Sand near Wertheim	6223-2	
	Northrhine-Westfalia	Riverbanks of the Main in Erlenbach near Wörth	6120-4	2000, Stumpf*
		„Drachenfelder Ländchen“, 20 km S of Bonn	5309-1	1996, Höfer*

* = unpublished

* = recorded here for the first time

Country	State	Locality	MTB-grid	Reference
Germany	Saxony-Anhalt	Dübener Heide	4340-1/2	BLISS (1993), BLISS (1990) and BLISS & TIETZE (1984), both cited in BLISS (1993)
		NSG "Porphyrylandschaft bei Gimritz"	4437-1	BLISS (1993), SCHNITTER (1991) cited in BLISS (1991)
		Sennewitz	4437-4	BLISS (1993), LANG (1989) and BLISS (1991b) both cited in BLISS (1993)
		Dresden	4949-3	HERBST (1799), cited in MARTENS 1978 and BLISS (1993)
	Saxony	Schlottwitz, "Hohler Stein	5148-2	Schiemenz, cited in MARTENS (1978) and BLISS (1993)
		Kunzwarda/Torgau (Elbaue)	4444-3	Tietze, cited in MARTENS (1978) and Bliss (1993)
		Freital-Zauckerode	4947-4	BLISS (1993)
		"Götzenbusch", Freital	5047-4	
		NSG "Pillnitzer Elbinsel", Dresden	4949-3	
		Prostitz "Ketzerbachtal"	4846-1	
		NSG "Ziegenbusch"	4847-1	
		Falkenberg	4342-4	
Czech Republic		"Wachtelberg"	4642-1	KLIMEŠ & ROUŠAR (1998)
		Wölpern, NSG "Torfwiesen"	4541-4	
		Povrly, near Ústí nad Labem	5350-2	
		Lipenec, near Louny	5648-3	
Hungary		South Moravia: Pálava: Devin, National Nature Reserve of Ranspürk, Nature Reserve of Krivé jezero, Nature Reserve of Slanisko	6573-2	KRISTEK (1991) and BEZDĚČKA & ROZKOŠÝ (1995) both cited in KLIMEŠ & ROUŠAR (1998)
		South Moravia	not to classify	
		River valley of river Rakaca-Cserehát	not to classify	BOKOR (1996)


* = unpublished

= recorded here for the first time

Although MARTENS (1978) discusses the possibility of a spreading in recent times when deforestation took place it is as well possible, that the species invaded its western range just after the last glaciation before the forest had covered most parts of Germany. Or it could have invaded its range along the streams, which cause open areas along their borders by flooding. The latter possibility does not restrict the spreading to a special period.

Astrobunus laevipes often has been confused with *A. bernardinus* (SCHENKEL 1926). They are believed to be sibling species (MARTENS 1978), and it is possible that these species hybridise in the Southern Alps. However there are clear differences in the ecology of the two species. *Astrobunus laevipes* is confined to low altitudes, while *A. bernardinus* has an alpine distribution, even occurring above the timberline. It has a wide range of habitats, but avoids dry, open, southern slopes. Another species occurring parapatric to both *A. laevipes* and *A. bernardinus* is *A. helleri*, occurring in the central and eastern parts of the Alps (KOMPOSCH 1997, 1999). It is very likely that the distribution of all three species is influenced by competition. *Astrobunus laevipes* and *A. bernardinus* seldomly occur in sympatry (Drau-valley near Spittal, leg. Ausobsky cited in MARTENS 1978). In areas where *A. helleri* and *A. laevipes* occur sympatric, *A. helleri* inhabits the higher altitudes but, in contrast to *A. bernardinus*, avoids the alpine region. It is very likely, that all three species arose from a single ancestor during the Pleistocene. To test this hypothesis, explain the present distribution of *A. laevipes*, and the relationship between populations we need further investigations and a phylogenetic analysis of this species.

Acknowledgments: Andreas MALTEN (Dreieich), Helmut STUMPF (Würzburg), Karl-Hermann HARMS (Rheinstetten) and Kerstin JACOBS (Boppard) kindly contributed unpublished records. Aloys STAUDT (Schmelz) prepared the distribution map. Theo BLICK (Hummeltal) supplied references and was always a helpful source of information.



REFERENCES

- BAUMANN, T., T.BLICK, F.FOECKLER & M.SCHLEUTER (1992): Erstnachweis von *Astrobunus laevipes* und *Nemastoma dentigerum* in Bayern (Opiliones: Phalangidae, Nemastomatidae). - Arachnol. Mitt. 3: 58-61
- BEZDĚČKA, P. & R.ROZKOŠNÝ (1995): Opilionida. In: R. ROZKOŠNÝ & J.VAÑHARA (eds.): Terrestrial invertebrates of the Pálava Biosphere Reserve of UNESCO. I. - Folia Fac. Sci. Nat. Univ. Masarykianae Brunensis, Biol. 92: 55-58
- BLISS, P. (1990): Ökologische Untersuchungen an Weberknechten (Arachnida, Opiliones) im Rauchschadensgebiet Dübener Heide. Diplomarbeit, Universität Halle, 156 p.
- BLISS, P. (1993): Neue Funde von *Astrobunus laevipes* (Arachnida, Opiliones, Palangiidae). - Boll. Soc. Neuchatel. Sci. Nat. 116 (1): 35-39
- BLISS, P. & H.HIEBSCH (1984): Verzeichnis der Weberknechte für das Gebiet der DDR. - Entomol. Nachr. u. Ber. 28:199-200
- BLISS, P. & F.TIETZE (1984): Die Struktur der epedaphischen Weberknechtfauna (Arachnida, Opiliones) in unterschiedlich immisionsbelasteten Kieferforsten der Dübener Heide. - Pedobiol. 26: 25-35
- BLICK, T. & A.HÄNGGI with co-operation of K.THALER (2000): Checklist of the arachnids of Germany, Switzerland and Austria (Arachnida: Araneae, Opiliones, Pseudoscorpiones, Scorpiones, Palpigradi). Preliminary version 2000-07-07. - Internet: http://www.arages.de/checklist_e.html
- BOKOR, Z. (1996): New Occurrence of *Astrobunus laevipes* (CANESTRINI, 1872) (Arachnoidea, Opiliones, Phalangidae) in Hungary (in the river valley of river Rakaca - Cserehát). - Acta Univ. Szegediensis Acta Biol. 41: 89-91
- HERBST, J.F.W. (1799): Natursystem der ungeflügelten Insekten. Drittes Heft. Naturgeschichte der Insectengattung *Opilio*. IV, 30 p., Berlin.
- HÖREGOTT, H. (1963): Zur Ökologie und Phänologie einiger Chelonethi und Opiliones (Arachn.) des Gonsenheimer Sandes und Waldes bei Mainz. - Senckenbergiana biol. 44: 545-541.
- KLIMEŠ, L. & A.ROUŠAR (1998): Remarkable harvestmen from the Czech Republic. - Arachnol. Mitt. 16: 33-39
- KOMPOSCH, C. (1997): Kommentierte Checkliste der Weberknechte (Opiliones) Kärntens. - Carinthia II 187/107: 597-608
- KOMPOSCH, C. (1999): Rote Liste der Weberknechte Kärntens (Arachnida: Opiliones). In: HOLZINGER, W.E.: Rote Liste gefährdeter Tiere Kärntens. - Schriftenreihe Naturschutz Kärnten 15: 1-718
- KRISTEK, J. (1991): Selected groups of insects and harvestmen. In: PENKA, M., M.VYSKOT, E.KLIMO & F.VAŠICEK (eds.): Floodplain forest ecosystem. II. After water management measures. Elsevier-Academia, Amsterdam, p. 451-468
- LANG, A. (1989): Ökofaunistische Untersuchungen in Pappelstandorten der Agrarlandschaft bei Halle (Saale) unter besonderer Berücksichtigung der Myriapoden (Diplopoda, Chilopoda) und Isopoda (Oniscoidae). Ph.D. Thesis, Pädagogische Hochschule Halle-Köthen, 154 p. + appendix.
- LANG, A., M.KLEIN, F.FOECKLER & M.SCHLEUTER (1993): On a recent record of *Astrobunus laevipes* and *Nemastoma dentigerum* in Bavaria, Germany (Opiliones: Phalangidae, Nemastomatidae). - Boll. Acc. Gioenia Sci. Nat. 26 (345): 227-235

- MARTENS, J. (1978): Spinnentiere, Arachnida – Weberknechte, Opiliones. - In: Die Tierwelt Deutschlands, 64. Teil. 464 p., Jena.
- SCHENKEL, E. (1926): Beitrag zur Kenntnis der Schweizerischen Spinnenfauna. - Rev. Suisse de Zool. 33 (5): 301-316
- SPELDA, J. (1999): Verbreitungsmuster und Taxonomie der Chilopoda und Diplopoda Südwestdeutschlands. Diskriminanzanalytische Verfahren zur Trennung von Arten und Unterarten am Beispiel der Gattung *Rhymogona* COOK, 1896 (Diplopoda: Chordeumatida: Craspedosomatidae). Ph.D. Thesis, University of Ulm. Part 1: 1-217, Part 2: 1-324

Anke Maria HÖFER, Australian National University, School of Botany and Zoology, AUS-0200 Canberra, A.C.T. Australia
e-mail: am.hoefer@anu.edu.au

Jörg SPELDA, Asangstrasse 49, D-70329 Stuttgart
e-mail: spelda@t-online.de

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Arachnologische Mitteilungen](#)

Jahr/Year: 2001

Band/Volume: [22](#)

Autor(en)/Author(s): Höfer Anke Maria, Spelda Jörg

Artikel/Article: [On the distribution of *Astrobonus laevipes* CANESTRINI, 1872 \(Arachnida: Opiliones\) in Central Europe 42-49](#)