

**A new species of *Tongeia* TUTT, [1908] from
Northeast Yunnan, China**

(Lepidoptera: Lycaenidae)

by

HAO HUANG¹ & ZHONG CHEN²

received 7.III.2006

1: Qingdao Vocational and Technical College, Qingdao, 266555, P. R. China
(email: cmdhlxx@hotmail.com)

2: 304, #14, #2 Xiang-Fu-Ying, Nanjing, 210018, P. R. China (email: czcloud@126.com)

Abstract: *Tongeia dongchuanensis* **spec. nov.** is described from Dongchuan, NE. Yunnan, China. It is closely allied to *Tongeia potanini* (ALPHERAKY, 1889) in phylogeny and is similar to *Tongeia zuthus* (LEECH, [1893]) and *Tongeia hainani* (BETHUNE-BAKER, 1914) in appearance. For comparison, all the known species of *Tongeia* TUTT, [1908] are discussed and a key is given.

Zusammenfassung: *Tongeia dongchuanensis* **spec. nov.** wird von Dongchuan aus dem Nordosten von Yunnan, China beschrieben. Die Art ist aufgrund der Phylogenie nahe mit *Tongeia potanini* (ALPHERAKY, 1889) und phaenotypisch mit *Tongeia zuthus* (LEECH, [1893]) und *Tongeia hainani* (BETHUNE-BAKER, 1914) verwandt. Zum Vergleich werden alle bekannten Arten der Gattung *Tongeia* TUTT, [1908] aufgeführt und in einem Bestimmungsschlüssel eingegliedert.

Introduction: Recently a strange looking ♂ specimen of the genus *Tongeia* TUTT, [1908] was collected by the junior author from Dongchuan, NE Yunnan. After a careful study, including dissection of the ♂ genitalia, the authors have concluded that this unique specimen represents a distinct new species, here described as

***Tongeia dongchuanensis* spec.nov.**

Holotype ♂ (colour plate 4: G), Length of forewing 14,5 mm. Dongchuan, NE Yunnan, China. No detailed collecting data.

Etymology. This new species is named after its type locality, Dongchuan.

Diagnosis

In external features, this new species is very similar to *T. zuthus* (LEECH, [1893]) from Sichuan and *T. hainani* (BETHUNE-BAKER, 1914) from Taiwan, but can be easily distinguished from them by the following combination of characters.

- 1: The ground color on the underside of both wings is uniformly whitish, whereas in *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER) it is more or less brownish.
- 2: On the underside of the forewing the discal spots are not followed by a dark shade on

their outer side, whereas in *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER) they are followed by such coloration in the postdiscal area.

On the underside of the hindwing, the orange color is fully developed in space 2 in association with the submarginal crescent marking, but absent in space 3, whereas in *T. zuthus* (LEECH) the orange color is fully developed in spaces 2 and 3, whereas in *T. hainani* (BETHUNE-BAKER) the color is completely absent.

- 4: On the underside of the hindwing there is no spot near base of space 1c adjoining the spot mid cell, which is present in both *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER). The $\sigma\sigma$ genitalia are very much different: the inner harpe of valva is much longer than in *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER), and the posterior angle of the valva is smooth, and without the hook or spine which is present in *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER).

This new species is similar to *Tongeia potanini* (ALPHERAKY, 1889) in that the σ genitalia have an elongate and sharply pointed inner harpe to the valva, but can be easily distinguished from the latter by the following combination of characters.

- 1: On the underside of the forewing, the upper discal spots in spaces 3-6 are well separated, not fused into a single smooth band as in *T. potanini* (ALPH.), and the lower discal spots in spaces 1b and 2 are well separated too, not conjoined as in *T. potanini* (ALPH.).
- 2: On the underside of both wings the submarginal crescents are well separated, not conjoined as a band as in *T. potanini* (ALPH.), and the marginal spots are well separated too, not conjoined as in *T. potanini* (ALPH.).
- 3: On the underside of the hindwing the discal spot in space 6 is placed inside of that in 7, the two spots are well separated and are not conjoined as in *T. potanini* (ALPH.), and are directed towards the discocellular bar, not at the anal angle of hindwing as in *T. potanini* (ALPH.).
- 4: The σ genitalia are different: valva is rather broad at caudal 1/3, not tapered from the base to the caudal end and phallus not tapered near the tip as in *T. potanini* (ALPH.).

Discussion: Although the new species is very similar to *T. zuthus* (LEECH) and *T. hainani* (BETHUNE-BAKER) in its external features, it is not closely allied because the species does not share any important genital structures. It is very possible that *T. potanini* (ALPH.) is the most closely allied species of *T. dongchuanensis* spec. nov. in phylogeny, because both share the following distinctive characters which are not found in other species: 1. the underside of the hindwing is without a spot near the base of space 1c adjoining the spot mid cell, 2. the inner harpe is overlapped with the ampulla and fully developed like a rifle bayonet. [In *T. bella* HUANG, 2001, the inner harpe is also developed like a bayonet, but is well, but well separated from the ampulla at the base, not overlapped by the ampulla as in *T. potanini* (ALPH.) and *T. dongchuanensis* spec. nov.]

It is also possible that *T. dongchuanensis* spec. nov. is similar to *T. potanini* (ALPH.) in nature, as the latter has been found in most areas of N. Yunnan (many specimens have been examined). The very little known *Evers arcana* LEECH, 1890 is only based upon a single σ type specimen from Hubei, Central China and cannot belong to *Tongeia* TUTT because the

butterfly illustrated in the original description has its upperside ground color bluish on the basal half of the forewing.

The recently described *T. shaolinensis* WANG & NIU, which is based upon a single ♀ from Henan, Central China is apparently only a **new junior synonym** of *T. fischeri* (EVERSMANN). According to its original figure and description, *T. shaolinensis* WANG & NIU differs from *T. fischeri* (EVERSMANN) only in that the underside forewing discal spots are in a straighter line and the underside hindwing discal spots are closer to the submarginal spots. However, such variation does not exceed that experienced in *T. fischeri* (EVERSMANN).

Hitherto 14 valid species have been known within *Tongeia* TUTT including the new species, and a checklist is as follows:

- 1: *T. amplifascia* HUANG, 2001; distribution: Nujiang valley at Yunnan-Tibet border. Specimens and ♂ genitalia illustrated in HUANG (2001).
- 2: *T. bella* HUANG, 2001; distribution: Nujiang valley at Yunnan-Tibet border. Specimen and ♂ genitalia illustrated in HUANG (2001).
- 3: *T. confusa* HUANG, 2003; distribution: Dulongjiang valley, NW Yunnan. Specimen and ♂ genitalia illustrated in HUANG (2003).
- 4: *T. davidi* (POUJADE, 1885); distribution: Sichuan. Specimens illustrated in LEECH (1893), SEITZ (1909), LEE & ZHU (1992), D'ABRERA (1993) and WANG & FAN (2002). ♂ genitalia illustrated in WANG & FAN (2002: 324) as *T. filicaudis filicaudis*.
- 5: *T. dongchuanensis* HUANG & CHEN **spec. nov.**; distribution: NE Yunnan. Specimen and ♂ genitalia here illustrated.
- 6: *T. filicaudis filicaudis* (PRYER, 1877); distribution: C. and S. China, Shaanxi and Sichuan. Specimens illustrated in various journals. ♂ genitalia illustrated in WANG & FAN (2002: 327) as *T. davidi* (POUJADE, 1885).
- 6 a: *T. filicaudis mushana* (TANIKAWA, 1940); distribution: Taiwan. Specimen and ♂ genitalia illustrated in SHIROZU (1960).
- 7: *T. fischeri fischeri* (EVERSMANN, 1843); type locality: Spasskoe, Russia; distribution: Russia, Japan, Korea, China. Specimens illustrated in various journals. ♂ genitalia illustrated in KAWAZOE & WAKABAYASHI (1979: 170) and WANG & FAN (2002: 323).
- 7 a: *T. fischeri caudalis* (BRYK, 1946); type locality: Korea.
- 7 b: *T. fischeri japonica* FUJIOKA, 1975; type locality: Japan.
- 7 c: *T. fischeri sachalinensis* (MATSUMURA, 1925); type locality: Sachalin Is.
- 7 d: *T. fischeri thyestes* (SEITZ, [1923]); type locality: not provided in original description.
- 7 e: *T. fischeri shaolinensis* WANG & NIU, 2002; type locality: Henan, Central China. **syn. nov.**

Notes: The subspecific division is questionable and not widely accepted. The taxon *T. fischeri thyestes* (SEITZ) was illustrated and described simply by SEITZ under the term of „*fischeri*“ to have distinct dark dots on underside „joined to strigiform chains“ The name „*thyestes*“ was attributed to FRUHSTORFER in SEITZ, but according to ICZN it should be attributed to SEITZ because he was the author of the description whilst FRUHSTORFER did not publish any text on this subject. The name „*thyestes*“ was incorrectly listed under *T. kala* (DE NICEVILLE) in BRIDGES' catalogue just because SEITZ wrongly treated „*kala*“ as a geographical variation of *T. fischeri* (EVERSM.) and „*kala*“ was listed

- under the term of *T. fischeri* (EVERSM.) and followed by „*thyestes*“ in SEITZ.
- 8: *T. hainani* (BETHUNE-BAKER, 1914); distribution: Taiwan, Hainan and Fujian. Specimens illustrated in SHIROZU (1960), D'ABRERA (1986), LEE & ZHU (1992) and CHOU & al. (1994). ♂ genitalia illustrated in SHIROZU (1960:343).
 - 9: *T. ion ion* (LEECH, 1891); distribution: Sichuan and N. Yunnan. Specimens illustrated in LEECH (1893), SEITZ (1909), LEE & ZHU (1992), D'ABRERA (1993), HUANG (1998 & 2001) and WANG & FAN (2002). ♂ genitalia illustrated in HUANG (1998 & 2001) and WANG & FAN (2002: 327).
 - 9a: *T. ion cellariusi* (BOLLOW, 1930); distribution: Gansu and Shaanxi. Specimens illustrated in CHOU & al. (1994).
 - 9 b: *T. ion cratylus* FRUHSTORFER, 1915; distribution: Sichuan-Tibet border & E. Tibet. Specimen and ♂ genitalia illustrated in HUANG (2001).
 - 10: *T. kala* (DE NICEVILLE, 1890); distribution: Naga Hills and Khasia Hills, NE India; S. Shan States, Burma. Specimen illustrated in EVANS (1932, plate 27). ♂ genitalia illustrated in CANTILE (1964).
Notes: According to CANTILE (1962) the two valvae of *T. kala* (DE NICEVILLE) were wrongly shown in CANTILE (1964) and the valva on the left should be placed on the right of the other, thus the valva should be pointed outwards and bifid at the apex in the ventral view. This bifid apex is unique and not found in other species of *Tongeia*.
 - 11: *T. menpae menpae* HUANG, 1998; distribution: Motuo and Tongmai, SE Tibet. Specimens and ♂ genitalia illustrated in HUANG (1998).
 - 11 a: *T. menpae pseudoion* HUANG, 1998; distribution: Linzhi and Milin, SE Tibet. Specimens and ♂ genitalia illustrated in HUANG (1998).
 - 12: *T. potanini potanini* (ALPHERAKY, 1889); distribution: China and Indo-China. Specimens illustrated in various literatures. ♂ genitalia illustrated in WANG & FAN (2002).
 - 12 a: *T. potanini umbriel* DOHERTY, 1889; distribution: N. Burma. Specimen illustrated in EVANS (1932, plate 27).
 - 12 b: *T. potanini glycon* CORBET, 1940; distribution: Peninsular Malaya and Langkawi. Specimen illustrated in CORBET & PENDLEBURY (1978: plate 20) & D'ABRERA (1986: 650).
 - 13: *T. pseudozuthus* HUANG, 2001; distribution: Chayu, SE Tibet. Specimens and ♂ genitalia illustrated in HUANG (2001).
 - 14: *T. zuthus* (LEECH, [1893]); distribution: Sichuan. Specimen illustrated in LEECH (1893), SEITZ (1909), LEE & ZHU (1992), D'ABRERA (1993), CHOU & al. (1994), HUANG (1998). ♂ genitalia illustrated in HUANG (1998 & 2001).

To distinguish the new species from all others, a key is provided as follows. The authors have examined specimens of almost all species, except only *T. kala* (DE NICEVILLE) from Burma, which is known to us only from the literature.

Key to species of *Tongeia* TUTT

1. Upperside forewing discocellular bar clearly marked in pale blue; underside hindwing all submarginal spots broadly associated with orange coloring; inner harpe of valva separated from ampulla deeply at base, not overlapped with ampulla—————*bella*

- Upperside forewing discocellular bar absent, at most bearing some blue scales; underside hindwing without orange coloring in submarginal area or at most only lower submarginal spots broadly associated with orange coloring; inner harpe of valva overlapped with ampulla ————— 2
2. Underside of hindwing without a spot near base of space 1c adjoining the spot mid cell; inner harpe developed like a long and straight sword bayonet pointing posteriorly— 3
Underside of hindwing with a spot near base of space 1c adjoining the spot mid cell; inner harpe not developed like a long and straight rifle bayonet pointing posteriorly- 4
3. Underside forewing discal spots in spaces 3-6 fused into a straight and smooth band; underside hindwing discal spots in spaces 6 and 7 conjoined and directed to anal angle of hindwing; valva tapered from base to tip in lateral view, phallus tapered near tip ————— *potanini*
Underside forewing discal spots in spaces 3-6 well separated; underside hindwing discal spots in spaces 6 and 7 well separated and directed to the discocellular bar; valva not tapered from base to tip in lateral view, phallus stout and downcurved near tip ————— *dongchuanensis*
4. Valva bifid at apex ————— *kala*
Valva not bifid at apex ————— 5
5. Discal spots on both wings uniform black; inner harpe obtuse at tip ————— 6
Discal spots blacker on forewing but more or less paler on hindwing; inner harpe pointed at tip ————— 8
6. Forewing underside without black spot in cell ————— *fischeri*
Forewing underside with black spot in cell ————— 7
7. On underside of forewing the submarginal markings are comparatively narrower, underside ground color more whitish; distal margin of ampulla serrate ————— *dauidi*
On underside of forewing the submarginal markings are comparatively broader, underside ground color more or less brownish; distal margin of ampulla smooth ————— *filicaudis*
8. Distal margin of ampulla more or less upright to posterior margin of valva ————— 9
Distal margin of ampulla considerably oblique to posterior margin of valva ————— 11
9. Valva gently curved at anterior angle, harpe not needle-like at tip ————— *hainani*
Valva strongly produced at anterior angle, harpe needle-like at tip ————— 10
10. Submarginal spots in spaces 2 and 3 on underside of hindwing broadly associated with orange coloring; harpe shorter and not exceeding the caudal end of ampulla ————— *zuthus*
Submarginal spots in spaces 2 and 3 on underside of hindwing not associated with orange coloring; harpe longer and exceeding the caudal end of ampulla ————— *menpae*
11. Size of male genitalia bigger, valva and inner harpe relatively longer ————— *ion*
Size of male genitalia smaller, valva and inner harpe relatively shorter ————— 12
12. Discal spot in space 4 of forewing underside nearly horizontal to forewing dorsum; distal margin of ampulla strongly serrate ————— *amplifascia*
Discal spot in space 4 of forewing underside only slightly oblique; distal margin of ampulla smooth or minutely serrated ————— 13
13. Subbasal spots on hindwing underside mostly black; harpe remarkably narrow at middle, valva longer ————— *pseudozuthus*
Subbasal spots on hindwing underside brown; harpe remarkably broader at middle, valva shorter ————— *confusa*

References

- BRIDGES, C. A. (1988): Catalogue of Lycaenidae and Riodinidae. - Urbana, Illinois.
- CANTILE, K. (1962): The Lycaenidae portion (except the *Arhopala* group) of Brigadier EVANS' The Identification of Indian butterflies, 1932. - Bomb. Nat. His. Soc., Bombay.
- CANTILE, K. (1964): Genitalia of the butterfly genera *Surendra* MOORE and *Everes* HÜBNER. - J. Bomb. nat. Hist. Soc. **61** (1): 210-212, 17 figs., Bombay. [Apr. 1964]
- CHOU, I (Editor) (1994): Monographia Rhopalocerorum Sinensium. - Henan Science and Technology.
- CORBET, A. S. & H. M. PENDLEBURY (1978): The butterflies of the Malay Peninsula. 3rd edition revised by J. N. ELIOT. - Kuala Lumpur.
- D'ABRERA, B. (1986): Butterflies of the Oriental Region.3. - Hill House, Melbourne.
- D'ABRERA, B. (1993): Butterflies of the Holarctic Region.3. - Hill House, Melbourne.
- Draeske, J. (1925): Die Schmetterlinge der Stotznernschen Ausbeute. 2-4. - Dt. Ent. Z. Iris. 38: 1-8; 39: 48-57, 211-231, Dresden.
- EVANS, W. H. (1932): The identification of Indian butterflies. 2nd edition. - Madras.
- HUANG, H. (1998): Research on the butterflies of the Namjagbarwa region, S.E.Tibet. - Neue Ent. Nachr. 41:207-263, Marktleuthen.
- HUANG, H. (2001): Report of H.HUANG's 2000 Expedition to SE Tibet for Rhopalocera. - Neue Ent. Nachr. **51**: 65-151, Marktleuthen.
- HUANG, H. (2003): A list of butterflies collected from Nujiang and Dulongjiang, China with descriptions of new species, new subspecies and revisional notes. - Neue Ent. Nachr. **55**: 3-114, 160-177, Marktleuthen..
- KAWAZOE, A. & M. WAKABAYASHI (1979): Coloured illustrations of the butterflies of Japan. - Osaka.
- LEE, C.-L. & B.-Y. ZHU (1992): Atlas of Chinese butterflies. - Shanghai.
- LEECH, J.H. (1892-1894): Butterflies from China, Japan and Corea. - London.
- MONASTYRSKII, A.L. & A.L. DEVYATKIN (2003): Butterflies of Vietnam (systematic list). - Moscow.
- OSADA, S., UEMURA, Y. & J. UEHARA (1999): An illustrated checklist of the butterflies of Laos P.D.R. - Mokuyo-sha, Tokyo, 240 pp. (143 pls.)
- SEITZ, A. (Editor) (1909): Macrolepidoptera of the world. Vol. 1. The Palaearctic Butterflies. - Alfred Kern, Stuttgart.
- SEITZ, A. (1927): Macrolepidoptera of the world. Vol. 9. The Indo-Australian Butterflies. - Alfred Kern, Stuttgart.
- SHIROZU, T. (1960): Butterflies of Formosa in colour. - Osaka.
- TONG, X.-S., QIAN, Z.-Q & L.-S. WANG (1993): Butterfly fauna of Zhejiang [in Chinese]. Zhejiang Science & Technology.
- WANG, M. & X.-L. FAN (2002): Butterflies Fauna Sinica: Lycaenidae. - Henan Science & Technology, Zhengzhou.
- WANG, Z.-G. & Y. NIU (2002): New species of butterflies from China 2. - Entomotaxonomia **24** (4): 276-284, Yangling.
- WANG, Z.-G., NIU, Y. & D.-H. CHEN (1998): Insect fauna of Henan Lepidoptera: Butterflies




Fig. 1: σ genitalia of *Tongeia dongchuanensis* **spec. nov.** consisting of dorsum in dorsal view (D), of uncus and brachium in lateral view (U+B), of phallus in lateral view (P), of right valva in inner lateral view (VL), of juxta in lateral view (J), and of valvae in ventral view (VV).

Colour plate 4/ Farbtafel 4, p. 284

Fig. G: *Tongeia dongchuanensis* **spec. nov.**, Holotype σ upperside (left half) and underside (right half).

Colour plate 3/ Farbtafel 3 (p.282)

HUANG, H. & K. SONG: New or little known elfin lycaenids from Shaanxi, China (Lepidoptera: Lycaenidae). - Atalanta (August 2006) 37 (1/2): 261-167, Würzburg

Fig. 1: *Ahlbergia luoliangi* **spec.nov.** Holotype ♂ upperside (♂ brand outlined in white on left forewing).

Fig. 2: *Ahlbergia luoliangi* **spec. nov.** Holotype ♂ underside.

Fig. 3: *Ahlbergia luoliangi* **spec.nov.** Paratype ♀ upperside (left half) and underside (right half).

Fig.4: *Ahlbergia leei* JOHNSON, 1992 ♀ (Chang-an, Shaanxi) upperside (left half, with ♀ brand outlined in white) and underside (right half).

Fig. 5: *Ahlbergia leei* JOHNSON, 1992, ♀(Chang-an, Shaanxi) upperside (left half) and underside (right half).

Fig. 6: *Ahlbergia hsui* JOHNSON, 2000, ♂ (Xing-long-sfan, S. Gansu) upperside (left half) and underside (right half).


Fig. 7: *Ahlbergia hsui* JOHNSON, 2000, Holotype ♂ (Kang-xian, S. Gansu, deposited in IZAS) upperside (left half) and underside (right half).

Fig. 8: *Novosatsuma collosa* JOHNSON, 1992, ♂ (Chang-an, Shaanxi) upperside (left half, with ♂ brand outlined in white) and underside (right half).

Fig. 9: *Novosatsuma collosa* JOHNSON, 1992, ♂ (Chang-an, Shaanxi) upperside (left half) and underside (right half).

Fig. 10: Costal area of forewing showing ♂ brand. C: *A. clarofacia* JOHNSON; Le: *A. leei* JOHNSON, 1992; Lu: *A. luoliangi* **spec.nov.**; H: *A. hsui* JOHNSON, 2000; A: *Cissatsuma albilinea* (RILEY, 1939); D: *A. dongyui* HUANG & ZHAN, 2006.

Colour plate 3/ Farbtafel 3



Colour plate 4/ Farbtafel 4




Fig. A: *Ahlbergia dongyui* **spec.nov.** Holotype ♂ upperside (left half, with ♂ brand outlined in white) and underside (right half).

Fig. B: *Ahlbergia dongyui* **spec.nov.** Paratype ♂ upperside.


Fig. C: *Ahlbergia dongyui* **spec.nov.** Paratype ♂ underside.

Fig. D: *Ahlbergia leei* JOHNSON, 1992 ♂ (Chang-an, Shaanxi) upperside (left half, with ♂ brand outlined in white) and underside (right half).

Fig. E: *Ahlbergia confusa* **spec.nov.** Holotype ♀ upperside (left half) and underside (right half).











Fig. F: *Ahlbergia confusa* **spec.nov.** Paratype ♂ upperside (left half, with brand outlined in white) and underside (right half).

Fig. G: *Tongia dongchuanensis* **spec.nov.**, Holotype ♂ upperside (left half) and underside (right half).



Agonopterix ferocella (CHRÉTIEN, 1910)

Colour plate 5/ Farbtafel 5



Colour plate 6/ Farbtafel 6




Fig. 1: *Tatargina* (s. str.) *picta* (WALKER, [1865] 1864), China, Yunnan, Haba mts., Hutiao vill., h=2100 m, 21-26.VI.1996, S. MURZIN leg.

Fig. 2: *Tatargina* (*Hindargina*) *pannosa* (MOORE, 1879), Nepal, Mahakali, Banku, 660 m, 20.VI.1995, anonymous leg.

Fig. 3-4. *Tatargina* (*Hindargina*) *sipahi* (MOORE, 1872), lectotype, India, Bombay, upperside (3) and underside (4).

Colour plate 5/ Farbtafel 5 (p. 285)

Atalanta (September 2006) 37 (1/2): 284-285, Würzburg, ISSN 0171-0079

DUBATOLOV, V. V.: On the generic status of the Afrotropical *Nyctemera* species (Lepidoptera, Arctiidae). - Atalanta (August 2006) 37 (1/2): 191-205, Würzburg.

Fig. 1: *Podomachla antinorii* (OBERTHÜR, 1880), ♂, Tanzania, 5 km S from Bukoba, Forest Kibira, 1.VI.1912, TROITSKII leg. (ZIN).

Fig. 2: *Podomachla antinorii* (OBERTHÜR, 1880), ♀, Cameroun, Bitye Ja River, 2000 ft, X-XI.1910 (MMUM). **Fig. 3:** *Podomachla apicalis* (WALKER, 1854), ♂, South Africa, Natal, Weenen, coll. by G.H. BURN (MMUM). **Fig. 4:** *Chiromachla restricta* (BUTLER, 1894), ♂, Kenya, Kibwezi, 31.VII.1917 (MMUM). **Fig. 5:** *Chiromachla torbeni* (WILTSHIRE, 1983), ♂, holotype, Yemen Arab Republic (N Yemen): Wadi Dhabab, 19.X.1981, TORBEN B. LARSEN leg., from: WILTSHIRE (1983). **Fig. 6:** *Chiromachla torbeni* (WALKER, 1983), ♀, allotype. Yemen Arab Republic (N Yemen): Wadi Annah, 1400 m, 22.V.1980, TORBEN B. LARSEN leg., from: WILTSHIRE (1983). **Fig. 7:** *Chiromachla insulare* (BOISDUVAL, 1833), ♂, [Madagascar], without label (MMUM). **Fig. 8:** *Chiromachla perspicua* (WALKER, 1854), ♂, without label, probably from West Africa (MMUM). **Fig. 9:** *Afronyctemera itokina* (AURIVILLIUS, 1904), ♂, Rwanda, Butare, XII.1976, A. POPOUDINA leg. (SZMN). **Fig. 10:** *Xylecata hemixantha* (AURIVILLIUS, 1904), ♂, Rwanda, Butare, I.1977, A. POPOUDINA leg. (SZMN).

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Atalanta](#)

Jahr/Year: 2006

Band/Volume: [37](#)

Autor(en)/Author(s): Huang Hao, Chen Zhong

Artikel/Article: [A new species of *Tongia* Tu tt. \[1908\] from Northeast Yunnan, China 184-190](#)