

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
<i>IGCP 596 Opening Meeting</i>		Graz, 19-24 th September 2011	

Early to Middle Devonian ostracodes from the Western Dra Valley (Morocco): first eventstratigraphical implication

DOJEN, C.¹, ABOUSSALAM, S.² & BECKER, R.T.²

(1) Landesmuseum Kärnten, Museumsgasse 2, A-9021 Klagenfurt, Austria;
claudia.dojen@landesmuseum.ktn.gv.at

(2) Westfälische Wilhelms-University, Institute for Geology and Palaeontology, Corrensstr. 24, D-48149 Muenster, Germany; *taghanic@uni-muenster.de*; *rbecker@uni-muenster.de*

The Devonian of the Anti-Atlas in southwest Morocco is world-famous for its extensive outcrops that are rich in well-preserved faunas, but those of the Western Dra Valley are still under study and especially the lower Devonian ostracodes are poorly known. We propose first eventstratigraphical results on ostracodes of earliest Emsian to basal Givetian age from the sections Bou Tserfine, Rich Tamelougou and Hassi Mouf near Assa, which are well dated by conodonts. All ostracode associations indicate an offshore position below wave base. Paleobiogeographically, most taxa belong to the Paleotethysian Province, but few North-American taxa are also present and corroborate migration paths between both areas via North Africa.

At present, some hundred ostracode individuals belonging to about 45 benthic taxa have been tentatively identified. Lower Emsian samples with ostracodes are from sections Rich Tamelougou and Bou Tserfine. They yielded taxa such as *Polyzygia kroemmelbeini* and *Bollia azagora*, which are in these sections restricted to the Early Emsian although their global range is longer. But most of the ostracodes are from the lower Upper Emsian *Hollardops* Limestone (basal Khebchia Fm) and the Eifelian *Pinacites* Limestone (lower Yeraifa Fm).

The *Hollardops*-Limestone is a regionally distinctive limestone that has furnished frequently *Caudicriodus culicellus* - *Icriodus corniger ancestralis* conodont assemblages, which indicate early Late Emsian age. Only its basal bed is latest early Emsian as indicated by monospecific occurrences of the *bilatericrescens* conodont assemblage. The ostracode assemblages belong to the so-called "mixed faunas" with only few palaeocopes but rich in metacopes (some of them with small spines), indicating an offshore position probably in deeper and less agitated water below wave base. The *Pinacites* Limestone contains goniatites (e.g., *Pinacites*, *Fidelites*), and conodonts of the *costatus* Zone (Eifelian). The ostracode fauna is partly similar to those of the *Hollardops*-Limestone and long ranging taxa such as *Ulrichia* ex gr. *acricula* and *Jenningsina planocostata* occur throughout the sections. But several taxa as e.g. *Bufina* aff. *bicornuta*, *Bufanchiste bufinoides*, *Polyzygia symmetrica* or *Favulella frankenfeldi* are despite their global ranges restricted here to the Eifelian. Besides these, several spiny taxa such as *Semibolbina*, *Loquitzella*, *Berounella* or *Tricornina* occur, thus, reflecting slightly deeper and calmer water conditions than during the sedimentation of the *Hollardops*-Limestone. So far, Givetian strata with ostracodes have been found in the studied sections only at Hassi Mouf. The occurring taxa are long-ranging and globally widely distributed such as *Jenningsina planocostata* or *Praepilatina*, and the assemblages present low diversities.

As regards Devonian events and their effect on ostracodes both the Kellwasser-Event and the Hangenberg-Event are studied thoroughly. However, the smaller-scale events such as the Daleje, Choteč or Kačák Event have not been studied in detail so far. Without covering the eventhorizons in detail, our study gives nonetheless preliminary information on the possible influence of these events on benthic ostracodes faunas in the W Dra Valley:

- After the Daleje-Event horizon (approx. Early/Late Emsian boundary), eight of thirteen taxa still occur; *Polyzygia vinea* disappears worldwide, whereas *P. kroemmelbeini* disappears locally.
- After the Choteč-Event horizon (approx. Emsian/Eifelian boundary), thirteen of twenty-four taxa still occur; *Jenningsina thuringica* disappears worldwide; *Bufina sotoi* and *Tricornina* ex gr. sp. A survive only in Morocco.

Ber. Inst. Erdwiss. K.-F.-Univ. Graz	ISSN 1608-8166	Band 16	Graz 2011
<i>IGCP 596 Opening Meeting</i>	Graz, 19-24 th September 2011		

- After the Kačák-Event horizon (approx. Eifelian/Givetian) only five from twenty-seven taxa still occur; the five survivors are ubiquitous taxa, such as the long-ranging and globally widely distributed *Jenningsina planocostata*; even some long ranging taxa such as *Ulrichia* ex gr. *fragilis*, *U.* ex, gr. *spinifera* and "*Cyterellina*" *inconstans* disappear locally.

Thus, the Kačák-Event seems to be the most effective one of the minor events as regards the ostracodes of the Western Dra Valley. However, further studies have to include more material covering the event horizons in detail. In addition, the results have to be compared with the studies on Devonian ostracodes from other Moroccan and North African areas as described e.g. by CASIER or by BECKER.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2011

Band/Volume: [16](#)

Autor(en)/Author(s): Dojen Claudia, Aboussalam Sarah, Becker Ralph Thomas

Artikel/Article: [Early to Middle Devonian ostracodes from the Western Dra Valley \(Morocco\): first eventstratigraphical implication. 41-42](#)