

Paleogeography, paleobiogeography and bathymetry based on fish, particularly otoliths

Rostislav BRZOBOHATY

Department of Geology and Palaeontology, Masaryk University Brno, Faculty of Science, Kotlářská 2, 611 37 Brno, Czeckia

Neogene otoliths have to be taxonomically compared with recent species. Otoliths from nearly 6000 Recent species are known (NOLF 1985). For localities with 50-60 species several hundred or thousand kg of sediment is usually needed, therefore samples from drilling cores or boreholes are not reliable for a sufficient interpretation of otolith faunas. All over the world there are only 10 specialists working on otoliths, but the knowledge of the European Neogene basins is quite good. The North Sea Basin belongs to the best investigated basins. In the Aquitanian Basin there is a low level of knowledge in the Serravallian and a very high one in the Langhian and earlier times (Fig. 2).

Ma	EPOCH	Otolith faunas - level of knowledge (investigation)			
		AQUITAINE BASIN		NORTH SEA BASIN	
15-16.4	Middle MIOCENE	SERRAVALLIAN	low	"Saubrigues - Paleocanyon"	LANGENFELDIAN
		LANGHIAN	very high		REINBEKIAN
20-23.8	Early MIOCENE	BURDIGALIAN	very high		HEM MOORIAN
		AQUITANIAN	very high		VIERLANDIAN
		CHATTIAN	very high		NEOCHATTIAN

Fig. 1: Level of knowledge of otoliths in North Sea and Aquitanian Basin.

Ma	EPOCH	Otolith faunas - level of knowledge (investigation)				
		MEDITERRANEAN		CENTRAL PARATETHYS		
5-5.3	Pliocene	GELASIAN	high	Spain (Papirol) Sicily, Calabria SE France (Le Puget) Morocco (Dar-Bel Hamri)	very low	PONTIAN
		PIACENZIAN				
		ZANCLLEAN				
10-11.4	Late MIOCENE	MESSINIAN	high	Torremondo, Moncuco Borelli	low	PANNONIAN
		IURTONIAN	high	Strat. area etc.	intermediate	SARMATIAN
15-16.4	Middle MIOCENE	SERRAVALLIAN	intermediate	Madona della Neve (Piemonte)	high	BADENIAN
		LANGHIAN	low	Janiro Baldissero	intermediate	KARPATIAN
20-23.8	Early MIOCENE	BURDIGALIAN	intermediate	Complesso Termo Fora, Sciolze, Baldissero, Valle Cepi	intermediate	OTTNANGIAN
		AQUITANIAN	low	Moleto Prera	very low	EGGENBURGLIAN
		CHATTIAN			very low	EGERIAN

Fig. 2: Level of knowledge of otoliths in the Mediterranean and in the Central Paratethys.

In the Mediterranean there is a low level of knowledge in the Aquitanian and Langhian, intermediate in Burdigalian and Serravallian and high in Tortonian, Messinian and younger stages. In the Central Paratethys there is a low level of knowledge in the Egerian, Eggenburgian, Pannonian, Pontian, intermediate in Sarmatian, Karpatian, Ottnangian and high in Badenian (Fig. 2). Knowledge of otolith faunas is quite insufficient in the Eastern Paratethys. Otolith associations can be used for bathymetric reconstruction. A generalized approach to the paleobathymetry (neritic – bathyal) can be refined on the basis of the method suggested by NOLF & CAPPETTA (1989) and discussed by NOLF & BRZOBOHATY (1994). This method was used for a paleobathymetric evaluation of the Lower Badenian Carpathian Foredeep (20 localities) in South Moravia (BRZOBOHATY 1997). Four bathymetric associations were proved in the Lower Badenian clay (Fig. 3). Two graphs (Fig. 4) document examples of the deepest association (Brno - Kralovo Pole) and the shallowest one (Kralice n. O.). Results in the map of the Carpathian Foredeep illustrate a proposed paleobathymetry for the Lower Badenian Sea (Fig. 5) and preliminary isobaths (Fig. 6).




Fig. 3: Actual bathymetric repartition of important teleost taxa represented in the Lower Badenian and their reflexion in localities of the Carpathian Foredeep (South Moravia).




Fig. 4: Actual bathymetric repartition of teleost taxa represented in the Lower Badenian from Brno-Kralovo Pole and Kralice n. O.




Fig. 5: Position of the Lower Badenian localities in the Carpathian Foredeep (South Moravia) and the paleobathymetric reconstruction.




Fig. 6: Preliminary isobath lines in the Lower Badenian Sea of the Carpathian Foredeep (South Moravia).

The Lower Badenian Sea flooded all the relief and archibenthic (Macrouridae) and mesopelagic (e.g., Myctophidae) fauna penetrated through deep depressions to the north and

western margin of the basin. Mesopelagic taxa could end up in shallow depths (usually with juvenile specimens, e. g., Kralice n. O.) (Fig. 7).




Fig. 7: a = Lower Badenian Sea (A – macrourids, M – mesopelagic fauna), b = recent Carpathian Foredeep with denudation remnants of Lower Badenian deposits.

Comparison of eastern and western parts in the Lower Badenian Central Paratethys shows that in the western parts there are higher numbers of genera and species in Macrouridae and Myctophidae, while in the eastern part Macrouridae are missing and only a low number of genera and species of Myctophidae is present. The Polish Foredeep and the Romanian part of the Central Paratethys seem to be generally shallower than the western part of the Central Paratethys.

References

- BRZOBOHATY, R., 1997: Paleobathymetry of the Lower Badenian (Middle Miocene, Carpathian Foredeep, South Moravia) based on otoliths. – In: HLADILOVA, S.: Dynamika vztahu marinního a kontinentálního prostředí, Sborník příspěvků, Masarykova universita Brno 1997, 37 – 45, Brno (in Czech, with English resume).
- NOLF, D., 1985: Otolithi Piscium. – In: SCHULTZE, H. P.: Handbook of Palaeoichthyology 10. – 145 pp., Fischer Verlag, Stuttgart, New York.
- NOLF, D. & BRZOBOHATY, R., 1994: Fish otoliths as paleobathymetric indicators. – Paleontologia i evolucion, 24-25 (1992), 255 – 264, Sabadel.
- NOLF, D. & CAPPETTA, H. C., 1989: Otolithes de poissons pliocenens du Sud-Est de la France. – Bull. Inst. Roy. Sci. Nat. Belgique, Sci. Terre, 58, 209 – 277, Bruxelles.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte des Institutes für Geologie und Paläontologie der Karl-Franzens-Universität Graz](#)

Jahr/Year: 2001

Band/Volume: [4](#)

Autor(en)/Author(s): Brzobohaty [Brzobohatý] Rostislav

Artikel/Article: [Paleogeography, paleobiogeography and bathymetry based on fish, particularly otoliths 37-40](#)