TALK

## Geological Transsect Across the Northwestern Himalaya in Eastern Ladakh and Lahul – A Model for the Continental Collision of India and Asia

A. STECK\*, L. SPRING\*, J.-C. VANNAY\*, H. MASSON\*, E. STUTZ\*, H. BUCHER\*, R. MARCHANT\* & J.-C. TIÈCHE\*



## Chronology of the Tertiary Himalayan structures

- 1- Continental collision, underthrusting of India below Asia and initiation of the Nyimaling-Tsarap Nappe (Eocene)
- 2 NE vergent Shikar Beh Nappe
- 3 SW vergent Nyimaling-Tsarap Nappe
- 4 SW vergent "Crystalline Nappe" (MCT and MBT)
- 5 Dextral transpression
- 6 NE vergent folding
- 7 Late extension

(Eocene) (Eocene) (Oligocene-Miocene) (Oligocene) (Miocene) (Miocene)

The detailed geological mapping and structural study of a complete transect across the north western Himalaya, allow to describe the tectonic evolution of the north Indian continental margin during the Tethys ocean opening and the Himalayan Orogeny.

The Late Paleozoic Tethys rifting is associated with several tectonomagmatic events. In Upper Lahul and SE Zanskar, this extensional phase is recorded by Lower Carboniferous synsedimentary transtensional faults, a Lower Permian stratigraphic unconformity, a Lower Permian granitic intrusion and middle Permian basaltic extrusions (Panjal Traps). In eastern Ladakh, a Permian listric normal fault is also related to this phase. The scarcity of synsedimentary faults and the gradual increase of the Permian syn-rift sediments thickness toward the NE, suggest a flexural type margin.

The collision of India and Asia is characterized by a succession of contrasting orogenic phases. South of the Suture Zone, the initiation of the SW vergent Nyimaling-Tsarap Nappe corresponds to an early phase of continental underthrusting. To the S, in Lahul, an opposite underthrusting within the Indian plate is recorded by the NE vergent Tandi Syncline. This structure is associated with the newly defined Shikar Beh Nappe, now partly eroded, which is responsible for the high grade (amphibolite facies) regional metamorphism of South Lahul.

The main thrusting of the Nyimaling-Tsarap Nappe followed the formation of the Shikar Beh Nappe. The Nyimaling-Tsarap Nappe developed by ductile shear of the upper part of the subducted Indian continental margin and is responsible for the progressive regional metamorphism of SE Zanskar, reaching amphibolite facies below the frontal part of the nappe, near Sarchu. In Upper Lahul, the frontal parts of the Nyimaling-Tsarap and Shikar Beh nappes are separated by a zone of

<sup>\*)</sup> Section des Sciences de la Terre, Université de Lausanne, BFSH2, CH-1015 Lausanne, Switzerland





low grade metamorphic rocks (pumpellyite-actinolite facies to lower greenschist facies). At high structural level, the Nyimaling-Tsarap Nappe is characterized by imbricate structures, which grades into a large ductile shear zone with depth. The related crustal shortening is about 87 km.

The root zone and the frontal part of this nappe have been subsequently affected by two zones of dextral transpression and underthrusting: the Nyimaling Shear Zone and the Sarchu Shear Zone. These shear zones are interpreted as consequences of the counterclockwise rotation of the continental underthrusting direction of India relative to Asia, which occured some 45 Ma ago, according to plate tectonic models.

Later, a phase of NE vergent "backfolding" developed on these two zones of dextral transpression, creating isoclinal folds in SE Zanskar and more open folds in the Nyimaling Dome and in the Indus Molasse sediments.

During a late stage of the Himalayan Orogeny, the frontal part of the Nyimaling-Tsarap Nappe underwent an extension of about 15 km. This phase is represented by two type of structures, responsible for the tectonic unroofing of the amphibolite facies rocks of the Sarchu area: the Sarchu high angle Normal Fault, cutting a first set of low angle normal faults, which have been created by reactivation of older thrust planes related to the Nyimaling-Tsarap Nappe.

## REFERENCES

ARGAND, E. 1916: Sur l'Arc des Alpes Occidentales. Eclogae geol. Helv. 14, 145-204.

- FRANK, W., HOINKES, G., MILLER, C., PURTSCHELLER, F., RICHTER, W. & THÖNI, M. 1973: Relations between metamorphism and orogeny in a typical section of the Indian Himalayas. Tscherm. mineral. petrogr. Mitt. 20, 303-332.
- ---, THÖNI & PURTSCHELLER, F. 1977: Geology and petrography of Kulu-South Lahul area. Collog. int. C.N.R.S., 268, 2, 147-172.
- HIRN, A. et al. (12 authors) 1984a: Crustal structure and variability of the Himalayan border of Tibet. Nature (London), 307, 25-27.
- --- et al. (12 authors) 1984b: Lhasa block and bordering sutures--a continuation of a 500 km Moho traverse through Tibet. Nature (London), 307, 23-25.
- MARGERIE, de E. & HEIM Albert, 1888: Les dislocations de l'écorce terrestre. J. Wurster, Zürich. PATRIAT, P. & ACHACHE, J. 1984. India-Eurasia collision chronology has implications for crustal shortening and driving mechanism of plates. Nature (London) 311, 615-621.
- SPRING, L. & CRESPO, A. 1992: Nappe tectonics, extension and metamorphic evolution in the Indian Tethys Himalaya. Tectonophysics, 11, 978-989.
- ----, BUSSY, F., VANNAY, J.C., COSCA, M. & HUON, S. 1992 (in press): Permo-Carboniferous "alkaline" granitic magmatism in the Indian High Himalaya (Upper Lahul - SE Zanskar): geochimical characterization and geotectonic implications. Spec. Publ. geol. Soc. London.

----, STUTZ, E., THELIN, P., MARCHANT, R., MASSON, H. & STECK, A. 1992. Inverse métamorphic zonation in very low-grade Tibetan series of SE Zanskar and its tectonic consequences (Himalaya, NW India). Schweiz. mineral. petrogr. Mitt. 72, in press.

- SRIKANTIA, S.V. & BHARGAVA, O.N. 1982: An outline of the structure of the area between the Rohtang pass in Lahaul an the Indus Valley in Ladakh. Geol. Surv. India, misc. Publ. 41/3, 193-204, Calcutta.
- STUTZ, E. A., 1988: Géologie de la chaîne de Nyimaling aux confins du Ladakh et du Rupshu (NW-Himalaya, Inde). Mém, Géol. (Lausanne) 3.
- --- & STECK, A. 1986: La terminaison occidentale du Cristallin du Tso Morari (Haut-Himalaya; Ladakh méridional, Inde). Eclog. geol. Helv. 79, 253-269.
- STECK, A., SPRING, L., VANNAY, J.-C., MASSON, H., STUTZ, E., BUCHER, H., MARCHANT, R. et TIECHE, J.-C. 1993: Geological transect across the Northwestern Himalaya in eastern Ladakh and Lahul. Eclogae geol. Helv. 86 (in press).
- VANNAY; J.C. & SPRING, L. 1993: The geochemistry of the continental basalts within the Tethyan Himalaya of Lahul-Spiti and SE Zanskar (NW India). Spec. Publ. geol. Soc. London (in press).

## **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Berichte der Geologischen Bundesanstalt

Jahr/Year: 1993

Band/Volume: 43

Autor(en)/Author(s): diverse

Artikel/Article: <u>Geological Transsect Across the Northwestern Himalaya in Eastern</u> Ladakh and Lahul - A Model for the Continental Collision of India and Asia 51-53