

The Krappfeld microflora, Carinthia (Austria): A presumable ETM-2 flora, SEM investigation of palynomorphs.

**Christa-Ch. Hofmann¹, Hans Egger², Reinhard Zetter¹,
Stepan Čorić², Omar Mohamed³**

¹ University of Vienna, Department of Palaeontology, Vienna, Austria

² Geological Survey, Vienna, Austria

³ El Minia University, Department of Geology, El Minia, Egypt

Six samples of the Holzer Formation from a short time outcrop in a now refilled quarry near Pemberg (Krappfeld area, NNE of Klagenfurt, Carinthia) have been examined for palynomorphs using LM and SEM. The stratigraphic age of the section is indirectly indicated as Ypresian, because of the mass occurrence of the dinoflagellate genus *Apectodinium*, that is absent in younger strata. Despite the fact that nannoplankton is not preserved, the overlying nummulitic limestones of the Sittenberg Formation are correlative with the NP12. The terrestrial palynomorph assemblages are characterized by the bulk of Myricaceae, Juglandaceae (*Normapolles*-, post-*Normapolles*, *Engelhardia*-, *Platycarya*-types), and Fagaceae and to lesser extant by megathermal elements, of which some are quite common, whilst others occur only accessorial. Of particular interest are the frequently encountered monocots such as Araceae, Arecaceae, and Restionaceae. Further, eudicot taxa that can be assigned to the Alangiaceae, Anacardiaceae, Avicenniaceae, Calycanthaceae, Chloranthaceae, former Euphorbiaceae, Hamamelidaceae, Malvaceae Mastixiaceae, Olacaceae, Sapotaceae, ?Simaroubaceae, Styracaceae, Theaceae, Thymelaceae and many other families occur in smaller numbers.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Berichte der Geologischen Bundesanstalt](#)

Jahr/Year: 2011

Band/Volume: [85](#)

Autor(en)/Author(s): Hofmann Christa-Charlotte, Egger Hans, Zetter Reinhard, Coric Stjepan, Mohammed Omar

Artikel/Article: [The Krappfeld microflora, Carinthia \(Austria\): A presumable ETM-2 flora, SEM investigation of palynomorphs 91](#)