



Entomofauna

ZEITSCHRIFT FÜR ENTOMOLOGIE

Band 36, Heft 37: 505-512

ISSN 0250-4413

Ansfelden, 2. Januar 2014

A study on the ground beetles in Mazandaran province, northern Iran (Coleoptera, Carabidae)

Najmeh SAMIN & Hamid SAKENIN

Abstract

The fauna of ground beetles (Coleoptera: Carabidae) in Mazandaran province is studied in this paper. Totally 15 species from 13 genera and 9 subfamilies were collected during 2011-2012. Two species *Chlaenius steveni* Quensel and *Carabus cribratus porrectangulus* Gehin are new records for Iran.

Zusammenfassung

Die Laufkäferfauna der Mazandaran Provinz wurde studiert. Insgesamt konnten 15 Arten aus 13 Gattungen und 9 Unterfamilien während der Sammelperiode 2011-2012 erfasst werden. *Chlaenius steveni* Quensel und *Carabus cribratus porrectangulus* Gehin sind Neumeldungen für den Iran.

Introduction

The family Carabidae (Coleoptera) is among the dominant groups of terrestrial predators and includes more than 32,000 species worldwide (BOUSQUET & LAROCHELLE 1993), making it one of the largest families of beetles (OBER 2002). Almost of these insects are powerful predators of agricultural and forest pests and can have efficient role in biological control. Most ground beetles are found in the tropics. More than 30% of species are

arboreal, though in general temperate species are terrestrial, most are also flightless and predatory (LOVEI & SUNDERLAND 1996). They are commonly found under stones, logs, leaves, bark, debris, or foraging on the ground. They run rapidly, but seldom fly. Many species are equipped for digging and burrowing in the soil. Most species are nocturnal and a few are attracted to lights especially at night. They are predaceous on insects, worms, slugs, snails, caterpillars, grubs and maggots (THIELE 1977; STORK 1990; LUFF et al. 1992). Carabid habitat and microdistribution (the precise distribution of one or more kinds of organisms in a microenvironment or in part of an ecosystem) are governed by abiotic and biotic factors such as light, temperature, and humidity extremes, food supply, predator presence and distribution, and life history strategies (THIELE 1979; LOVEI & SUNDERLAND 1996). Also, carabid beetles are increasingly used as taxonomic study group in biodiversity and as bio-indicators in monitoring or site assessment studies for nature conservation purposes (LOREAU 1994; HEIJERMAN & TURIN 1995; DESENDER 1996).

The fauna of Iranian Carabidae was studied rather well. Some important studies on Iranian Carabidae were made by Jaeger (1990, 1992), HEJKAL (2000), LASSALLE (2001), HEINZ (2002), MAGRINI & PAVESI (2003), MOHAMMADZADEHFARD & HOJJAT (2005), JASKULA (2007), TOLEDANO & MARGGI (2007), MOHAMMADZADEH FARD (2008), GHAHARI et al. (2009 a, b, c, 2010), SAMIN et al. (2011), HASANISAADI & SADEGHI NAMAGHI (2011), REZAEI NODEH et al. (2012), ATAMEHR (2013), SALARI GOUGHERI et al. (2013). The aim of this research is to determining of carabids in Mazandaran province (northern Iran).

Materials and Methods

The specimens were collected from different regions of Mazandaran province. Many plastic pitfall traps, 8.5×10 cm (diameter × depth), were installed at 100 m intervals in different regions and were part-filled with mixture of Zolon and water. The traps were emptied weekly and the fallen beetles were collected and identified. The majority of carabid beetles are relatively easily collected in a more or less standardized way by means of pitfall trapping. Nevertheless, much discussion remains on the necessary methodologies in sampling as well as in data analyses or in diversity assessment (SOUTHWOOD 1978; DESENDER 1996). In addition to the pitfall traps, sweepings were conducted randomly in different regions. The nomenclature of the Carabidae is given sensu TRAUTNER & GEIGENMÜLLER (1987) and LÖBL & SMETANA (2003).

Results

In total 15 species from 13 genera and 9 subfamilies of Carabidae were collected and identified from some regions of Mazandaran province. The list of species is given below.

Subfamily **B e m b i d i n a e**

Genus ***Bembidion*** LATREILLE, 1802

Bembidion dalmatinum haupti REITTER, 1908

Material examined: Mazandaran province: Noor (8 m), 2♂♂, September 2011.

Subfamily **B r a c h i n i a e**

Genus *Brachinus* WEBER, 1801

***Brachinus cruciatus* QUENSEL, 1806**

Material examined: Mazandaran province: Savadkooh (543 m), 1♂, July 2012.

Subfamily **C a l l i s t i n a e**

Genus *Chlaenius* BONELLI, 1810

***Chlaenius steveni* QUENSEL, 1806**

Material examined: Mazandaran province: Babol (27 m), 1♀, August 2011. *New record for Iran.*

Subfamily **C a r a b i n a e**

Genus *Carabus* LINNEAUS, 1758

***Carabus cribratus porrectangulus* GEHIN, 1885**

Material examined: Mazandaran province: Sari (10 m), 2♀♀, April 2012. *New record for Iran.*

Subfamily **C i c i n d e l i n a e**

Genus *Cicindela* LINNEAUS, 1758

***Cicindela (Cicindela) rhodoterena* TSCHITSCHITSCHERINE, 1903**

Material examined: Mazandaran province: Kiasar (1082 m), 2♀♀, April 2011.

Subfamily **H a r p a l i n a e**

Genus *Harpalus* LATREILLE, 1802

***Harpalus cupreus* DEJEAN, 1829**

Material examined: Mazandaran province: Qaemshahr (9 m), 2♂♂, September 2011.

***Harpalus saxicola* DEJEAN, 1829**

Material examined: Mazandaran province: Noor (8 m), 1♀, September 2011.

Genus *Ophonus* DEJEAN, 1821

***Ophonus nitidulus* STEPHENS, 1828**

Material examined: Mazandaran province: Nooshahr (48 m), 1♀, 1♂, August 2012.

Genus *Stenolophus* STEPHENS, 1827

***Stenolophus steveni* KRYNICKI, 1832**

Material examined: Mazandaran province: Nooshahr (48 m), 3♀♀, August 2012. Mazandaran province: Kiasar (1082 m), 1♀, 2♂♂, April 2011.

Subfamily **L e b i i n a e**

Genus *Cymindis* LATREILLE, 1806

***Cymindis vaporarium* (LINNAEUS, 1758)**

Material examined: Mazandaran province: Savadkooh (550 m), 2♂♂, July 2012. Mazandaran province: Sari (10 m), 2♀♀, 3♂♂, April 2012.

Genus *Drypta* LATREILLE, 1796

***Drypta dentata* (ROSSI, 1790)**

Material examined: Mazandaran province: Amol (196 m), 1♀, 1♂, July 2012.

Genus *Parazuphium* JEANNEL, 1942

***Parazuphium chevrolati* (CASTELNAU, 1833)**

Material examined: Mazandaran province: Babol (23 m), 1♀, 1♂, August 2011; Amol (196 m), 1♀, July 2012.

Subfamily **N e b r i i n a e**

Genus *Leistus* FRÖLICH, 1799

***Leistus (Leistus) caucasicus* CHAUDOIR, 1867**

Material examined: Mazandaran province: Sari (12 m), 3♀♀, April 2012.

***Leistus (Pogonophorus) hermonis* PIOCHARD DE LA BRÛLERIE, 1875**

Material examined: Mazandaran province: Qaemshahr (9 m), 2♂♂, September 2011.

Subfamily **Pterostichinae**

Genus *Zabrus* CLAIRVILLE, 1806

***Zabrus spinipes* (FABRICIUS, 1798)**

Material examined: Mazandaran province: Savadkooch (543 m), 1♀, July 2012.

Discussion

This research shows that in despite of several conducted researches on the fauna of Carabidae in different regions of Iran, there are unknown species still. Continuing these faunistic researches will result to new data on species diversity and also distribution of Iranian Carabidae. Of course there are some problems in studying on the diversity of Carabidae. One of them is to assess which part of the collected species in a certain locality actually belongs to the local fauna and has reproducing populations. Another problem is the lack of knowledge on year-to-year variation in numbers of many carabid species, in other words data on the magnitude of population dynamics in more or less natural situations. Such studies of course require a continuous long term sampling program, which is probably the most obvious reason for their scarcity (DESENDER 1996). Additionally, although pesticides affect destructively on the survival of carabid beetles, but these beneficial insects can recover their population density gradually and can have efficient role in biological control of agricultural and forest pests (GHAHARI et al. 2005). Therefore, training the farmers in order to applying the safe or at least low-dangerous strategies under IPM (integrated pest management) is necessary for conservation of carabid beetles.

Acknowledgements

The authors are grateful to M. Kesdek and S.S. Avgin of Turkey, and J. Muilwijk of the Netherlands for determining of some specimens and providing the necessary papers. The research was supported by Islamic Azad University (Young Researchers Club and Elites, and Qaemshahr Branch).

References

- ATAMEHR A. (2013): Ground beetles (Coleoptera: Carabidae) of Azarbaijan, Iran. – Turkish Journal of Zoology **37**: 188-194.
- BOUSQUET Y. & A. LAROCHELLE (1993): Catalogue of the geadephaga (Coleoptera: trachypachidae, rhyssodidae, carabidae including Cicindelini) of America north of Mexico. – Memoirs of Entomological Society of Canada **167**: 1-397.
- DESENDER K.R.C. (1996): Diversity and dynamics of coastal dune carabids. – Annales Zoologici Fennici **33**: 65-75.

- GHAHARI H., SAKENIN H. & H. OSTOVAN (2005): Bioeffect of three insecticides, Diazinon, Etrifos, and Chlorpyrifos on density and activity of non-target arthropods. – *J. of Agriculture and Natural Resources Science* 12(4): 125-137 [In Persian with English Summary].
- GHAHARI H., KESDEK M., SAMIN N., OSTOVAN H., HAVASKARY M. & S. IMANI (2009a): Ground beetles (Coleoptera: Carabidae) of Iranian cotton fields and surrounding grasslands. – *Munis Entomology & Zoology* 4 (2): 436-450.
- GHAHARI H., JEDRYCZKOWSKI W.B., KESDEK M., OSTOVAN H. & M. TABARI (2009b): Ground beetles (Coleoptera: Carabidae) from rice fields and surrounding grasslands of Northern Iran. – *Journal of Biological Control* 23 (2): 105-109.
- GHAHARI H., SAKENIN H. & M. KESDEK (2009c): Ground beetles (Coleoptera: Carabidae) fauna and dominant species in fruit orchards of Mazandaran province, northern Iran. Proceedings of the Third International Symposium on Biological control of Arthropods, February 8-13, 2009, Christchurch, New Zealand, p. 597.
- GHAHARI H., AVGIN S.S. & H. OSTOVAN (2010): Carabid beetles (Coleoptera: Carabidae) collected from different ecosystems in Iran with new records. – *Türkiye Entomoloji Dergisi* 34 (2): 179-195.
- HASANISAADI, S. & H. SADEGHI NAMAGHI (2011): Species diversity of carabid beetles (Coleoptera: Carabidae) inhabiting agroecosystems in Kerman province. – Proceedings of 2nd Iranian Pest Management Conference (IPMC), Shahid Bahonar University of Kerman, pp. 77.
- HEIJERMAN T. & H. TURIN (1994): Towards a method for biological assessment of habitat quality using carabid samples (Coleoptera, Carabidae). – In: DESENDER, K., DUFRÈNE, M., LOREAU, M., LUFF, M.L. & MAELFAIT, J.-P. (eds.), *Carabid Beetles: Ecology and Evolution*: 305-312. Kluwer Academic Publishers, Dordredt, Boston, London.
- HEINZ W. (2002): Beschreibung einer neuen *Carabus (Lamprostus)*-Art aus Persien (Coleoptera: Carabidae). – *Zoology in the Middle East* 26: 157-162.
- HEJKAL J. (2000): *Amara (Amara) elborzensis* sp. n. (Coleoptera: Carabidae) from Iran. – *Klapalekiana*. č. 36 pp.
- JAEGER B. (1990): Zur Verbreitung von *Bradycellus csikii* Laczo, 1912 und *B. distinctus* Dejean, 1829 sowie Beschreibung von *Bradycellus heinzei* n.sp. aus dem Nord-Iran (Col., Carabidae). – *Entomologische Nachrichten und Berichte* 34: 9-13.
- JAEGER B. (1992): Beitrag zur Erforschung der Acupalpus-Fauna des Iran und der Türkei nebst Beschreibung von *Acupalpus turcicus* n.sp. (Col., Carabidae). – *Entomologische Nachrichten und Berichte* 36: 223-230.
- JASKULA R. (2007): First record of *Lebia trimaculata* from Iran (Coleoptera: Carabidae). – *Entomological Problems* 37 (1-2): 48.

- LÖBL I. & A. SMETANA (2003): Catalogue of Palaearctic Coleoptera. Volume I. Archostemata-Myxophaga-Adephaga. – Apollo Books. Stenstrup, Denmark, 819 pp.
- LOREAU M. (1994): Ground beetles in a changing environment: determinants of species diversity and community assembly. – In: BOYLE, T.J.B. & BOYLE, C.E.B. (eds.), Biodiversity, Temperate Ecosystems, and Global Change: 77-98. Springer-Verlag, Berlin, Heidelberg.
- LOVEI G.L. & K.D. SUNDERLAND (1996): Ecology and Behaviour of Ground Beetles (Coleoptera: Carabidae). – Annual Review of Entomology **41**: 231-256.
- LUFF M.L., EYRE M.D. & S.P. RUSHTON (1992): Classification and prediction of grassland habitats using ground beetles (Coleoptera, Carabidae). – Journal of Environmental Management **35**: 301-315.
- MAGRINI P. & M. PAVESI (2003): A new Iranian species of *Neoblemus* (Coleoptera Carabidae). – Bollettino della Societa Entomologica Italiana **135** (2): 71-78.
- MOHAMMADZADEH FARD S. (2008): Description of female *Axonya farsica* (Coleoptera: Carabidae: Broscinae). – Journal of Entomological Society of Iran (Supplement) **27** (2): 23-24.
- MOHAMMADZADEH FARD S. & S.H. HOJJAT (2005): New records on Carabid beetles in Iran. – Journal of Entomological Society of Iran **24** (2): 144.
- OBER K.A. (2002): Phylogenetic relationships of the carabid subfamily Harpalinae (Coleoptera) based on molecular sequence data. – Molecular Phylogenetics and Evolution **24**: 228-248.
- REZAEI NODEH M., AFSHARI A., YAZDANIAN M. & GH. A. ASSADEH (2012): Biodiversity of the ground beetles (Coleoptera: Carabidae) in orchard ecosystems of Azadshahr region, Golestan province, Iran. – Journal of Entomological Society of Iran **32** (1): 59-79.
- SALARI GOUGHERI M., HOSSEINI R. & J. HAJIZADEH (2013): A faunal study on the subfamilies Harpalinae and Lebiinae (Coleoptera: Carabidae) in Guilan province, Iran. – Entomofauna **34** (15): 193-200.
- SAMIN N., SAKENIN H., KESDEK M. & S. IMANI (2011): Ground beetles (Coleoptera: Carabidae) from some regions of Iran. – Linzer biologische Beitrage **43/1**: 873-880.
- SOUTHWOOD T.R.E. (1978): Ecological methods with particular reference to the study of insect populations. – Chapman and Hall, London.
- STORK N.D. (1990): The role of Ground Beetles in Ecological and Environmental Studies. Andover: Intercept 424 pp. Sunderland, K. and Vickerman, G.P. 1980: Aphid feeding by some polyphagous predators in relation to aphid density in cereal fields. – Journal of Applied Ecology **17**: 389-396.
- THIELE H.U. (1977): Carabid beetles in their environments. – Springer-Verlag, Berlin.

- THIELE H.U. (1979): Relationships between annual and daily rhythms, climatic demands and habitat selection in carabid beetles. In: ERWIN T.L., BALL G.E. & D.R. WHITEHEAD (eds.). Carabid beetles: Their evolution, natural history, and classification. – W. Junk, The Hague.
- TOLEDANO L. & W. MARGGI (2007): A new *Bembidion* subg. *Trepanes* Motschulsky, 1864 from Iran (Coleoptera: Carabidae). – Atti della Società italiana di Scienze naturali e del Museo Civico di Storia naturale di Milano **148** (I): 127-134.
- TRAUTNER J. & K. GEIGENMÜLLER (1987): Tiger beetles, ground beetles. Illustrated key to the Cicindelidae and Carabidae of Europe. – Margraf, Aichtal, 488 pp.

Authors addresses:

Najmeh SAMIN
Young Researchers Club and Elites, Science and Research Branch
Islamic Azad University
Tehran, Iran
E-mail: n_samin63@yahoo.com

Hamid SAKENIN
Department of Plant Protection, Qaemshahr Branch
Islamic Azad University
Mazandaran, Iran
E-mail: hchelave@yahoo.com

Druck, Eigentümer, Herausgeber, Verleger und für den Inhalt verantwortlich:
Maximilian SCHWARZ, Konsulent f. Wissenschaft der Oberösterreichischen Landesregierung, Eibenweg 6, 4052 Ansfelden, Austria. E-Mail: maximilian.schwarz@liwest.at.

Redaktion: Erich DILLER, ZSM, Münchhausenstraße 21, 81247 München;
Roland GERSTMEIER, Lehrstuhl f. Tierökologie, Hans-Carl-von-Carlowitz-Platz 2, 85350 Freising;
Fritz GUSENLEITNER, Lungitzerstr. 51, 4222 St. Georgen/Gusen, Austria;
Wolfgang SPEIDEL, MWM, Tengstraße 33, 80796 München;
Thomas WITT, Tengstraße 33, 80796 München.

Adresse: Entomofauna, Redaktion und Schriftentausch c/o Museum Witt, Tengstr. 33, 80796 München, Deutschland, E-Mail: thomas@witt-thomas.com; Entomofauna, Redaktion c/o Fritz Gusenleitner, Lungitzerstr. 51, 4222 St. Georgen/Gusen, Austria, E-Mail: f.gusenleitner@landesmuseum.at

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Entomofauna](#)

Jahr/Year: 2015

Band/Volume: [0036](#)

Autor(en)/Author(s): Samin Najmeh, Sakenin Hamid

Artikel/Article: [A study on the ground beetles in Mazandaran province, northern Iran \(Coleoptera, Carabidae\) 505-512](#)