SHORT NOTE

SHORT NOTE 191

Wiebelsheim (Aula). CRNOBRNJA-ISAILOVIĆ, J. & ALEKSIĆ, I. (1999): First record of Coluber najadum EICHWALD (1831) in Serbia.- Arch. Biol. Sci., Belgrade; 51 (3): 47P-48P. DIMOVSKI, A. (1963): Herpetofauna na skopska kotlina. I - zoogeografski i ekološki pregled.- Godišen zbornik Prirodno-matematičkog fakulteta, Univerziteta u Skoplju, Skoplje; knjiga 14, Biologija 2: 189-221. DIMOVSKI, A. (1966): Herpetofauna na skopska kotlina. II - faunistički del.- Godišen zbornik Prirodno-matematičkog fakulteta, Univerziteta u Skoplju, Skoplje; knjiga 16, Biologija 4: 179-188. DžUKIĆ, G. (1972): Herpetološka zbirka Prirodnjačkog muzeja u Beogradu. (Herpetological collection of the Belgrade museum of natural history).- Glasnik Prirodnjačkog muzeja, Beograd; (Ser. B) 27: 165-180. DŽUKIĆ, G. (1995): Diverzitet vodozemaća (Amphibia) i gmizavaca (Reptilia) Jugoslavije, sa pregledom vrsta od međunarodnog značaja; pp. 449-469. In: STEVANOvić, V. & VASIć, V. (eds): Biodiverzitet Jugoslavije. Beograd (Biološki fakultet & Ecolibri), 562 pp. DŽUKIĆ, G. & PASULJEVIĆ, G. (1979): O rasprostranjenju ljuskavog guštera - Algyroides nigropunctatus (DU-MÉRIL & BIBRON, 1839) Reptilia, Lacertidae.- Biosistematika, Beograd; 5 (1): 61-70. KARAMAN, S. (1931): Zoološke prilike skopske kotline.- Glasnik skopskog naučnog društva, Skoplje; knjiga 10, Odeljenje prirodnih nauka, 4: 1-16. KARAMAN, S. (1939): Über die Verbreitung der Reptilien in Jugoslavien.- Ann. Mus. Serbiae Meridionalis, Skoplje; 1 (1): 1-20. MATVEJEV, S. D. (1961): Biogeografija Jugoslavije. Biološki institut NR Srbije, Monograph. 9. Beograd (Naučna knjiga), 232 pp. MATVEJEV, S. D. & PUNCER, I. J. (1989): Karta bioma, predeli Jugoslavije i njihova zaštita --Glasnik Prirodnjačkog Muzeja, Beograd; Posebna izdanja 36: 1-76. PASULJEVIĆ, G (1968): Prilog poznavanju herpetofaune Kosova.- Zbornik Filozofskog fakulteta, Priština; 1: 61-74. RADOVANOVIĆ, M. (1951): Vodozemci i gmizavci naše zemlje. Beograd (Naučna knjiga), 250 pp. RADOVANOVIĆ, M. (1964): Die Verbreitung der Amphibien und Reptilien in Jugoslawien .-Senckenbergiana biol., Frankfurt a. Main; 45 (3/5): 553-561.

KEY WORDS: Reptilia: Testudines: Testudinidae: *Testudo graeca*, distribution, Serbia and Montenegro, new country record

### SUBMITTED: April 19, 2004

AUTHORS: Ljiljana TOMOVIĆ, Institute of Zoology, Faculty of Biology, Studentski trg 16, 11000 Belgrade, Serbia and Montenegro < lili@bf.bio.bg. ac.yu >; Rastko AJTIĆ, Institute for Nature Conservation of Serbia, Dr Ivana Ribara 91, 11070 Belgrade, Serbia and Montenegro, Đorđe ĐOKOVIĆ, Saša ANTIĆ, Biology Students Association "Josif Pančić", Studentski trg 16, 11000 Belgrade, Serbia and Montenegro.

## Range extensions of Lycodon capucinus BOIE, 1827 in eastern Indonesia

The colubrid snake Lycodon capucinus BOIE, 1827 is widely distributed throughout Southeast Asia, southeastern China, the Philippines, and Indonesia (DE ROOIJ 1917; DE HAAS 1950; BOSCH 1985; ISKANDAR & COLIJN 2001). A recent colonization of Christmas Island, about 320 km south of Java, was reported by L. A. SMITH (1988). In eastern Indonesia, L. capucinus has been known from central, southwestern, and southeastern Sulawesi (DE ROOIJ 1917; ISKANDAR & TJAN 1996) and from the Lesser Sunda Islands of Sumbawa, Sumba, Savu, Roti, Timor, Flores, Lomblem, Alor, Lembata, and Wetar (DE ROOIJ 1917; How et al. 1996). Recent new island records for L. capucinus include Lombok, Moyo, Komodo, Adonara, Pantar, Kisar and Semau in the Lesser Sunda Islands, and Babar and Kai in Southeastern Maluku (How et al. 1996; MONK et al. 1997; ISKANDAR & COLIJN 2001 [politically, the islands of Wetar and Kisar also belong to Southeastern Maluku regency]). Here we report the presence of L. capucinus on the islands of Ambon, Seram and Buru (Central Maluku regency) from where we examined a total of six specimens.

Vouchers are deposited in the following collections: Forschungsinstitut und Naturmuseum Senckenberg, Frankfurt am Main, Germany (SMF); Texas Memorial Museum, The University of Texas at Austin, USA (TNHC). On Seram we collected two specimens on 14 April 1998 between 19:00 and 21:00 on the road between the villages Kairatu and Wae Samu on the western coast (SMF 82999), and 28 km NW (by road) of the Kairatu ferry port on the same road (TNHC 59520). On Buru, we collected a specimen (SMF 83000) on 18 April 1998 near the village of Labuan, Namrole, on the south coast. The Buru specimen regurgitated a partly digested, scincid lizard a few hours after capture. Three specimens of L. capucinus from the harbour town Waai in northeastern Ambon (SMF, uncatalogued) document its presence on that island as early as 1956.

The six specimens reported herein are the first records of *L. capucinus* and the genus *Lycodon* FITZINGER, 1826 for the islands of Ambon, Seram and Buru, and for the regency of Central Maluku and the northern Maluku biogeographical subregion (sensu HOW & KITCHENER 1997). In their external phenotype, these specimens closely resemble descriptions of this taxon as provided by DE ROOIJ (1917) and MERTENS (1930).

The Greater Sunda Islands on the Sunda shelf, Sulawesi and the Lesser Sunda Islands harbor a snake fauna originating from Asian genera and species, while the islands of southern and northern Maluku. Irian Jaya and adjacent islands are inhabited by genera and species that are more closely related to those in the Australo-Papuan region (How & KITCHENER 1997). Within the snakes of Indonesia, there is in addition evidence for considering the Lesser Sunda Islands and the southern and northern Maluku Islands as three unique biogeographic subregions with differing regions of origin for their assemblages, a relatively high degree of endemism and areas of incipient speciation (How & KITCHENER 1997). The currently known distribution of L. capucinus in Indonesia is clearly incongruent with these patterns: This species occurs in Sumatra, Java, Karimunjawa, Bawean and Bali, but had never been reported from Borneo until the discovery of a single specimen in a residential area (apparently in Sabah [STUEBING & INGER 1999]; ISKANDAR & COLIJN [2001] also list this species for Kalimantan). From Bali it ranges east across Wallace's line and well into the southern and northern Maluku biogeographical subregions (How et al. 1996; this paper). Throughout this biogeographically unlikely range, morphological differentiation in L. capucinus is remarkably low. Among four snake species from the Lesser Sunda Islands studied by How et al. (1996), L. capucinus showed the least morphological variation, with most populations overlapping considerably. These authors noted no variation in body colour or pattern in the specimens collected for their study, a fact also commented on by MERTENS (1930) and confirmed by the specimens from Ambon, Buru and Seram, which are very similar to L. capucinus from western Indonesia (e.g., urban Jakarta). The extensive distribution of L. capucinus indicates that it has good dispersal abilities. This is confirmed by its (re-)colonization of Moyo Island, which had been obliterated by ash during the eruption of the nearby Gunung Tambora on Sumbawa in 1815, and more recently, its colonization of Christmas Island (L. A. SMITH 1988), and is thought to account for the lack of morphological differentiation of populations through the Lesser Sunda region (How et al. 1996). Similarly, in the Philippines LEVITON (1965) demonstrated that *L. capucinus* is widely distributed but showed no apparent tendency to evolve morphologically distinct populations on different islands. He attributed these findings to a presumed recent arrival and dispersal in the Philippines, where *L. capucinus* was mainly associated with human habitations.

The specimens from Seram were collected from an asphalted road in an area used for timber production and related industries. The nearby port of Kairatu is served several times per day by ferries from Waai in Ambon, and has been used by international timber companies as well as by transmigrants arriving from western Indonesia. No specimens of L. capucinus had been collected during several biological surveys in Manusela National Park in central Seram in the late 1980s and early 1990s (EDGAR & LILLEY 1993; MONK et al. 1997). The collecting locality of the Buru specimen is near the main port of southern Buru (Labuan) which is several times per week served by supply ships from other Maluku islands, namely Ambon. In view of its recently discovered presence in Buru and Seram, the existence of L. capucinus in collections made in Ambon in the 1950s is not surpris-For centuries, Ambon served as the ing. principal regional basis for European colonialization, travel, commerce, and research throughout the Maluku and western New Guinean regions. Ambon's ports continue to connect the region, thus also bearing considerable potential for dispersing introduced fauna and flora in Maluku and beyond. We therefore propose the hypothesis, for testing in a phylogenetic context, that the presence of L. capucinus in Ambon is due to human activities, and that this species was introduced to Buru and Seram during the past few decades.

Lycodon capucinus attains a maximum length of 76 cm (MANTHEY & GROSSMAN 1997), lays 3-11 eggs in a clutch, possibly breeds twice a year, climbs well on rocks and trees, and lives on a diet of geckos, skinks, and small mammals (DUNN 1927; M. A.

SHORT NOTE	HERPETOZOA 17 (3/4)	Wien 30	Dezember 2004	SHORT NOTE	193
SHOKINOIL		, wien, 50	. Dezember 2004	SHOKINOIL	1/5

SMITH 1943). These snakes could therefore pose a threat to the autochthonous small vertebrate fauna of the islands where they have been introduced. Although it is presently unknown whether *L. capucinus* will colonize undisturbed habitats on these various islands, the potential consequences of the introduction of such exotic predators on island ecosystems should not be underestimated (FRITTS & RODDA 1998).

ACKNOWLEDGMENTS: We are grateful to Frank Bambang YUWONO (Jakarta), Bulquiny BAADIL-LA (formerly of Paso, Ambon) and Bapak ELI (Labuan, Buru) for help with logistics and assistance in the field.

REFERENCES: DE HAAS, C. P. J. (1950): Checklist of the snakes of the Indo-Australian archipelago (Reptiles - Ophidia).- Treubia, Bogor; 20: 511-625. DE ROOIJ, N. (1917): The reptiles of the Indo-Australian Archipelago; II. Ophidia. Leiden (E. J. Brill), 334 pp. DUNN, E. R. (1927): Results of the Douglas Burden expedition to the islands off Komodo. II. Snakes from the East Indies.- American Museum Novitates, New York City; 287: 1-7. EDGAR, P. W. & LILLEY, R. P. H. (1993): Herpetofauna survey of Manusela National Park; pp. 131-141. In: EDWARDS, I. D. & MACDONALD, A. A. & PROCTOR, J. (Eds.): The natural history of Seram; Andover, Hampshire, UK (Intercept Ltd.). FRITTS, T. H. & RODDA, G. H. (1998): The role of introduced species in the degradation of island ecosystems: a case history of Guam.- Annual Review of Ecology and Systematics, Palo Alto; 29: 113-140. How, R. A. & KITCHENER, D. J. (1997): Biogeography of Indonesian snakes.- Journal of Biogeography, Oxford; 24: 725-735. How, R. A. & SCHMITT, L. H. & SUYANTO, A. (1996): Geographical variation in the morphology of four snake species from the Banda Arcs, eastern Indonesia.- Biological Journal of the Linnean Society, London; 59: 439-456. IN DEN BOSCH, H. A. J. (1985): Snakes of Sulawesi: a checklist, keys and additional biogeographic remarks.-Zoologische Verhandelingen, Leiden; 217: 1-50. Iskandar, D. T. & Colijn, E. (2001): A Checklist of Southeast Asian and New Guinean Reptiles - Part I. Serpentes. Jakarta (Biodiversity Conservation Project, Indonesian Institute of Sciences, Japan International Cooperation Agency, The Ministry of Forestry, The Gibbon Foundation and Institute of Technology Bandung), 195 pp. ISKANDAR, D. T. & TJAN, K. N. (1996): The amphibians and reptiles of Sulawesi, with notes on the distribution and chromosomal number of frogs; pp. 39-46. In: KITCHENER, D. J. & SUYANTO, A. (Eds.): Proceedings of the first international conference on eastern Indonesian-Australian vertebrate fauna, Manado, Indonesia, November 22-26, 1994. Perth (Western Australian Museum). LEVITON, A. E. (1965): Contributions to a review of Philippine snakes, VIII. The snakes of the genus Lycodon H. BOIE.- Philippine Journal of Science, Manila; 94: 117-140. MANTHEY, U. & GROSSMANN, W. (1997): Amphibien & Reptilien Südostasiens. Münster (Natur und Tier-Verlag), 512 pp. MERTENS, R. (1930): Die Amphibien und Reptilien der Inseln Bali, Lombok, Sumbawa und Flores.- Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft, Frankfurt am Main; 42: 117-344. MONK, K. A. & DE FRETES, Y. & REKSODIHARJO-LILLEY, G.

(1997): The ecology of Nusa Tenggara and Maluku. Singapore (Periplus), 923 pp. SMITH, L. A. (1988): *Lycodon aulicus capucinus* a colubrid snake introduced to Christmas Island, Indian Ocean.- Records of the Western Australian Museum, Perth; 14: 251-252. SMITH, M. A. (1943): The Fauna of British India, Ceylon and Burma, including the whole of the Indo-Chinese Sub-Region; Reptilia and Amphibia Vol. III Serpentes. Reprint 1981. New Delhi (Today's & Tomorrow's Printers and Publishers), xii, 583 pp. STUEBING, R. B. & INGER, R. F. (1999): A field guide to the snakes of Borneo. Kota Kinabalu (Natural History Publications [Borneo]), 254 pp.

KEY WORDS: Squamata: Serpentes: Colubridae: Lycodon capucinus; geographical distribution; new island records; biogeography; invasive species; Ambon; Buru; Seram; Maluku; Indonesia

#### SUBMITTED: May 7, 2004

AUTHORS: Ulrich KUCH, Sektion Herpetologie, Forschungsinstitut und Naturmuseum Senckenberg, Senckenberganlage 25, D-60325 Frankfurt am Main, Germany < u.kuch@em.uni-frankfurt.de >; Jimmy A. McGUIRE, Museum of Vertebrate Zoology, 3101 Valley Life Science Building, University of California, Berkeley, CA 94720-3160, USA < mcguirej @socrates.berkeley.EDU >.

## Record of the Black-striped Snake, *Rhinoplocephalus nigrostriatus* (KREFFT, 1864), from Papua, Indonesia

The genus *Rhinoplocephalus* MÜLLER, 1885 contains six species of small to moderately sized elapid snakes found in the coastal and adjacent regions of Australia (COGGER 2002). Two of these, the Carpentaria Snake Rhinoplocephalus boschmai (BRONGERSMA & KNAAP-VAN MEEUWEN, 1964) and the Black-striped Snake Rhinoplocephalus nigrostriatus (KREFFT, 1864) also occur in southern New Guinea: The type locality of R. boschmai is Merauke in the southeast of the Province of Papua, Indonesia, and this species is suspected to also occur in adjacent parts of Papua New Guinea (O'SHEA 1996), in addition to its more extensive range in eastern Queensland, Australia (COGGER 2002). Rhinoplocephalus nigrostriatus, on the other hand, inhabits the coast and adjacent ranges of northeastern Australia from Cape York Peninsula to southern Queensland and has been collected in southern Papua New Guinea (Western Province: southern Trans-Fly, Oriomo River to Bensbach River: O'SHEA 1996), but has so far not been recorded from the Indonesian Province of Papua (DE ROOIJ

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Herpetozoa

Jahr/Year: 2004

Band/Volume: 17\_3\_4

Autor(en)/Author(s): Kuch Ulrich, McGuire Jimmy A.

Artikel/Article: <u>Range Extensions of Lycodon capucinus BOIE, 1827 in</u> <u>eastern Indonesia 191-193</u>