

bridge; 50 (2): 87-265. BÖHME, W. & WAGNER, P. & MALONZA, P. & LÖTTERS, S. & KÖHLER, J. (2005): A new species of the *Agama agama* group (Squamata: Agamidae) from western Kenya, East Africa, with comments on *Agama lionotus* BOULENGER, 1896.—Russian J. Herpetol., Moskva; 12 (2): 83-90. BROADLEY, D. G. & HOWELL, K. M. (1991): A check list of the reptiles of Tanzania with synoptic keys.—Syntarsus, Bulawayo; 1: 1-70. BURMANN, A. (2006): Phylogenie & Taxonomie der Agamen (*Agama lionotus*-Komplex) Ostafrikas: morphologische & genetische Untersuchungen.—Unpublished MSc. thesis, University of Bonn, pp. 1-244. DAUDIN, F. M. (1802): Histoire naturelle, générale et particulière des reptiles, vol. 3. Paris (F. Dufart), 452 pp.

KEY WORDS: Reptilia, Squamata, Sauria, Agamidae, *Agama lionotus* complex, *Agama lionotus usambarae*, synonymisation, taxonomy, Tanzania, Usambara Mountains

SUBMITTED: November 14, 2006

AUTHOR: Philipp WAGNER, Zoologisches Forschungsmuseum Alexander Koenig, Adenauerallee 160, D-53113 Bonn, Germany, < philipp.wagner.zfmk@uni-bonn.de >

First record of *Hemidactylus mindiae* BAHÄ EL DİN, 2005 from Jordan

Hemidactylus turcicus turcicus (LINNAEUS, 1758) was traditionally believed to be the only Jordanian taxon of the genus *Hemidactylus* OKEN, 1817. It is considered as a common reptile species distributed widely in all main ecozones of Jordan (e.g. DISI 1996; DISI & AMR 1998; DISI et al. 1999, 2001, 2004). However, the situation seems to be more complex. Using mitochondrial DNA sequences, CARRANZA & ARNOLD (2006) found 4.4 % genetic divergence between their specimens from Dair al Khaf (32°19'N, 36°53'E; northern Jordan) and Wadi al Burbeytah (30°59'N, 35°42'E; central part of western Jordan) and all other investigated samples of *H. t. turcicus* which covered an essential part of the range of this form. This finding indicates that the taxonomic status of the Jordanian form of *H. turcicus* is not clear.

CARRANZA & ARNOLD (2006) used the name *Hemidactylus turcicus lavadeserticus* MORAVEC & BÖHME, 1997 for the specimen of *Hemidactylus* originating from Dair al Khaf. However, determination of this individual was not based on an appropriate examination of the relevant material. *Hemidactylus t. lavadeserticus* was described

from southern Syria from the area of the black lava desert east of Jabal Duruz (ca. 30 km northwest of the Jordanian border) and its occurrence in the basalt desert of Jordan and Saudi Arabia has been expected (MORAVEC & BÖHME 1997). The three specimens of *Hemidactylus* collected at Dair al Khaf (margin of basalt area of northern Jordan), which are deposited in the collection of the National Museum in Prague (NMP6V 72130/1-3) and include also the specimen sequenced by CARRANZA & ARNOLD (2006) resemble *H. t. lavadeserticus* in colouration. A similar colour form can be found also among the three specimens (NMP6V 72131, 72740/1-2) obtained from the nearby locality of Jawa (32°20'N, 37°02'E). Nevertheless, all these individuals clearly differ from the type specimens of *lavadeserticus* (NMP6V 34831/1, 35540/1-4; ZFMK 64409) in the following morphological features (measurements in mm): larger body size (max. male SVL = 49.5 versus 48.0, max. female SVL = 50.5 versus 48.5); less depressed head (head depth: 44.4-54.8 % of head length measured from snout to posterior edge of ear opening, n = 6 versus 35.8-44.6 %, n = 6); larger and more prominent dorsal tubercles; lower number of lamellae under 1st toe (6-7, n = 6 versus 7-8, n = 6) and 4th toe (10-11, n = 6 versus 10-12, n = 6); higher number of preanal pores (7-8, n = 2 versus 6, n = 3); higher number of tail segments bearing 6 tubercles ((3)6-8, n = 5 versus 2-5, n = 6); and anterior postmentals being in contact with 1st lower labials (n = 6, there is a punctual contact with the 2nd lower labial on the left side in NMP6V 72740/1). Thus, the occurrence of *H. t. lavadeserticus* in Jordan remains to be proved.

Recently, BAHÄ EL DİN (2005) described a new gecko *Hemidactylus mindiae* BAHÄ EL DİN, 2005 from the mountain range of southern Sinai. A thorough examination of the material of *Hemidactylus* available from the area of southern Jordan revealed occurrence of this species (Fig. 1) also within the Jordanian borders. In all, five voucher specimens of *H. mindiae* from two localities are deposited in the collection of the National Museum Prague: NMP6V 71323/1-2, 1 female, 1 subadult, Jordan, Jabal Ghazali, 29°31'N, 35°25'E, 16-26 June 2000, Leg. D. MODRÝ; NMP6V 72739/



Fig. 1: *Hemidactylus mindiae* BAHA EL DIN, 2005 from Wadi Ramm, southern Jordan.

1–3, 1 female, 1 male, 1 subadult, Jordan, Nughra-Raddet Salem, Wadi Ramm, 29° 35'N, 35°26'E, 22 June 2005, Leg. A. BAKER and D. MODRÝ). This series has the following morphological characteristics (measurements in mm): max. male SVL = 50.0, max. female SVL = 56.5; head depth = 40.5–46.7 % of head length, $n = 5$; lamellae under 1st toe = 6–7, $n = 5$; lamellae under 4th toe = 10, $n = 5$; number of preanal pores = 4, $n = 1$; number of tail segments bearing 6 tubercles = 4–7, $n = 5$; and anterior post-



Fig. 2: Habitat of *Hemidactylus mindiae* BAHA EL DIN, 2005 in Wadi Ramm, southern Jordan.

mentals being in contact with 1st lower labials (1 specimen) and 2nd lower labials (4 specimens).

Our records extend the known range of *H. mindiae* from its type locality in the southern Sinai for ca. 200 km linear distance to the northeast and suggest a more extensive distribution of the species in the area of the Gulf of Aqabah and the southern part of Wadi Araba and northern Arabia. This should be substantiated by re-examining materials from these regions.

Wadi Ramm area consists of precipitous sandstone and granitic mountains, which are isolated from each other by flat corridors covered with mobile sand-dunes. The altitude varies from 800–1750 m. Wadi Ramm massifs are formed by several types of sandstone, which was deposited by braided rivers running from south to north during the lower Cambrian and lower Ordovician. Sedimentation continued during the Ordovician and Silurian periods. The area was subjected to the present day phase of lift (ABED 2002). Ramm area is a mosaic of diverse habitats, ranging from sand dunes, open sand, rocky slopes and cliffs, mud flats, open gravel, caves, etc. (ABU BAKER et al. 2004).

The specimens of *H. mindiae* were collected along the walls of canyons in Khazali Mountain and Al Nughra (Fig. 2). Both canyons are narrow and extend over 500–1500 m deep into the rocky slopes reaching a maximum width of 4–1 m, respectively, with relatively high humidity and cool temperature during day time. The altitude is about 1150 m a.s.l. The ground flora consists of a few solitary wild fig trees, *Ficus pseudo-sycomorus*, and bushes of Venus Maidenhair fern, *Adiantum capillus-veneris*. Within the same habitat, *Lacerta* cf. *kulzeri* was also collected. A total of 34 species of reptiles representing nine families were collected from different habitats in Wadi Ramm and its vicinity (ABU BAKER et al. 2004). In their paper these authors considered all the collected *Hemidactylus* from rocky areas as *H. turcicus*.

ACKNOWLEDGEMENTS: We would like to express our gratitude to the Aqaba Special Economic Zone Authority (ASEZA) for their support and encouragements. Also, we would like to thank Prof. Dr. Ratib AL-ORAN, president of Al Hussein Bin Talal University, Ma'an, and the Royal Society for

the Conservation of Nature for the logistic support. The work of the last author (JM) was supported by the Grant Agency of the Czech Republic, project no. 206/05/ 2334, and by the Ministry of Culture of the Czech Republic, projects nos. MK00002327201 and DE06P04OMG008.

REFERENCES: ABED, A. M. (2002): An overview of geology of Jordan; pp. 21-24. In: DISI, A. (Ed.): Jordan country study on biological diversity: The herpetofauna of Jordan; United Nations Environment Programme, Amman. ABU BAKER, M. & QARQAZ, M. & RIFAI, L. & HAMIDAN, N. & AL OMARI, K. & MODRÝ, D. & AMR, Z. (2004): Results of herpetofaunal inventory of Wadi Ramm Protected Area, with notes on some relict species.- Russian Journal of Herpetology, Moskva; 11 (1): 1-5. BAHA EL DIN, S. (2005): An overview of Egyptian species of *Hemidactylus* (Gekkonidae), with the description of a new species from the high mountains of South Sinai.- Zoology in the Middle East, Heidelberg; 34: 11-26. CARRANZA, S. & ARNOLD, E. N. (2006): Systematics, biogeography, and evolution of *Hemidactylus* geckos (Reptilia: Gekkonidae) elucidated using mitochondrial DNA sequences.- Molecular Phylogenetics and Evolution, San Diego; 38: 531-545. DISI, A. M. (1996): A contribution to the knowledge of the herpetofauna of Jordan. VI. The Jordanian herpetofauna as a zoogeographic indicator.- Herpetozoa, Wien; 9 (1/2): 71-81. DISI, A. M. & AMR, Z. S. (1998): Distribution and ecology of lizards in Jordan (Reptilia: Sauria); pp. 43-66. In: FRITZ, U. & OBST, F. J. & ANDREAS, B. (Eds.): Contribution to a "Herpetologia arabica", Faunistische Abhandlungen Staatliches Museum für Tierkunde Dresden, 21, Suppl., 182 pp. DISI, A. M. & AMR, Z. S. & MARTENS, H. (2004): On a collection of amphibians and reptiles made by J. KLAPPERICH in Jordan.- Herpetozoa, Wien; 16 (3/4): 141-150. DISI, A. M. & MODRÝ, D. & NEČAS, P. & RIFAI L. (2001): Amphibians and reptiles of the Hashemite Kingdom of Jordan. An atlas and field guide. Frankfurt a. M. (Chimaira), 408 pp. DISI, A. M. & MODRÝ, D. & BUNIAN, F. & AL-ORAN, R. M. & AMR, Z. S. (1999): Amphibians and reptiles of the Badia region of Jordan.- Herpetozoa, Wien; 12 (3/4): 135-146. MORAVEC, J. & BÖHME, W. (1997): A new subspecies of the Mediterranean gecko *Hemidactylus turcicus* from the Syrian lava desert (Squamata: Sauria: Gekkonidae).- Herpetozoa, Wien; 10 (3/4): 121-128.

KEY WORDS: Reptilia: Squamata: Sauria: Gekkonidae, *Hemidactylus mindiae*, *Hemidactylus turcicus*, new country record, Jordan

SUBMITTED: September 08, 2006

AUTHORS: Zuhair S. AMR, Department of Biology, Faculty of Sciences, Jordan University for Science and Technology, P.O. Box 3030, 22110 Irbid, Jordan < amrz@just.edu.jo >; David MODRÝ, Department of Parasitology, University of Veterinary and Pharmaceutical Sciences, 612 42 Brno, Czech Republic; Mohammad ABU BAKER, Dept. Biological Sciences (MC066), 845 West Taylor, St. University of Illinois at Chicago, Chicago, IL 60607 USA; Mayas QARQAS, The Royal Society for the Conservation of Nature, Amman, Jordan; Jamaal AL Zaidanyen, Ramm Protected Nature Reserve, Aqaba Special Economic Zone Authority; Jiří MORAVEC, Department of Zoology, National Museum, 115 79 Prague 1, Czech Republic < jiri.moravec@nm.cz >.

Cochranella punctulata RUIZ-CARRANZA & LYNCH, 1995 – new record from the department of Antioquia (Colombia), with comments on the conservation status

During a herpetological survey carried out in the Magdalena River valley in October 2005, the authors found an individual of the poorly-known glassfrog *Cochranella punctulata* RUIZ-CARRANZA & LYNCH, 1995; Fig. 1) at Hacienda Santa Bárbara (ca. 06° 32'N / 74°38'W, between 515-575 m above sea level; 1 in Fig. 2), vereda Las Brisas, Maceo municipality, Department of Antioquia. This location is ca 70 km northeast of the nearest known record of this species (vereda La Granja, Cocorná municipality, Department of Antioquia, 06°3'N / 75°11' W, 900-930 m a.s.l.; RUIZ-CARRANZA & LYNCH 1995; 2 in Fig. 2). The female found (Museo de Herpetología Universidad de Antioquia, MHUA 4071; coll. P. D. GUTIÉRREZ-C.) was observed at night (2042 h) on herbaceous vegetation attached to rock boulders near to a small waterfall inside a secondary forest.

This glassfrog species has been listed as vulnerable (IUCN category VU B1ab(iii)), based on its small range area, fragmented distribution and the decline of its forest habitat quality (GIL & LYNCH 2004). The first two arguments are valid considering that until now there are documented records only from three disjunctive localities as indicated in the original description (departments of Antioquia, Caldas and Tolima – RUIZ-CARRANZA & LYNCH 1995; 2-4 in Fig. 2). However, we consider that the previous arguments are only an artifact of lack of surveys, since our new record suggests that there may be other unknown northern populations along the eastern flank of the Cordillera Central, to which *C. punctulata* is apparently restricted (ACOSTA-GALVIS et al. 2006; FROST 2006). This new locality indicates that this species has a more extended distribution range. On the other hand, the adverse effect of declining habitat quality on the frog's distribution range reported by Gil & Lynch (2004) is well substantiated since in some localities of *C. punctulata* such as Samaná (Department of Caldas) and Cocorná

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2007

Band/Volume: [20_1_2](#)

Autor(en)/Author(s): Amr Zuhair S., Modry David, Abu Baker Mohammad, Qarqas Mayas, Al Zaidanyen Jamaal, Moravec Jiri

Artikel/Article: [First record of Hemidactylus mindiae Baha El Din, 2005 from Jordan 73-75](#)