

Redescription of *Rhacophorus tuberculatus* (Anderson, 1871) and the validity of *Rhacophorus verrucopus* Huang, 1983

R. S. Naveen^{1,2}, Shuo Liu^{3,4}, S. R. Chandramouli⁵, S. Babu¹, P. V. Karunakaran¹, Honnavalli N. Kumara¹

¹ *Sálim Ali Centre for Ornithology and Natural History, Anaikatty, Coimbatore, Tamil Nadu, India*

² *EDGE of Existence Programme, Conservation and Policy, Zoological Society of London, London, NW1 4RY, UK*

³ *Kunming Natural History Museum of Zoology, Kunming Institute of Zoology, Chinese Academy of Sciences, Yunnan 650223, Kunming, China*

⁴ *Kunming Institute of Zoology, Chinese Academy of Sciences, Yunnan 650201, Kunming, China*

⁵ *Department of Ecology and Environmental Sciences, School of Life Sciences, Pondicherry University, Puducherry, India*

<https://zoobank.org/15047501-D9F4-4F96-B55A-E6778EBAB3F0>

Corresponding authors: S. Babu (sanbabs@gmail.com); S. R. Chandramouli (findthesnakeman@gmail.com)

Academic editor: Günter Gollmann ♦ Received 4 October 2023 ♦ Accepted 5 December 2023 ♦ Published 22 December 2023

Abstract

Rhacophorus tuberculatus and *Rhacophorus verrucopus* are two morphologically similar species described in 1871 and 1983 respectively. Their taxonomic distinctiveness has been questioned in the past. In the current study, we encountered frogs that we confer to *R. tuberculatus* based on morphological similarity to a syntype of this species. We redescribe the species based on a re-examination of a syntype, which is designated as a lectotype here, and additional specimens from Garo hills of Meghalaya. We also present molecular data, natural history notes, and report a range extension of this species. Molecular phylogenetic analysis based on the 16S rRNA fragment revealed minimal genetic divergences (0.20–1.74% uncorrected p-distance) between specimens identified as either *R. tuberculatus* or *R. verrucopus* from different locations. On the basis of molecular data and morphological characteristics, we conclude that *R. verrucopus* is a junior synonym of *R. tuberculatus*.

Key Words

Darwinian shortfall, Indo-Burma hotspot, range extension, Rhacophoridae, synonymy, systematics

Introduction

Frogs of the genus *Rhacophorus* Kuhl & Van Hasselt, 1822 occur across South and Southeast Asia and are represented by 44 extant species (Frost 2023). Despite many species of *Rhacophorus* being widespread, most of them are poorly studied, and information on their distribution is sparse (Ohler and Delorme 2005). *Rhacophorus tuberculatus* (Anderson, 1871) is one such poorly known species. It was described in 1871 from “Sebsaugor, Assam” (now Sivasagar 26.98515°N, 94.63878°E) and was subsequently reported from Arunachal Pradesh, Meghalaya, and West Bengal, India (Annandale 1912; Chanda

2002; Sen 2004; Das and Dutta 2007; Ahmed et al. 2009; Mathew and Sen 2010; Roy et al. 2018). Huang (1983) described *Rhacophorus verrucopus* from “Beibeng, Medo Xian” (Beibeng Township, Motuo County, Xizang Autonomous Region, China 29.23942°N, 95.17644°E), which closely resembles *R. tuberculatus* and this species was known only from its type locality and northern Myanmar (Fei 1999; Fei et al. 2009, 2010; Li et al. 2010; Fei et al. 2012; Liu et al. 2020; Fei 2020; Zug 2022). By studying the type and near-topotypical specimens of *R. verrucopus* and comparing them with the descriptions of *R. tuberculatus*, Che et al. (2020) considered that *R. verrucopus* and *R. tuberculatus* could be the same species.

However, since Che et al. (2020) did not obtain any molecular data or examined the types of *R. tuberculatus* from India and only made comparisons of external morphology, Frost (2023) does not currently adopt their view and still treats *R. verrucopus* as a valid separate species.

We encountered a *Rhacophorus* species, the identity of which was difficult to determine, during our recent surveys between 2021 and 2022 in the West Garo hills of Meghalaya, India. Herein, we ascertain the identity of that *Rhacophorus* species as *R. tuberculatus* based on comparison with the syntypes of *R. tuberculatus* and provide a redescription of the species after more than 150 years since its original description, provide photographs of live individuals, morphological measurements and genetic divergence of *R. tuberculatus* with other congeners using 16S rRNA gene and discuss its relationship with *R. verrucopus*.

Materials and methods

Study area

We conducted surveys in Sasatgre village (25.5250°N, 90.3350°E, ca. 940 m) and Baladingre village (25.514213°N, 90.398204°E, ca. 835 m) of West Garo Hills district of Meghalaya between 2020 and 2022.

Voucher collection

Frogs were caught by hand, photographed first and euthanized using 20% Benzocaine following Torreilles et al. (2009). A small portion of the liver tissue was extracted by making a narrow slit on the ventral aspect of the specimens and stored in Molecular Biology Grade Ethanol (BP2818). Specimens were later fixed in 90% Ethanol for two hours and then transferred to 70% Ethanol for long-term storage as museum specimens. In total, seven individuals were collected for this study: six adult males collected from near Sasatgre village (25.5250°N, 90.3350°E, 940 m asl.) and an adult female from near Baladingre village (25.514213°N, 90.398204°E, 835 m asl.). The specimens used for morphological and molecular analyses were deposited at the herpetological collection facility at Sálím Ali Centre for Ornithology and Natural history (SACON).

DNA extraction and molecular analysis

Total genomic DNA was extracted from tissue samples of two specimens of *R. tuberculatus* (SACON VA 148 and VA 800) with a DNA extraction and purification kit, following the manufacturer's protocols. 16S rRNA gene was amplified using the primers 16sAR-L (5'-CGCCT-GTTTATCAAAAACAT-3') and 16sBR-H (5'-CCG-GTCTGAACTCAGATCACGT 3') respectively (Kocher et al. 1989). Amplifications were performed in an Applied Bio Systems Veriti 96 well thermal cycler: 20 µl reactions with 4 µl of 5× Phusion HF buffer, 0.4 µl of 10mM dNTP,

0.2 µl of Phusion DNA Polymerase, 0.1 µl each of forward and reverse primers, 2.0 µl of DNA template and 13.2 µl of nuclease free water with the following procedure: initial denaturation of DNA at 95 °C for 5 min, 35 cycles of: denaturation at 95 °C for 1 m, annealing at 55 °C for 1 min, extension at 72 °C for 1 m and at last, final extension at 72 °C for 10 min. The amplicon was checked by running it through an agarose gel electrophoresis for a clear band of the desired region in the amplified PCR product. The amplified PCR product was purified and sequenced commercially (National Centre for Biological Sciences, Bengaluru). Sequences were edited and manually adjusted using SeqMan in Lasergene 7.1 (DNASTAR Inc., Madison, WI, USA) and MEGA 11 (Tamura et al. 2021). Species of the genus *Zhangixalus* were selected as outgroups following Liu et al. (2022). Homologous and outgroup sequences were obtained from GenBank (Table 1). The technical computation methods for sequence alignment, genetic distance calculation, the best substitution model selection, Bayesian Inference (BI) and Maximum Likelihood (ML) phylogenetic analyses were the same as those in Liu et al. (2021).

Morphometric measurements

The following measurements were recorded to the nearest 0.02 mm from the specimens using an INSIZE dial caliper: snout–vent length (SVL, from the tip of the snout to the anterior margin of the cloaca), axilla–groin distance (AG, from the posterior margin of the forelimb at its insertion point on the body to the anterior margin of the hind limb at its insertion point on the body), head length (HL, from the posterior edge of the mandible to the tip of the snout), head width (HW, the maximum width of the head at the angle of the jaws), head depth (HD, the maximum depth of the head), body width (BW, the maximum width of the body at the trunk), eye diameter (ED, the greatest horizontal diameter of the orbit), eye–nostril distance (EN, from the anterior border of the orbit to the middle of the nostril), eye–snout distance (ES, from the anterior border of the orbit to the tip of the snout), upper eyelid width (UEW, the maximum width of the upper eyelid), interorbital distance (IO, distance between the margins upper eyelids), internarial distance (IN, distance between the nostrils), upper arm length (UAL, from the axilla to elbow), lower arm length (LAL, from the posterior margin of the elbow to the base of the outer metacarpal tubercle), palm length (PAL, from the posterior border of the outer metacarpal tubercle to tip of the 3rd finger), femur length (FEL, from the cloaca to the knee), tibia length (TBL, from knee to heel), foot length (FOL, from inner metatarsal tubercle to the top of the 4th toe). Webbing formulae follows Savage and Heyer (1997).

Geographic range estimation

Geographic range of the target species was calculated by plotting the known occurrences of the species on a

Table 1. List of specimens and GenBank accession numbers for all 16S rRNA sequences included in this study.

Taxon	Voucher No.	Locality	GenBank No.
<i>Rhacophorus annamensis</i>	VNMN 4090	Dak Nong, Nam Nung, Vietnam	LC010566
<i>Rhacophorus baluensis</i>	FM235958	Sabah, Malaysia	KC961089
<i>Rhacophorus bengkulensis</i>	UTA A-62770	Lampung, Sumatra, Indonesia	KM212948
<i>Rhacophorus bipunctatus</i>	PUCZM/IX/SL360	Mizoram, India	MH087073
<i>Rhacophorus borneensis</i>	BORN:22410	Maliau Basin, Sabah, Malaysia	AB781693
<i>Rhacophorus calcaneus</i>	VNMN 4093	Dak Lac, Chu Yang Sin, Vietnam	LC010573
<i>Rhacophorus catamitus</i>	ENS 14726	Sumatra, Indonesia	KX398877
<i>Rhacophorus exechopygus</i>	VNMN 4107	Gia Lai, Kon Ka Kinh, Vietnam	LC010585
<i>Rhacophorus helenae</i>	AMS R 173230	Binh Thuan, Vietnam	JQ288087
<i>Rhacophorus hoabinhensis</i>	VNMN A.2016.16	Hoa Binh, Vietnam	LC331097
<i>Rhacophorus indonesiensis</i>	MZB: Amp:23619	Indonesia	AB983367
<i>Rhacophorus kio</i>	VNMN 4110	Gia Lai, Kon Ka Kinh, Vietnam	LC010589
<i>Rhacophorus lateralis</i>	SDB.2010.330	Karnataka, Bygoor, India	KC571277
<i>Rhacophorus malabaricus</i>	Rmal-In	Madikeri, India	AB530549
<i>Rhacophorus margaritifer</i>	ENS 16162	Java, Indonesia	KX398889
<i>Rhacophorus modestus</i>	ENS 16853	Sumatra, Indonesia	KX398904
<i>Rhacophorus napoensis</i>	GXNU YU000171	Napo, Guangxi, China	ON217796
<i>Rhacophorus nigropalmatus</i>	Rao081203	Malaysia	JX219438
<i>Rhacophorus norhayatie</i>	NNRn	Endau Rompin, Johor, Malaysia	AB728191
<i>Rhacophorus orlovi</i>	VNMN 3067	Huong Son, Ha Tinh, Vietnam	LC010598
<i>Rhacophorus pardalis</i>	FMNH273243	Sarawak, Bintulu, Malaysia	JX219454
<i>Rhacophorus poecilonotus</i>	ENS 16480	Sumatra, Indonesia	KX398920
<i>Rhacophorus pseudomalabaricus</i>	SDB.2011.1010	Kerala, Kadalar, India	KC593855
<i>Rhacophorus reinwardtii</i>	Rao081205	Malaysia	JX219443
<i>Rhacophorus rhodopus</i>	SCUM 060692L	Mengyang, Yunnan, China	EU215531
<i>Rhacophorus robertingeri</i>	VNMN 4123	Gia Lai, Kon Ka Kinh, Vietnam	LC010613
<i>Rhacophorus spelaeus</i>	IEBR A.2011.1	Khammouan, Lao	LC331095
<i>Rhacophorus translineatus</i>	Rao6237	Motuo, Xizang, China	JX219449
<i>Rhacophorus tuberculatus</i>	KIZ014154	Motuo, Xizang, China	MW111522
<i>Rhacophorus “verrucopus”</i>	Rao6254	Motuo, Xizang, China	JX219436
	SEABRI2019120056	Htamanthi, Sagaing, Myanmar	MW275978
<i>Rhacophorus tuberculatus</i>	SACON VA – 148	Meghalaya, India	OR836578
	SACON VA – 800	Meghalaya, India	OR836579
<i>Rhacophorus vampyrus</i>	VNMN 4125	Hon Ba, Khanh Hoa, Vietnam	LC010616
<i>Zhangixalus dennysi</i>	SCUM 060401L	Shaoguan, Guangdong, China	EU215545
<i>Zhangixalus dugritei</i>	SCUM 051001L	Baoxing, Sichuan, China	EU215541

distribution map generated using ARCGIS 10.5. The area within the minimum convex hull was computed by connecting the outermost occurrence points to calculate the extent of occurrence as defined by the IUCN (2001).

Results

The suggested best substitution model for BI was GTR+F+I+G4 and for ML was TIM2+F+I+G4, both analyses showed an essentially consistent topology (Fig. 1). The sequences of the newly collected specimens from Meghalaya, India clustered with the sequences of *R. verrucopus* from Myanmar with strong supports by both BI and ML (0.97/99), and they together clustered with the sequences of *R. verrucopus* from China with strong supports by both BI and ML (1/100). The genetic divergence (uncorrected p-distance) between the sequences of the newly collected specimens and the sequences of *R. verrucopus* from China ranged from 1.50% to 1.74%, the genetic divergence (uncorrected p-distance) between the sequences of the newly collected specimens and the sequence of

R. verrucopus from Myanmar ranged from 0.20% to 0.44% (Table 2).

Morphologically, the newly collected specimens (*R. tuberculatus*) from Meghalaya, India agree with the Syntype (ZSI 10154) and subsequent descriptions and figures by Annandale (1912) and Mathew and Sen (2010) of *Rhacophorus tuberculatus* in most aspects, especially in having a distinct tympanum, almost half as large as the eye; absence of vomerine teeth; presence of partial, sheath-like webbing on fingers and fully developed webbing on toes; pointed projection at tibio-tarsal articulation; well-developed and expanded discs on toe. Hence, we consider these newly collect-

Table 2. Genetic divergences (uncorrected p-distance in %) between specimens identified as either *Rhacophorus tuberculatus* or *R. verrucopus* from different locations.

Species	Voucher	1	2	3	4
<i>Rhacophorus tuberculatus</i>	SACON VA – 148 India				
	SACON VA – 800 India	0.22			
<i>Rhacophorus “verrucopus”</i>	SEABRI2019120056 Myanmar	0.20	0.44		
	Rao6254 China	1.50	1.74	1.24	
	KIZ014154 China	1.54	1.74	1.28	0.00




Figure 1. Bayesian inference tree of the genus *Rhacophorus* based on partial 16S rRNA fragments. Numbers before slashes indicate Bayesian posterior probabilities (≥ 0.90 remain) and numbers after slashes indicate ultrafast bootstrap support for ML analyses (≥ 90 remain).

ed specimens to belong to *R. tuberculatus*. In addition, there were also no obvious morphological differences between the newly collected specimens of *R. tuberculatus* and *R. verrucopus* from China and Myanmar (see Table 3).

Integrating the results of morphological data and also considering the shallow genetic divergence that is usually considered as intraspecific variation in the genus *Rhacophorus*, we second Che et al. (2020) in stating that *R. tuberculatus* and *R. verrucopus* are conspecific, and formally place *R. verrucopus* under the subjective synonymy of *R. tuberculatus*. Below, we give a formal synonymy list and provide the description of the specimen ZSI 10154 and designate it as the lectotype of *Rhacophorus tuberculatus*. Additionally, we also provide measurements of the newly collected material for comparison.

Systematics

Rhacophorus tuberculatus (Anderson, 1871)

Polypedates tuberculatus Anderson, 1871.

Rhacophorus tuberculata – Boulenger, 1882.

Rhacophorus (*Rhacophorus*) *tuberculatus* – Ahl, 1931.

Rhacophorus schlegelii tuberculatus – Wolf, 1936.

Rhacophorus verrucopus Huang, 1983, syn. nov.

Rhacophorus tuberculatus – Inger, 1985.

Rhacophorus (*Rhacophorus*) *verrucopus* – Dubois, 1987 «1986».

Redescription of *Rhacophorus tuberculatus* (Anderson, 1871). Specimens examined: ZSI 10154, lectotype by present designation, adult female, collected by Anderson from “Sebsaugor, Assam”; SACON VA – 143, 144, 145, 146, 147 and 148, adult males collected

Table 3. Morphometric measurements (in mm) of *R. tuberculatus* and *R. turpes* from the current study (*Designated as lectotype).

Species	<i>Rhacophorus tuberculatus</i> (Voucher no., sex and values)								<i>Rhacophorus turpes</i> (Voucher No., sex and values)	
Voucher no: Morphometric variables	*ZSI 10154	SACON VA 143	SACON VA 144	SACON VA 145	SACON VA 146	SACON VA 147	SACON VA 148	SACON VA 800	BMNH 1940.6.1.30	BMNH 1974.828-832
	Female	Male	Male	Male	Male	Male	Male	Female	Female	Female
SVL (mm)	39.10	30.32	29.38	30.88	28.16	27.02	30.76	34.02	37.42	35.80
AG	17.44	12.80	12.20	14.06	12.84	12.48	13.74	16.82	21.28	18.02
BW	16.92	7.52	5.82	7.28	6.76	6.10	8.44	7.44	14.22	11.46
HL	11.40	8.74	9.18	10.00	8.40	7.92	10.02	11.62	12.22	10.76
HW	11.10	9.38	9.00	9.34	8.92	8.86	9.40	12.00	9.74	9.78
HD	6.04	4.32	4.08	4.60	3.92	3.62	3.54	6.32	4.68	4.28
ED	4.42	4.28	4.10	3.70	3.70	3.84	4.18	4.90	3.54	2.98
EN	2.80	2.98	2.00	2.32	2.84	2.84	2.56	3.08	2.74	3.30
ES	4.42	3.78	3.64	4.32	4.52	4.22	4.32	5.62	4.90	5.43
TYD	1.82	1.68	1.80	2.08	1.20	1.72	1.70	2.38	2.44	2.76
ET	1.24	0.48	0.80	0.52	0.60	0.40	0.72	0.66	1.12	1.18
UEW	2.86	3.70	4.20	3.70	3.66	3.68	3.70	5.54	2.44	2.02
IO	4.64	3.12	3.70	2.82	2.98	2.72	3.02	4.14	4.28	4.12
IN	3.50	2.40	2.36	1.74	2.50	2.80	2.98	3.22	2.80	3.08
UAL	6.46	5.72	5.62	4.72	4.92	4.04	4.50	5.42	5.24	6.98
LAL	7.62	5.64	5.42	5.66	5.24	5.94	5.34	8.00	7.00	7.68
PAL	9.42	6.84	7.32	7.20	6.90	6.70	7.40	9.58	9.18	8.64
FEL	16.92	13.56	13.62	11.72	11.74	11.72	13.96	15.06	14.88	16.22
TBL	18.80	15.48	14.82	13.8	14.4	13.74	14.32	17.28	16.90	18.28
TAL	12.44	8.48	9.34	7.34	8.44	7.20	9.14	9.90	8.94	9.32
FOL	19.76	11.12	10.88	9.38	9.84	9.28	11.88	12.96	10.82	10.36

from near Sasatgre (25.5250°N, 90.3350°E, ca. 940 m asl.) by RSN between 13th–26th May 2020 and SACON VA – 800 Adult female collected from near Baladingre (25.514213°N, 90.398204°E, ca. 835 m asl.) by RSN on 26th February 2022.

Diagnosis. *Rhacophorus tuberculatus* can be differentiated from all known congeners by the following suite of external morphological characters: small to medium adult size (mean SVL 29.4 mm; range 27.0–30.9 mm); distinct tympanum, almost half as large as the eye; absence of vomerine teeth; presence of a prominent calcar at tibio-tarsal articulation; presence of partial, sheath-like webbing on fingers and fully developed webbing on toes; well-developed and expanded discs on toes; and a dorsal colouration of uniform pale brown with mild traces of an irregular patch on the head and mossy greenish patches near the shoulders in some individuals.

Description of Lectotype (Fig. 2). Head flat, almost as long as wide (HL:HW 1.03); snout slightly pointed in dorsal view, rounded in lateral aspect, projecting slightly beyond margin of the lower jaw; canthus rostralis distinct, bluntly angular; nostrils much closer to tip of snout than eye; eyes large (ED:HL 0.39); tympanum distinct almost half as large as eye (TYD:ED 0.41); Supra-tympanic fold distinct, originating from the posterior of eye to the axilla; Upper eyelids wide, (UEW 2.86), narrower than the interorbital space (UEW:IO 0.62). Inter-orbital space broader than the inter-narial space (IO:IN 1.33). Upper arms short (UAL:SVL 0.17), shorter than the lower arms (UAL:LAL 0.85); palm length longer than the upper arms (UAL:PAL 0.69); pointed projection (calcar) at tibio-tarsal articulation; relative length of fingers I < II < IV < III, tips of all fingers with well-developed discs

with distinct circum-marginal grooves. Fingers partially webbed. Relative length of toes I < II < III < V < IV; tips of toes with well-developed disks with distinct circum-marginal grooves; disks smaller than those of toes. Dorsal skin smooth; flanks wrinkled; underside of chin and chest smooth, abdomen and thigh coarsely granular; the granulation much denser around the cloacal region; outer margin of both limbs with low dermal ridges.

Colouration in preservative. Dorsal colouration uniformly pale brown with mild traces of an irregular bluish black patch on the head. Ventral aspect of body pale cream coloured (Fig. 2).

Colouration in life of frogs recorded from Meghalaya. Dorsum, overall pale to medium brown in colour with small scattered black dots and three or four dark blackish brown transverse bands across the thigh and tibial region, tarsus feet and webbing between toes orangeish-red in colour. Webbing in fingers translucently yellow. Ventral sides mild brownish white and groins, thighs and the rest of the legs brownish yellow to dark red in colour. Some individuals with irregular florescent green patches on the head and mid body region (Fig. 4).

Natural history. The specimens of *R. tuberculatus* from Meghalaya examined during this study were collected from two different locations within West Garo Hills. A small shallow stream running parallel to the eastern boundary of the Sasatgre community reserve, the stream was bounded on both side by cardamom and banana plantations. The frogs were encountered at 1900 – 2300 hrs in the month of May, found perched on leaves of yam and cardamom plants, one to two meters above ground level. The other location was a similar habitat from another cardamom plantation near a forest patch near Baladingre village.




Figure 2. The ventral and dorsal view of the lectotype of *Rhacophorus tuberculatus* ZSI 10154.



Figure 3. The ventral and dorsal view of a syntype of *Rhacophorus turpes* BMNH 1940.6.1.30.



Figure 4. *Rhacophorus tuberculatus* in life from Garo hills, Meghalaya.

Table 4. Morphological (in mm) comparisons between our newly collected specimens of *Rhacophorus tuberculatus* from India and *R. verrucopus* from China and Myanmar. Data for *R. verrucopus* from China were obtained from Huang (1983) and Che et al. (2020), and data for *R. verrucopus* from Myanmar were from Liu et al. (2020) (“–” data unavailable).

Morphological variables	<i>R. tuberculatus</i>		<i>R. verrucopus</i>		<i>R. verrucopus</i>
	India		China		Myanmar
	Mean (Range) Male, n=6	Female, n=1	Mean (Range) Male, n=7	Female, n=1	Female, n=1
SVL	29.4 (27.0–30.9)	34.00	37.9 (36.0–40.6)	41.60	52.00
HL	9.0 (7.9–10.0)	11.60	12.5 (12.0–13.0)	12.20	17.60
HW	9.2 (8.9–9.4)	12.00	11.4 (10.9–12.3)	11.60	15.70
ED	4.0 (3.7–4.3)	4.90	4.3 (3.9–4.9)	5.20	5.40
ES	4.1 (3.6–4.5)	5.60	5.7 (5.1–6.7)	5.90	7.70
TYD	1.7 (1.2–2.1)	2.40	2.2 (2.0–2.4)	2.70	3.10
UEW	3.8 (3.7–4.2)	5.50	2.9 (2.5–3.6)	3.40	–
IO	3.1 (2.7–3.7)	4.10	4.3 (3.8–5.0)	3.60	–
IN	2.5 (1.7–3.0)	3.20	3.5 (3.2–4.1)	3.10	4.30
LAL+ PAL	12.6 (12.1–12.9)	17.60	16.5 (16.0–17.0)	18.80	17.80
PAL	7.1 (6.7–7.4)	9.60	10.6 (10.1–11.1)	11.70	–
FEL	12.7 (11.7–14.0)	15.10	17.3 (16.3–17.9)	20.50	22.90
TBL	14.4 (13.7–15.5)	17.30	17.9 (17.3–18.3)	21.20	24.20
FOL	10.4 (9.3–11.9)	13.00	15.3 (14.6–16.2)	19.00	20.40
HL/SVL	0.31 (0.29–0.33)	0.34	0.33	0.29	0.34
HL/HW	0.99 (0.89–1.07)	0.97	1.10	1.05	1.12
ED/HL	0.44 (0.37–0.49)	0.42	0.35	0.43	0.31
TYD/HL	0.19 (0.14–0.22)	0.21	0.17	0.22	0.18
FEL/SVL	0.43 (0.38–0.46)	0.44	0.46	0.49	0.44
TBL/SVL	0.49 (0.45–0.51)	0.51	0.47	0.51	0.47
TBL/FEL	1.14 (1.03–1.23)	1.15	1.04	1.03	1.06

Discussion

“Darwinian shortfall” is a major challenge faced by conservationists today, the lack of availability of molecular data for several extant species is a common phenomenon across taxa, leading to a situation where phylogenetic information is absent for most organisms, thus inhibiting a robust understanding of phylogenetic relationships within a particular group (Diniz-Filho et al. 2013). *Rhacophorus tuberculatus* is one such species which was described by Anderson (1871), based on three adult specimens. The original description was brief and lacked photographs or diagrams since it was from more than 150 years ago. Although several subsequent studies such as Ahmed et al. (2009) and Roy et al. (2018) reported photographic records of this species, there had been no studies involving a detailed taxonomic assessment of the species to date. As a result, the species had been overlooked in several studies, including the description of *Rhacophorus verrucopus* by Huang (1983). The validity of this species has been doubted by Che et al. (2020) who suggested that *R. verrucopus* could be a junior synonym of *R. tuberculatus* but due to the lack of molecular evidence this was not accepted. In the current study, we provide genetic divergence between *R. tuberculatus* and other congeners, including ‘*R. verrucopus*’ using 16s rRNA gene, thus filling an important knowledge gap. Further examination of *R. verrucopus* from China and Myanmar and *R. tuberculatus* from India based on re-examination of types and also freshly collected specimens used in this study revealed that there is an extensive overlap in morphology, and shallow molecular divergence (of a level that usually qualifies to be considered an intraspecific variation in the genus) between *R. verrucopus* and *R. tuberculatus*. Based on these

lines of evidence we endorse the conclusions of Che et al. (2020) and formally place *R. verrucopus* under the junior synonymy of *R. tuberculatus*. Examination of the types (BMNH 1940.6.1.30 and BMNH 1974.828-832) of *Rhacophorus turpes* described from Kachin region of Northern Myanmar (26.24972°N, 97.23878°E) by Smith (1940) revealed that these specimens also closely match the morphological characters of *R. tuberculatus* except for a slightly truncated snout, a relatively less prominent tibio-tarsal projection, and fewer granulations on the ventral surface (Fig. 3). However, further studies utilizing molecular approaches based on fresh material are necessary to assess the taxonomic status of this species.

The current study also addresses the re-assessment of the threat status of *R. tuberculatus* as per IUCN Red List criteria of this Data Deficient species. *Rhacophorus tuberculatus* was known with certainty only from its type locality until now. However, in this study, we were able to resolve the taxonomic confusion with this species thus mapping its actual distribution range for the first time. The new records of this species from West Garo hills of Meghalaya mark the westernmost limit of the distribution of the species (Fig. 5). The up-listing or down-listing of species from one threat category to another of the IUCN Red List requires an assessment against all the five criteria (A–E, with 11 sub-criteria) but only one criterion needs to be fulfilled for designation of threat categories (IUCN 2001). In the case of *R. tuberculatus*, the information on geographic distribution seems the most accurate and reliable among all other criteria and hence, used for a conservative estimate of the extent of occurrence (Criteria B, B1). The current Extent Of Occurrence for this species estimated based on all known localities is about 1,07,600 km² and hence,




Figure 5. Updated distribution map of *R. tuberculatus* including records from Meghalaya (Current study) in green and black circle, from Myanmar in green and black square, type locality of “*R. verrucopus*” in red circle and type locality of *R. tuberculatus* in green circle.

with all the updated information presented here, we recommend transferring the species from Data Deficient to Least Concern.

Acknowledgements

First and foremost, we are grateful to Mr. Lising G. Momin and the villagers of Sasatgre village for their invaluable assistance during the fieldwork. Our sincere thanks are due to the Additional Principal Chief Conservator of Forests (Wildlife) and Chief Wildlife Warden (CWLW), Meghalaya, and officers of the Department of Forest and Environment, Government of Meghalaya, for facilitating permission from the Community Reserve Management Committees to carry out the field study (No. FWC/Research/15/603-04 dated 31st May 2019). We extend our thanks to the Technology Platform Services, C-CAMP, National Centre for Biological Sciences, for their expertise in sequencing samples used in this study. Additionally, our gratitude goes to Dr. Kaushik Deuti for helping with the examination of specimens from the Zoological Survey of India, and to Dr. David Gower, Dr. Jeff Streicher, Dr. Simon Loader and Dr. Rikki Gumbs for facilitating the examination of specimens from the British Natural History Museum.

References

- Ahmed F, Das A, Dutta SK (2009) Amphibians and Reptiles of Northeast India: A Photographic Guide. Aaranyak Guwahati, India, 169 pp.
- Anderson J (1871) A list of the reptilian accession to the Indian Museum, Calcutta from 1865 to 1870, with a description of some new species. *Journal of the Asiatic Society of Bengal* 40: 12–39.
- Annandale N (1912) Zoological results of the Abor Expedition, 1911–1912. I. BatrachiaAmphibia. *Records of the Indian Museum* 8: 7–36. <https://doi.org/10.5962/bhl.part.1186>
- Chanda SK (2002) Handbook. Indian Amphibians. Zoological Survey of India, Kolkata, 335 pp.
- Che J, Jiang K, Yan F, Zhang Y (2020) Amphibians and Reptiles in Tibet–Diversity and Evolution. Science Press, Beijing, 803 pp. [In Chinese with English abstracts and species descriptions]
- Diniz-Filho JA, Loyola RD, Raia P, Mooers AO, Bini LM (2013) Darwinian shortfalls in biodiversity conservation. *Trends in Ecology & Evolution* 28(12): 689–695. <https://doi.org/10.1016/j.tree.2013.09.003>
- Das I, Dutta SK (2007) Sources of larval identities for amphibians of India. *Hamadryad* 31: 152–181.
- Fei L (1999) Atlas of Amphibians of China. Henan Press of Science and Technology, Zhengzhou, 432 pp.
- Fei L (2020) Atlas of Amphibians in China (Field Edn). Henan Science and Technology Press, Zhengzhou, 837 pp.

- Fei L, Hu SQ, Ye CY, Huang YZ (2009) Fauna Sinica (Vol. 2). Amphibia Anura. Science Press, Beijing, 957 pp.
- Fei L, Ye CY, Jiang JP (2010) Colored Atlas of Chinese Amphibians. Sichuan Publishing House of Science and Technology, Chengdu, 519 pp.
- Fei L, Ye CY, Jiang JP (2012) Colored Atlas of Chinese Amphibians and Their Distributions. Sichuan Publishing House of Science and Technology, Chengdu, 620 pp.
- Frost DR (2023) Amphibian Species of the World: An Online Reference. Version 6.1 (Date of access 23 Nov 2023). Electronic Database accessible at American Museum of Natural History New York, USA. <https://doi.org/10.5531/db.vz.0001>
- Huang YZ (1983) A new species of flying frog from Xizang – *Rhacophorus verrucopus*. Acta Herpetologica Sinica 2(4): 63–65.
- Kocher TD, Thomas WK, Meyer A, Edwards SV, Pääbo S, Villablanca FX, Wilson AC (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. Proceedings of the National Academy of Sciences of the United States of America 86: 6192–6200. <https://doi.org/10.1073/pnas.86.16.6196>
- Kumar R, Jayanta B, Ramie H, Ahmed MF (2018) Amphibians of the Dibang River Basin, Arunachal Pradesh: An annotated checklist with distribution records. Journal of Threatened Taxa 10: 12940–12952. <https://doi.org/10.11609/jott.4249.10.15.12940-12952>
- Liu S, Hou M, Mo MZ, Rao DQ (2021) A new species of *Micryletta* Dubois, 1987 (Anura, Microhylidae) from Yunnan Province, China. Herpetozoa 34: 131–140. <https://doi.org/10.3897/herpetozoa.32.e69755>
- Liu S, Lwin YH, Quan R, Li S (2020) First record of *Rhacophorus verrucopus* Huang, 1983 from Myanmar. Herpetozoa 33: 207–211. <https://doi.org/10.3897/herpetozoa.33.e60214>
- Mathew R, Sen N (2010) Pictorial Guide to Amphibians of North East India. Zoological Survey of India, Kolkata, 144 pp.
- Ohler A, Delorme M (2006) Well known does not mean well studied: morphological and molecular support for existence of sibling species in the Javanese gliding frog *Rhacophorus reinwardtii* (Amphibia, Anura). Comptes rendus biologies 329: 86–97. <https://doi.org/10.1016/j.crv.2005.11.001>
- Savage JM, Heyer WR (1997) Digital webbing formulae for anurans: a refinement. Herpetological Review 28: e131.
- Sen N (2004) Further notes on state-wise distribution of the amphibian fauna of North East India. Records of the Zoological Survey of India 102: 105–112. <https://doi.org/10.26515/rzsi/v102/i3-4/2004/159512>
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis version 11. Molecular Biology and Evolution 38(7): 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Torreilles S, McClure D, Green S (2009) Evaluation and Refinement of Euthanasia Methods for *Xenopus laevis*. Journal of the American Association for Laboratory Animal Science (JAALAS) 48: 512–516.
- Zug GR (2022) Amphibians and reptiles of Myanmar: Checklists and keys I. Amphibians, crocodilians, and turtles. Smithsonian Contributions to Zoology 653: 1–113. <https://doi.org/10.5479/si.19098995>

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Herpetozoa](#)

Jahr/Year: 2023

Band/Volume: [36](#)

Autor(en)/Author(s): Naveen R.S., Liu Shuo, Chandramouli Sumaithangi R., Babu Santhanakrishnan, Karunakaran P. V., Kumara Honnavalli N.

Artikel/Article: [Redescription of *Rhacophorus tuberculatus* \(Anderson, 1871\) and the validity of *Rhacophorus verrucopus* Huang, 1983 325-333](#)