

Notiser ur Sällskapets pro Fauna et Flora fennica,
Förhandlingar. IX. 1868. Musci novi scandinavici.
S. O. Lindberg.

1) *Bartramia* (*Glyphocarpus*) *breviseta* *Lindb.* *Synoica*,
densissime caespitosa, valde fusco-radiculosa, caule densifolio,
foliis e basi brevi et obovata fere sensim in subulam erectam
acutissimam, margine planam et serrulatam attenuatis, nervo
continuo, applanato latissimo, subulamque fere solo formante,
seta brevi crassiuscula, inter bracteas inclusa, capsula magna
suberecta et regulari, subglobosa, plicata, ore parvo, peristomio
nullo vel simplici, maxime rudimentario et irregulari
hyalino, laevissimo, sporis majuseulis, papillulosis brunneis,
operculo minuto, convexo, medio paulo elevatiore, calyptora
submitriformi, uno latere fisso. Ins. Maasöe prope Mageröe
Finnmarkiae occ. (Th. Fries. S. Henschen.)

2) *Orthotrichum scopulorum* *Lindb.* Autoicum, caule
rigido, foliis obscure viridibus, angustius ovato-lanceolatis,
obtusis, margine valde revolutis, carinatis, cellulis omnibus
chlorophylliferis, papillosis, seta sat longa, capsula magna
emergente, collo ad basin abrupto, cylindrico-oblonga, sicca
sub ore strangulata, suburceolata, jugis latis, distinctissimis,
ore lato, peristomii dentibus per paria arcte connatis, hya-
linis, laevissimis, apice grosse perforatis, cristatis et trabe-
culatis, linea mediana perforata, ciliis octo a dupli serie
cellularum, hic illic appendiculatis, calyptora magna, lurido-
straminea, nitidula, glaberrima, campanulato-conica, sporis
fusco-viridulis, pellucidis, papillulosis, androecio in ramulo
proprio apicali. — In rupib. granit. insul. Kattön, 2 mill. ex
oppid. Lovisa in prov. Nyland. Fenniae. (Autor.)

3) *Seligeria crassinervis* *Lindb.* Autoica, fusca, foliis
abrupte angustatis in subulam brevem, patulam, obtusiuscu-
lam, integerrimam et a nervo crassissimo solo formatam in-
ferne subcanaliculatam, superne plano convexam, seta 2—3
mm. longa, crassa, leniter curvata, capsula indistincte obliqua,
pyriformi, sicca suburceolata, pachyderma, dentibus peristomii
acutis. Rup. calcar. mont. Billingen prov. Westergötland
Speciae. (Cleve.)

4) *S. acutifolia* *Lindb.* Autoica, perparva, foliis viri-
dissimis, superioribus ut et bracteis perichaetii e basi plus
minus vaginante abrupte angustatis in subulam subteretem
setiformem acutissimam et pungentem fragilem crenulatam
nervo totam fere subulam formante, seta gracillima brevissima
1 mm. longa, capsula parva, apices bractearum orificio vix
superante, leptoderma, pallida et pellucida, breviter pyriformi,
collo brevi, dentibus peristomii brevibus, fere obtusiusculis,

rostro operculi brevissimo, capsula quadruplo breviore, vix obliquo. Rup. calc. ins. Gotland. (Cleve.)

B. longiseta Lindb. planta major, seta 2—3 mm. longa, capsula alte emergente, rostro operculi longiore, magis obliquo.

5) *Bryum mamillatum* Lindb. Autoicum, dense caespitosum densifolium, foliis rigidis, erecto-patentibus, latissime ovalibus, breviter acuminatis, valde concavis, limbatis, margine fere ad medium revoluto, superne serrulato nervo crasso, ut apiculo serrulato excedente, cellulis magnis, eisdem limbi in 2—3 stratis dispositis, capsula pendula, plus minusve turgido-pyriformi, brunneo-lutea, opaca, pachyderma, sicca rugulosa et sub ore minuto leniter constricta, operculo convexo-plano, mamillato, peristomii dentibus rufescensibus, processibus e membrana basilari, supra medium dentium producta, angustissimis, e dentibus liberis, ciliis subnullis, sporis permagnis, androecio in ramulo proprio apicali. In ripa arenos. amnis Gothemån ins. Gotland. — in litor. maris ad Fågelvik et Othem insulae. (Cleve.) — Ins. Aland. (Bomansson.)

6) *Anomodon apiculatus* B. S. Dioicus, obscure viridis, ramis strictis subteretibus, obtusis, foliis e basi erecta ovato-cordata et auriculata patenti-divergentibus, abrupte anguste lingulatis, obtusis, vulgo minute apiculatis, ad basin undulatis et spinoso-superne denticulato-papillosis, margine recto, nervo supra medium dissoluto, cellulis, praecipue auricularum, valde elevato-papillosis, capsula brevius pedicellata, cylindrico ovata, operculo breviter rostrato, annulo nullo. Rupes silic. mont Kissakallio ad Kaitas paroeciae Asikkala prov. Tavastland Fenniae. (Norrlin. Silén.) — Billingen. (Cleve.)

7) *Thuidium gracile* B. S. Autoicum repens, pallido-luteum, tenui, subpinnatum, paraphylliferum, caule dense radiculoso, foliis caulinis et rameis erecto-patentibus, profunde plicato-concavis, cordato-ovatis, abrupte subulato-acuminatis, acumine serrulato, margine revoluto, nervo in acumine sensim dissoluto, eisdem ramulinis ovatis, acuminatis, minus concavis, margine recto, superne serrato, bracteis perichaetii radicantis profunde plicatis, margine valde revoluto, integro, nervo crasso, in subula dissoluto, seta 1,5 cm. alta, capsula elliptica, operculo convexo-conico, alte acuteque mamillato. In radice arbor. prope Skarpneck ditionis Holmiae. (Lindb.)

(Schluß folgt.)

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Hedwigia](#)

Jahr/Year: 1868

Band/Volume: [7_1868](#)

Autor(en)/Author(s): Milde Julius [Carl August]

Artikel/Article: [Literaturbesprechung 143-144](#)