

# Engineering Geological Map

Miloš KOVÁČIK (ed.)<sup>1</sup>  
 Pavol LIŠČÁK<sup>1</sup>  
 Péter SCHAREK<sup>2</sup>  
 Tibor TULLNER<sup>3</sup>  
 Gerhard SCHÄFFER<sup>3</sup>

<sup>1</sup>Geological Survey of Slovak Republic  
 Mlynská Dolina 1, 817 04 Bratislava

<sup>2</sup>Geological Institute of Hungary  
 11-11/3 Budapest, Stefánia út 14.

<sup>3</sup>Geological Survey of Austria  
 Rastumofskygasse 23, A-1031 Vienna

## Introduction

### Types of rocks and soils

*Mostly solid rocks*

*Mostly semi-solid rocks*

*Alternation of hard and soft rocks*

*Mostly gravelly soils*

*Mostly sandy soils*

*Mostly fine soils*

*Alternation of gravelly and fine soils*

*Alternation of sandy and fine soils*

*Loess and loess-like sediments*

*Organic soils and peat*

*Anthropogenic (man made) deposits,  
 replenishment*

### Engineering geological zones

*Zone of municipal (communal)  
 wastes (Ao)*

*Zone of industrial wastes (Ai)*

*Zone of loessy sediments (L)*

*Zone of aeolian sands (E)*

*Zone of proluvial sediments (P)*

*Zone of deluvial (slope) sediments (D)*

*Zone of alluvial (fluvial) deposits (F)*

*Zone of river terrace sediments (T)*

*Zone of oxbow deposits (F's)*

*Zone of organic sediments (O)*

*Zone of organic sediments on alluvial  
 (deposits above flood plain)  
 deposits (OF)*

*Zone of aeolian sands on alluvial  
 deposits (EF)*

*Zone of clayey-silty sediments (Ni)*

*Zone of sandy sediments (Np)*

*Zone of gravelly sediments (Ng)*

*Zone of alternating clayey-silty  
 and gravelly sediments (Nk)*

*Zone of pyroclastic rocks (Vp)*

*Zone of effusive rocks (VI)*

*Zone of claystone-siltstone  
 rocks (Si)*

*Zone of flyschoid rocks (Sf)*

*Zone of conglomerate rocks (Sz)*

*Zone of sandstone rocks (Sp)*

*Zone of dolomitic rocks (Sd)*

*Zone of limestone rocks (Sv)*

*Zone of carbonate and clastic  
 rocks (Sk)*

*Zone of magmatic and intrusive  
 igneous rocks (Ih)*

*Zone of low metamorphosed  
 rocks (Mr)*

*Zone of high metamorphosed  
 rocks (Mv)*

*Zone of residual sediments (R)*



## Introduction

The Engineering Geological Map is based on the genetic and lithological classification of rocks and soils. The map has been constructed on the basis of the Surface Geology Map 1:100 000 of the DANREG area and the basic principles are in accordance with the guidelines given by the International Association of Engineering Geology [Annon. 1976, MATULA *et al.* 1979, MATULA (ed.) 1981]. The map 1:200 000 displays the engineering geological zones and the important tectonic features. In order to prevent the duplicity of data displayed in other DANREG maps (Tectonic Map, Neotectonic Map, Environmental Geohazards Map, and Hydrogeological Map), only limited data are shown in Engineering Geological Map. The geodynamic phenomena are shown in the Environmental Geohazard Map: the slope deformations, erosional and accumulation features, subsidence manifestations, sagging of loesses and karstic features, also including important major earthquake events. Neotectonic phenomena are shown on the Neotectonic Map. The hydrogeological data on the Environmental Geohazard map show the depth of water level and the chemistry of the groundwater, and partly the hydroisobaths and important springs. Another reason for this concept was to render the map as readable as only possible. Some endogenous and exogenous geodynamic data, the hydrogeological conditions and some environmental features which usually appear in the standard map are described in the text. During the compilation of our map we were collecting and using the earlier geological and engineering geological maps as well of the concerned area. These were:

— Austria: BRIX & PASCHER 1994; FUCHS 1965, 1984, 1985a, b, c, d; GRILL 1968; HERMANN *et al.* 1993; HOCH & HACKER 1971; HUBER 1978; JELEM 1975; KAPOUNEK & FINK 1978; KÜMMEL *et al.* 1957; Österreichisches... 1979; SCHMID 1970, 1972; VETTERS 1910; WESSELY 1968.

— Slovakia: MATULA *et al.* 1989, NEMÉOK 1982, AJGALÍK & MODLITBA 1983.

— Hungary: OVH 1984; RAINCSÁKNÉ KOSÁRY (ed.) 1984; SCHAREK (ed.) 1990, 1991 a, b, 1993; SCHAREK & TULLNER (eds) 1992, 1993.

The engineering geological zones (pre-Quaternary basement, or Quaternary deposits) are regional units with identical or similar genesis and lithology [FODORNÉ (ed.) 1971, FODORNÉ & KLEB 1986]. Rocks and soils down to the depth of 5 metres were taken into account. The boundaries of zones are shown on the map in full black line and marked by using symbols (*e.g.* VI, or D etc.).

Different colours indicate the characteristic lithological composition of the uppermost 5 m of the section. Altogether 11 colours appear in the map representing the following lithologies: mostly solid rocks (with the uniaxial strength > 50 MPa), mostly semi-solid rocks (with the uniaxial strength < 50 MPa), alternation of hard (solid) and soft rocks, mostly gravelly soils, mostly sandy soils, mostly fine soils, alternation of gravelly and fine soils,

alternation of sandy and fine soils, loess (and loess-like sediments), organic soils and peat, and anthropogenic (man-made) deposits.

The zones are described in brief in the following text. Only limited data of the geotechnical issues were available on parameters from the Austrian project area. Slovak technical standards (STN) were used for characterisation of some geotechnical parameters. Sporadic investigations using the Schmidt Hammer Test were carried out, to improve the correlation of the rocks. The fracture zones were partly checked in the field and verified also from the satellite and airborne images.

A brief description of the engineering geological characteristics is given in Tab.1. The following items are described: (1) Symbol of zone, (2) Name of the zone, (3) Lithological content and age of rocks/soils, (4) Hydrogeological conditions, (5) Geomorphological conditions, geodynamic phenomena, (6) Geotechnical classification of rocks/soils, (7) Geotechnical description of rocks/soils, (8) Stability of slopes/cuttings, (9) Usage of rocks/soils, raw materials, (10) Recommendations for constructions, environmental hazard.

## Types of rocks and soils

### *Mostly solid rocks*

These are shown in dark-blue colour on the map and include Palaeozoic crystalline (both intrusive and metamorphosed) rocks, Mesozoic limestones, dolomites, sandstones, as well as Miocene carbonate rocks and volcanics essentially consisting of andesite lavas. Although different in their origin, the “mostly solid rocks” feature uniformly favourable uniaxial strength characteristics (> 50 MPa).

The rock types are found in the uplifted areas of the Hundsheim Mts, (including the Braunsberg), the Leitha Hills, and the Rust Upland in Austria. The Malé Karpaty Mts. represent a large occurrence of this type of rocks in Slovakia (mostly igneous, metamorphic and carbonate rocks). The largest area of solid rocks is in the Eastern part of the DANREG area (Krupinská planina in Slovakia, and the Börzsöny Mts and Pilis Mts in Hungary).

### *Mostly semi-solid rocks*

This group includes semi-solid (soft) rocks with uniaxial strength lower than 50 MPa. It comprises sandy sediments and the claystone facies of Palaeogene basin sediments (in the Austrian part of Badenian and Pannonian ages).

Occurrences: West slope of the Hundsheim Mts., in the area surrounding of the Arbetsal Upland and their easterly hills, the area of the Leitha Hills, the area between the Leitha River and the Parndorfer Platte (Austria) and the area south of Sopron (Hungary).



waste disposal

disposal

Kf = hydraulic conductivity

*Alternation of hard and soft rocks*

Non-uniform formations of variable lithological and stratification pattern are assigned to this group. The uniaxial strength of this type of rock is lower than 50 MPa. They incorporate Palaeozoic low-grade metamorphic rocks — gneiss and micaschist series, Lower Miocene and Pannonian sediments bearing sandstone banks, as well as Badenian and Lower Pannonian marls and sandstones.

Rocks crop out in the Leitha Hills in Austria and along the northern periphery of the Bakony Mts, Quaternary freshwater limestones in the Gerecse Mts and in Sopron Hills in Hungary.

*Mostly gravelly soils*

They are depicted by dark-green colour. They represent terrace formations of the Danube River and its tributaries uplifted to different altitudes. Lithologically, sandy gravel occurs most commonly with lenticular and cross-bedded intercalations of medium- and coarse-grained sand, including layers of organic clay along the Danube River. Sometimes the gravel is covered by loamy sediments up to several metres (*e.g.* the Gänserndorf terrace in the North of the Marchfeld including the Schlosshofer Platte).

*Mostly sandy soils*

Typical occurrences of sandy soils are shown in yellow colour. This group consists essentially of fine-grained aeolian and fluvial sands. The deluvial slope deposits, which were derived from Pannonian sediments also belong into this category.

Larger occurrences are known from Marchfeld, from the Hundsheim Mts, from Seewinkel, from the northern and western part of the Braunsberg Mt and from spots along the Danube (mostly its right bank in Hungary) and Little Danube rivers throughout the whole DANREG area. There are essential occurrences of aeolian sands in the Žitavská and Pohronská pahorkatina uplands in Slovakia.

*Mostly fine soils*

Apart from unconsolidated Upper Pannonian formations which consist of sandy silt and aleuritic loess, sandy loess and loess and fine sand assemblages belong to this group. Undivided fluvial sediments (NW of Sopron in the region of the border) contain mostly fine soils. Due to its specific textural features, loess is quite different from silts which are deposited in water with the same grain-size distribution pattern. Loess and loess-like sediments are therefore distinguished by a specific code sign.

Fine soils occur along the rivers Leitha and Fischa, in the surroundings of Lake Neusiedl in Austria, they cover the very most part of the Podunajská nížina Lowland in Slovakia, occurring along the River Hron and also cover large parts of the Kisalföld (Little Hungarian Plain) in Hungary.

*Alternation of gravelly and fine soils*

Holocene alluvial deposits in basins and intramountain valleys (Danube, March, Wulka), are assigned into this category. Gravels are quite common at the base of their typical section overlain by sand and flood-laid silt, rich in organic matter. Consequently, the thickness of this unit is mostly intermediate and varies from site to site.

This type of deposit occurs mostly along the rivers Danube and March. The gravels are overlain in the flood plain of the tributaries of the River Danube by up to 5 m offlood plain clay and silt and along the River March by up to 3 m of loam.

*Alternation of sandy and fine soils*

Fine-fraction members of loose, slightly reworked proluvial fills (without matrix) of dry valleys are combined in this group. Also included are deluvial clay, silt and sand, and lacustrine-paludal sediments. By their specific lithology they can easily be distinguished from the basement.

The occurrence of this lithological type is rare. It was documented on local faint morphologic highs in the Seewinkel area in Austria.

*Loess and loess-like sediments*

Loess and loess-like sediments are always distinguished from fine (aleuritic/silty soils) due to their specific engineering geological properties. They are mostly of aeolian and also polygenetic genesis. Typical loess is relatively homogeneous, bedded soil with high content of silt fraction (0.05–0.005 mm), calcareous and porous. The sediment is shown in yellow-brown colour in the map.

Loess and loess-like sediments cover large parts in Slovakia and Hungary mostly along the Danube River.

*Organic soils and peat*

The typical locality for organic clay, as well as peat and muck assigned to this category occurs around the Lake Neusiedl/Fertő and the Hanság. The maximum thickness is in the Austrian part between 1 and 2 m. They are commonly underlain by sand and gravel. The whole assemblage is highly irregular in extent and thickness and has low and very low bearing capacity. This assemblage occurs locally in some small, abandoned river channels in the Szigetköz area and in a number of intramountain basins without an outlet.

Organic soils are common in ox-bows along the Little Danube and Hron rivers in Slovakia and along the whole main course of the Danube River. There is a relatively large occurrence of peat east of the Malé Karpaty Mountains as well.

*Anthropogenic (man-made) deposits, replenishment*

These occur commonly in the vicinity of large cities (Wien, Bratislava, Budapest, Győr) and industrial areas



(Tatabánya, Esztergom), as well as in operating or abandoned pits. Owing to their alternating composition they have generally irregular low to very low load-bearing capacity. In open pits and central districts of capitals their thickness can exceed even 10 m.

### Engineering geological zones

29 zones have been delineated in the map: 13 zones of Quaternary deposits and 16 zones of pre-Quaternary basement. When the knowledge of the study area allowed to detach two zones in superposition, combined zones were marked in the map (OF and EF).

The zones are the following: zone of municipal wastes (Ao), zone of industrial wastes (Ai), zone of loessy sediments (L), zone of aeolian sands (E), zone of proluvial sediments (P), zone of deluvial (slope) sediments (D), zone of alluvial deposits (F), zone of river terrace sediments (T), zone of oxbow deposits (Fs), zone of organic sediments (O), zone of organic sediments on alluvial deposits (OF), zone of eolian sands on alluvial deposits (EF), zone of clayey-silty sediments (Ni), zone of sandy sediments (Np), zone of gravelly sediments (Ng), zone of alternating clayey-silty and gravelly sediments (Nk), zone of pyroclastic rocks (Vp), zone of effusive rocks (Vi), zone of claystone-siltstone rocks (Si), zone of flyschoid rocks (Sf), zone of conglomerate rocks (Sz), zone of sandstone rocks (Sp), zone of dolomitic rocks (Sd), zone of limestone rocks (Sv), zone of carbonate and clastic rocks (Sk), zone of magmatic and intrusive rocks (Ih), zone of low-grade metamorphic rocks (Mn) and zone of highly metamorphosed rocks (Mv), zone of residual soils (R).

#### *Zone of municipal (communal) wastes (Ao)*

This zone is composed of isolated areas in the vicinity of large settlements. The landfill material consists mainly of: household wastes, ash, clinker, local industry wastes, redundant soils and construction debris. Although, the thickness of wastes varies, it is generally less than 10 m. The landscape depressions (such as pits, ox-bows, etc.) have been used generally to dispose of local wastes. Even grading and recultivation of the surface render the area within this zone hardly suitable for construction works. Its unsuitability is due to high compressibility, irregular sagging, low bearing capacity and health hazard, or even danger (methane).

#### *Zone of industrial wastes (Ai)*

It incorporates technological wastes of industrial estates (frequently involving dangerous elements) and overburden of the mining branch. The material is very variable, mostly inert rocks of various granulometric composition. The thickness of these accumulations reaches 10 metres, locally even more. Specific cases are the wastes from the chemical industry, with special physical and

mechanical properties and chemical composition. These require special care because of their possible ecological harmfulness. The zone is not suitable for any type of construction.

#### *Zone of loessy sediments (L)*

The loess is widely distributed in a thickness of several tens of metres. In the Austrian part of the project it is mostly less than 5 m. The thickness of the loess sediments in Slovakia vary between 5 and 15 m (greatest thickness in Slovakia occur in the surroundings of Svodín (up to 40 m). Lithologically variable, the zone is a sequence of aeolian and fluvio-lacustric-aeolian deposits, as well as fossil soil horizons.

The age, genesis, thickness as well as the fossil soil horizons are partially varied, while the age, genesis, thickness, character and properties of loessy sediments are varied. The loess and loam was formed mainly during the Middle and Late Pleistocene, with the greatest representation during the Mindel, Riss and Würm glacials. Typical loess is a relatively homogeneous bedded soil with a high content of silt fraction (0.05–0.005 mm). The loess is calcareous and very porous. The porosity and the bearing capacity decrease with age of the sediments. Sagging depends on the thickness of this zone. The colour is mostly yellow to pale-brown.

Loess is the most common of all Quaternary sediments in the DANREG region. Larger amounts of loess cover the area in the N of Gänserndorf terrace (Oberweiden, Strasshof ad. Nordbahn), near the airport Vienna (Schwechat), Schwadorf and S of Sopron. Loess sediments (typical loesses and loessy loams) occur often along the Danube River on both riverbanks in Slovakia and in Hungary. Extensive deposits occur in the Trnavská, Nitrianska, Žitavská, Pohronská and Ipeľská pahorkatína uplands. They are subordinately present over the Záhorská nížina Lowland, together with fluvial sediments.

From the hydrogeological point of view the zone is important to protect groundwater horizons. Hydrocompaction occurs sometimes while the scour erosion (on slopes), or lateral erosion occurs to a limited extent only.

The loess is suitable for brick-making and pottery. The zone is suitable, or limitedly suitable (sagging, furrow/gully erosion) for building construction.

#### *Zone of aeolian sands (E)*

The deposit is fine-grained sand (fraction 0.5–0.25 mm prevails), well sorted, with predominance of quartz (more than 90%). The thickness of the zone varies from a few metres up to 25 m (Marcelová). In Austria it is mostly less than 5 m. The sediment has medium bearing capacity. The morphology depends on the preservation of sand dunes. When preserved, sand dunes are suitable for building construction especially in inundation areas.

The zone of aeolian sands occupies the southern part of the Záhorská nížina Lowland and isolated areas in Žitn

ostrov, in Rábaköz in the proximity of the Danube River. The aeolian sand is well permeable.

Due to the predominantly low thickness and limited extent of the sands, pore water is usually absent. Accumulations of water can be expected to occur in thicker sand beds, within depressions of the Záhorská nížina Lowland.

Wind erosion (deflation) is increasingly active due to the removal of vegetation (exploitation of wood or agriculture in the Záhorská nížina Lowland).

Aeolian sands can be used for building construction (cement), for casting, or for the glass industry. The zone comprises mostly dryer, inferior site index soils. The zone is unsuitable for landfills.

#### *Zone of proluvial sediments (P)*

This zone includes sediments with poorly rounded components deposited in dry valleys with seasonal flooding. The sediment consists of coarse-grained gravels alternating with sandy and loamy beds. They may be seldom boulder-size, like at the foothills of the Malé Karpaty Mts. The gravels are generally unsorted, the pebbles are semi-rounded to angular. The thickness of the sediments varies between 2 and 10 m. The slope inclination has a range of 1.5–5 degrees. Scour erosion and landsliding affect the slopes dipping 4–5°.

The zone occurs in the Parndorfer Platte in Austria, W and E of the foothills of the Malé Karpaty Mts and in the smaller Pleistocene and Holocene accumulations from side valleys along the Little Danube River, at the foothills of Krupinská vrchovina Highland in Slovakia.

Significant water-bearing can be expected only in thicker gravelly horizons. In general, proluvial sediments are more favourable for the concentration of groundwater than the deluvial ones.

The zone is only limitedly suitable for building constructions. One can expect an uneven compressibility within the zone. The contained clastic fraction will increase the strength after compaction. The zone is suitable for landfills. Flat slopes are suitable, or conditionally suitable for any kind of building construction.

#### *Zone of deluvial (slope) sediments (D)*

Slope deposits of mountainous and hilly regions belong to this zone. The composition of slope loams reflects the lithology of the basement rocks. If the basement is made up of solid or semisolid rocks, the loam has a variable content of fragments.

The deluvial sediments overlying Neogene gravels usually contain pebbles. The thickness of loams ranges between 2 and 10 m. Higher thickness is rare. The thickness of loams increases downhill. Hard rocks (limestones, dolomites, and granodiorites) are overlain by thin deluvial loams. In case of semisolid rocks (such as claystones, tuffs *etc.*) the loamy cover is thicker. Depending on the underground, the inclination of slopes rarely exceeds 20–25 degrees. The deluvial sediments related to volcanic rocks

contain a higher percentage of clay particles. The deluvial sediments in the Malé Karpaty Mts more often contain clastic components. Scour erosion and also landsliding occur locally. Owing to their lithological variability, the compressibility in the zone is irregular. Most of the sediments of this zone are highly congelative (frost susceptible).

Only small occurrences of deluvial deposits were identified in Austria (*e.g.* in the depression south of the Leitha Hills in the N of Winden). Deluvial slope loams occur at the foothill of the Malé Karpaty Mts and in the Krupinská vrchovina Highland and at foothills of the Börzsöny and Pilis Mts.

The deluvial loams are highly water-bearing. Springs at the base of the slopes stop in most cases during the dry seasons.

This zone is used mostly as pasture or arable land.

#### *Zone of alluvial (fluvial) deposits (F)*

The zone of fluvial deposits is the dominant zone in the region. It covers at least 40% of the total DANREG area (estimation). It is present over the whole braided river-system (Danube, Váh, Hron, Ipeľ/Ipoly, Žitava *etc.*). The zone incorporates alluvial deposits of fluvial plains and intramountain valleys. These have a mixed lithological composition (gravel, sand, fine particles). The total thickness of the deposits ranges between 5–10 m. The flood deposits are clayey to loamy. The gravels are coarse and loamy. The river deposits show a wide range of grain size categories. The plasticity ranges from low to high. The fine sediments are of variable consistency. The morphology is completely flat (in most cases does not reach even one degree).

Logically the zone covers areas along main rivers of the region: Danube, Leitha, Fischa, Little Danube, Nitra, Hron, Ipeľ/Ipoly *etc.*

The alluvial gravel is an important aquifer. The maximum and minimum of groundwater level is depicted on the Environmental Geohazard Map. Due to the occurrence of aquifers this zone is used for exploitation of groundwater and is often protected from any other activities except agriculture.

Mixed sand and gravel are suitable as infillings, or for road construction. Pure gravels are suitable for stabilising of gravity banquettes. The alluvial gravels are used for local building construction.

#### *Zone of river terrace sediments (T)*

The terrace sediments along the main rivers and their tributaries are composed of Pleistocene gravels and sandy gravels. The younger terraces have the largest areal extent and are dated as Würm and later. Most of the older terrace sediments have been eroded. The sediments contain coarse gravels and sands. The main fraction of the terrace gravels is medium- to coarse-grained. The older terrace sediments are highly weathered. The surface of terraces is flat. The

boundaries towards the surrounding zones are morphological visible.

Terrace sediments cover the majority of the Austrian part of the map. Along the main water courses (Danube, Váh, Hron, Ipeľ/Ipoly, Zitava, etc.) the terrace steps at various levels are preserved.

The groundwater is usually bound to the proximity of the gravel base (some 5 m below the surface). The permeability of the gravels varies, depending on loam and clay admixtures. The zone offers a good quality for concrete addition.

The area is almost unsuitable to dispose waste (due to local high permeability). The reality is adverse. There are a lot of illegal local dumps there, mostly with household waste.

#### *Zone of oxbow deposits (Fs)*

The oxbow fill is composed of sediments of variable consistence (solid clay to slurry) with a high content of organic matter. The thickness of the sediments exceeds often 5 m. The oxbow zone represents morphologic lows in the lowlands. The peat deposits occur locally in this zone. The fill in the remnants of dead channels in the Szigetköz area is characterised by silts with high organic content. This zone is delineated by the Danube River and the Mal' Dunaj River (Little Danube) and the Leitha River. This zone includes isolated lake areas as well.

The groundwater level is seasonally fluctuating, in pace with the rate of flow in the rivers and their inundated areas.

Certain areas are used for agricultural purposes in order to preserve the fertile soils. The groundwater is close to the surface. Therefore this places are not suitable to store waste (as it was done in the past).

#### *Zone of organic sediments (O)*

Isolated areas in the Podunajská nížina Lowland, adjacent to the main course of the Danube and Mal' Dunaj rivers, belong to this zone. The occurrences of organic silt, clay and peat (5–10 m thick) are typical for this zone. They are present mainly in the Hanság and Rába Valley and around the Lake Neusiedl/Fertő. Minor deposits of organic sediments occur in the zone of oxbow lakes. In Austria the maximum thickness is 2 m. The top of the groundwater sometimes rises above the surface (swamps). Because of their fauna and flora, the areas with peat and bogs are often protected due to the rare flora and fauna associations. The peat has been used as a fertiliser in the agriculture.

#### *Zone of organic sediments on alluvial (deposits above flood-plains) deposits (OF)*

This zone includes young, uncompacted sediments with variable amounts of organic matter occurring above flood-plains, and at the periphery of flat areas, that are filled with peat. In the Podunajská nížina Lowland, there

is an isolated zone adjacent to the Danube and Mal' Dunaj river valleys. Minor zones of organic sediments occur also in the zone of oxbow lakes. Occurrences: in the Hanság area and around Lake Neusiedl/Fertő.

#### *Zone of aeolian sands on alluvial deposits (EF)*

This zone has several small occurrences in Záhorská nížina Lowland in Slovakia. There are wind-blown sands lying over alluvial deposits of the Morava River. The thickness of sands is up to 5 m. The other characteristics are identical with those of zones 3.4 and 3.7.

#### *Zone of clayey-silty sediments (Ni)*

The zone is represented by fine-grained Pannonian compact rocks with variable content of clayey, silty and sandy (in the eastern part often tuff) sedimentary rocks. Sporadically, they include beds with carbonate matrix. Gravel beds occur occasionally. Occurrences in Austria are concentrated especially in the N between Parndorf and Potzneusiedl.

The rocks of this zone have varied consistency. The Pliocene clays are soft while the oldest Badenian sediments are compact to hard. The plasticity index is generally high to very high and increases with the distance from the mountains margins. In compliance with the Standard STN 73 1001 the compacted soils in this zone can be assigned to classes F6 to F8. According to the Standard STN 73 3050 we rank the mentioned rocks in the 3<sup>rd</sup> exploitability class. The most unfavourable engineering geological property of the rocks in the zone is their generally risky congealing capacity and their overall increased susceptibility in contact with water.

With regard to the permeability of the soils, we rank them as aquicludes with a very low permeability coefficient.

The weathering susceptibility is high. If a slope is undercut, either by river erosion or due to human activity slope movement might be triggered. Such geodynamic phenomena can be observed, e.g. in the area between Sered and Hlohovec. At this location the Váh River has undercut the adjoining slope, which is covered by compacted Neogene sediments. The slope movement disrupts the road body and jeopardises the buildings. Similarly, the landslide near Viťtuk was caused by the action of the Viťtuk potok brook. In this case, the road surface has been damaged and the dam of the Viťtuk water works has been threatened. In both cases the landslide bodies were more or less drained by means of dewatering boreholes. In the area north of Sered a monitoring system has been established.

The rocks of the zone can be used for building constructions and for brick-making (Devínska Nová Ves, Pezinok etc.) as well as for blending clays in for the cement production (in Rohoňník). The thickness of clays and absence of sandy intercalations is generally sufficient to assure safe waste disposal.

### *Zone of sandy sediments (Np)*

The sand beds of the Pannonian are unconsolidated. They are mainly well-stratified and bear intercalations of sandstone with carbonate matrix at the bottom. This zone also includes unconsolidated or slightly consolidated Neogene sediments. Occurrences in Austria are of only minor importance. The Pliocene sands are moderately compacted to compacted. The Miocene sands are always compacted. The sands are often cemented by carbonate or clayey matrix. The bearing capacity of Neogene sands increases in accordance with the distance from the mountains and with the higher clay content.

According to the Standard STN 73 1001 the rocks of the mentioned zone belong to the classes S1 to S3 and the sands with clayey admixture to classes S4 up to S5. In compliance with the Standard STN 73 3050 we rank them into 2<sup>nd</sup> and 3<sup>rd</sup> mineability classes.

The permeability coefficient of the sands in this zone ranges from  $10^{-7}$  to  $10^{-4}$  m/s<sup>1</sup>. It decreases in function of the increasing compactness of sands (porosity drop) and with the increase of clay content, or silty fraction. The groundwaters in Neogene sands from the margin of the Malé Karpaty Mts are slightly mineralised. The groundwaters of the Podunajská nížina Lowland are often characterised by sulphate aggressiveness. The sulphates derive from agricultural activities.

The area covered by Neogene sands is characterised by a very smoothly modelled morphology. Only river erosion causes steep slopes with landslide scars. The stability of the slopes is threatened during technical works (cuts and excavations).

The area of the Záhorská nížina Lowland is generally afforested, while the Podunajská nížina Lowland is cultivated by agriculture. If the sands are less compacted and the groundwater is close to the surface, the zone offers foundation soils unsuitable or conditionally suitable for building constructions. The congelation (frost sensitivity) of Neogene sands with higher content of clay has to be investigated with respect to road construction. The rocks of this zone are unsuitable for waste dumping.

### *Zone of gravelly sediments (Ng)*

Neogene gravels are more compacted than the younger terrace gravels. The material is mostly strongly weathered. In comparison to Quaternary gravels, they contain minor part of loamy admixture and are coarser.

This zone is areally rather restricted. It occurs locally at the base of the Pannonian deposits with gravels west of Fertőrákos and in minor occurrences in Austrian part.

Rill and gully erosion is relatively intense on the slopes.

Usually coarse, sandy gravels are sources of groundwater.

Material is suitable for infillings, gravity banquettes and for road construction. Due to the weathered material, their use for making concrete is only limited.

### *Zone of alternating clayey-silty and gravelly sediments (Nk)*

The zone includes mainly terrestrial Oligocene–Miocene sediments of varied composition, as well as formations underlying the volcanic complex of the Börzsöny Mountains, at the margins of the Malé Karpaty Mts, and in the area of the Krupinská planina Plain. The Volkovce, Beladice, Ivanky, Záhorie, Vráble and Studienka formations are part of this zone. In Austria this zone is present in the southern part of the Leitha Mts. (Auwaldschotter).

The rocks of this zone are characterised by almost chaotic alternations of beds and lenses of clays, sands and gravels with various ratio and grain size fraction. From the engineering geological point of view, the thickness and the areal extents are too small to allow differentiation among the lithological types on the small scale map. The individual lithological types are in compliance with the soils in the zones Ni and Np.

Although the gravelly and sandy beds have the highest permeability, the largest yields can be expected in the zone that is in contact with permeable underground or surroundings.

The zone is characterised by smooth slopes, locally disturbed by out-wash erosion. The slopes might be disrupted by erosional impact or by inappropriate building constructions.

The utilisation of the zone depends on the lithological type corresponding to zones of sandy or fine-grained sediments.

### *Zone of pyroclastic rocks (Vp)*

It occupies eastern part of the area under study. Various types of pyroclastic rocks are represented — sandy and aleuritic tuffs and tuffites, tuffitic sandstones and pyroclastic breccias.

The rocks are almost exclusively semi-solid, with uniaxial compressive strength values between 2 and 40 MPa.

Quite distinct is the susceptibility of the mentioned rocks, mainly of syngenetically, or postgenetically altered ones, to weathering. The intensity of weathering causes in several cases that the rock breakdown to form sand. Fresh rocks we assign, according to the Standard STN 73 1001, into classes R3 and R4, and in case of intense weathering into R5 and R6.

The water-bearing of the pyroclastic rocks is conditioned by their genesis and grain size. Pore permeability predominates, but the fissure permeability is also important. Mineral springs also occur. The aggressiveness is usually of the hydrocarbonate type.

Despite the fact that the relief is characterised by smooth slopes, the dissection is relatively high. A decisive factor, responsible for the breakdown of the Krupinská planina Plain into discrete blocks, is the outwash, generally selective erosion. The andesite pyroclastics in the Burda area are being quarried as construction materials. The zone is predominantly afforested, the site index of soils ranges between 3 and 7. It offers good conditions to found build-



ings, a limiting factor may be the outwash erosion. If sufficiently large and thick, the silty and pelitic tuffs and tuffites provide good environment for waste disposal.

#### *Zone of effusive rocks (VI)*

The rocks of this zone occur mainly in the eastern part of the area under study, in the Krupinská planina plain, Burda Mts in the Slovak part, in Börzsöny and Pilis Mts. The main types of rocks are extrusive and intrusive andesites of the Mátra, Vinica and Burda Formations.

Depending on the genesis, mineral composition and postgenetic alterations the rocks of the zone show distinct variations in physical and mechanical properties. The compressive strength of "sound" andesites reaches up to 200 MPa. According to the Standard STN 73 1001 we rank such rocks into R1 class, the mineability class according to the Standard STN 73 3050 is 7. Altered rocks of the zone are of semisolid rocks character in some places.

The relief is relatively intensely dissected and the exposed lava flows create usually positive relief.

Among geodynamic phenomena, the weathering of hydrothermally altered rocks occurs.

The water-bearing of the zone depends on tectonic jointing. Shallow circulation of groundwaters is typical for the weathering zone, the springs have relatively low yield due to the small areal extent of the zone.

The andesites at the Vinica and Hrušov sites are mined for local purposes and for road maintenance. The area of the zone is afforested. Due to jointing the rocks of the zone are unsuitable for waste disposal. The conditions for constructions are limited because of the dissected relief.

#### *Zone of claystone-siltstone rocks (Si)*

The zone includes the Eocene and Oligocene marine marl and sandstone sequence. Also Miocene formations with varied lithological composition contain some sequences belonging to this category. Mesozoic rock types of this zone belong to the Marianka formation (also including sandstones, marls and limestones). They occur at the western margin of the Malé Karpaty Mts, between Marianka and Turecká vrch Mts. The rocks are black shales with sandstone and crinoidal limestone intercalations, known as the Mariánka shales. Occurrences in Austria are known from the slopes of the Leitha Hills, in the surrounding of the Arbustal Hills and in the area of Neckenmarkt.

The uniaxial compressive strength reaches up to 60 MPa. Therefore they belong to the category of solid rocks. According to the Standard 73 1001 the rocks of this zone belong into class R2 and according to the Standard STN 73 3050 into the classes 4 and 5. The shaly beds are less resistant against weathering.

The clayey shales are practically impermeable. The clastic limestone intercalations with fracture permeability are negligible, due to their limited distribution.

At the W margin of the Malé Karpaty Mts the zone has a smooth morphology. Erosion occurs due to the effect of water and weathering. Most of the zone is afforested. The zone offers suitable conditions for building constructions. The conditions for waste disposal are to some degree favourable.

#### *Zone of flyschoid rocks (Sf)*

To this zone we have assigned tiny occurrences of the flyschoid series of the Borinka sequence in the Slovak part of the DANREG area, represented by alternation of sandy limestones and sandstones and silicites. The properties of individual engineering geological types are described in zone of sandstone rocks and zone of limestone rocks.

#### *Zone of conglomerate rocks (Sz)*

This zone includes the conglomerates and volcanic tuffs of the Krupinská planina plain, the breccias and conglomerates of the Devínska Nová Ves Member (Badenian) and the feldspathic greywackes, arkoses and polymictic Permian conglomerates of the Devín Formation as well as several minor occurrences in Austria and Hungary.

In compliance with the Standard STN 73 1001 they are ranked into class R2, and in compliance with Standard STN 73 3050 into classes 5 and 6. The technical properties of the zone are similar to those of the zone of sandstone rocks (Sp).

#### *Zone of sandstone rocks (Sp)*

We include into this zone Neogene sandstones with carbonate matrix between Sopron and Fertőrákos, the quartzites and arkoses (hard rocks) of the Lower Triassic, the lower terrigenous envelope unit of the Malé Karpaty Mts (Lúžna Formation), which occurs in isolated outcrops in the area of Devínska Kobyla and W and NW of Pezinok. Furthermore, the Jurassic siliceous sandstones and bioclastic limestones at the W margin of the Malé Karpaty Mts are also included. In Austrian part of the area under study only minor occurrences are known.

The Middle Triassic quartzites and arkoses for example are characterised by a very high uniaxial compressive strength reaching as much as 200 MPa. According to the Standard STN 73 1001 the rocks of this zone are assigned into class 1 and according to the Standard STN 73 3050 into the classes 6 and 7. The permeability of the mentioned rocks is relatively high, but due to the limited distribution of the zone the yield of springs is low. A highly uneven morphology is typical (Traja jazdci in the Malé Karpaty Mts).

The zone is afforested. It provides good conditions for building constructions, but it is unsuitable for waste disposal.

### *Zone of dolomitic rocks (Sd)*

The Triassic dolomites in Austria and Slovakia are comprised in this zone. The dolomites have usually thick banks. The exposed surfaces are strongly affected by weathering. Other characteristics are similar to those of the zone of limestone rocks.

### *Zone of limestone rocks (Sv)*

Several lithological rock types belong to this category: In Austria the Dachstein Limestone, the Lower Jurassic limestones, the Miocene reef limestones and the Pliocene–Pleistocene freshwater limestones. Miocene limestones occur mainly in the surroundings of the Leitha Hills and in the Rust Hills. Minor occurrences are found on the NW margin of the Hundsheim Mts limestones on the Braunsberg Mt, the hill with the castle of Hainburg and the western part of the Hundsheim Mts. In the Mesozoic carbonate rocks of the Slovak part of the DANREG area the limestones prevail over dolomites. The zone has a minor occurrence in the Malé Karpaty Mts, east of Borinka. Lower Jurassic limestones occur in the Borinka sequence, of the Malé Karpaty envelope unit. Dolomite intercalations occur locally in limestones.

We rank the limestones among the hard rocks: uniaxial compressive strength reaching approximately from 90 to 150 MPa. According to the Standard STN 73 1001 they belong to the classes R1 and R2, while the Standard STN 73 3050 ranks them into class 6.

Permeability is connected with jointing or karst phenomena.

The limestones in the quarry East of Borinka are resistant to weathering. They contain tectonically affected zones and karstic features. They are used as underground material for roads and railways. The zone is afforested. It provides a suitable foundation soils for massive building constructions, but it is not suitable for waste disposal.

### *Zone of carbonate and clastic rocks (Sk)*

The rock complexes in this zone, floored by a variegated sandstone-marl-limestone formation, occur in the Malé Karpaty Mts, in the area between Borinka and Turecká vrch. There are marls, sandy-crinoidal to calcareous sandstones, locally chertstones of Early Jurassic age, of the Malé Karpaty envelope unit. The rocks in this zone are characterised by slab-like jointing, and for marls a shaly breakdown is typical.

The greatest weathering resistance have the chertstones with a tendency to form positive features in the relief. The marly limestones are less resistant to weathering. The limestones and chertstones are ranked, in compliance with the Standard STN 73 1001 with the classes R1 and R2, and have the uniaxial compressive strength exceeding 100 MPa, while the marly rocks are characterised by smaller resistance (some 60 MPa) and belong to classes R2 to R4. According to the Standard STN 73 3050 we assign the rocks of this zone into the classes 5 and 6.

The permeability of the mentioned rocks is of fissure type, or fissure-karstic, respectively. The springs occur mainly where the limestones contact marls.

Several quarries are in operation in this zone. The stone from them is being used in road and railway construction. Large parts of the zone are afforested, the meadows occur to a lesser extent. The zone offers suitable conditions to found common types of constructions. To dispose wastes, the zone is mostly unsuitable, excepting the areas floored by sufficiently large marl bodies.

### *Zone of magmatic and intrusive igneous rocks (Ih)*

This zone is characterised by granites and granodiorites, to a lesser extent by diorites in the Malé Karpaty Mts which form an important part of the crystalline basement. The two-mica granite to granodiorite of the Bratislava type (often intersected by abundant pegmatite and aplite dikes) is the most common lithology. The biotitic, mostly fine-grained granodiorite occurs especially in the surroundings of Bratislava.

This type of rock is also outcropping in the Hundsheim Mts. A system of galleries was established for dumping waste. An area of approximately 10 km<sup>2</sup> is without springs. At present the rock is not used in the industry.

The amphibole — and biotite-amphibole diorites are known from the Horská park area in Bratislava.

The rocks of this zone belong to the hardest of the whole Slovak part of the DANREG region. The uniaxial compressive strength reaches 120 to 180 MPa. According to the Standard STN 73 1001 we assign the rocks of this zone into classes R1 and R2 and according to the Standard STN 73 3050 into the classes 6 and 7.

The geomorphology in the zone is high to rolling. Water erosion occurs distinctively in tectonically affected areas.

Fracture permeability is predominant, but the springs have small yields only. The groundwater is only weakly mineralised, but chemical effects can be recognised.

In tectonically affected areas outwash erosion and weathering is effective. For instance, in the area of the Devínska Kobyla Mt clayey fracture fills are known down to a depth of several tens of metres.

Several quarries located in the granodiorites and granites were operated in the past. Presently only the quarry at the southern foothill of Devínska Kobyla near Bratislava, is in operation. The rock, formerly used for building constructions, is nowadays used for roads and railways, irrigation and drainage purposes and river embankments. The zone is suitable for building construction, and conditionally suitable in the tectonically disrupted and weathered zones. The zone is to some extent suitable for dumping waste. Tectonically disturbed areas are unsuitable in all respects.

### *Zone of low metamorphosed rocks (Mn)*

In Hungary, the gneiss and mica schist complexes of the Sopron Hills belong to the zone of low-epimetamor-

phic rocks. Another small patch can be observed north of Fertőrákos at the state boundary. In Austria, small occurrences of the zone are known in the Leitha Mts and S of Sopron. In Slovakia, the rocks of this zone have only limited distribution in the area of the Malé Karpaty Mts. The rocks are regionally metamorphosed and their contact is epimetamorphic. They occur predominantly in the Harmónia series, NW of Pezinok. The bulk of them consist of sericitic-chloritic phyllites, sericitic-biotitic and biotitic phyllites, the green schists of Marianka and Harmónia successions and graphitic schists and metaquartzites and metapelites. Also the schists and schistose gneisses from the Dúbravka area belong to this zone. A small occurrence of the zone of low-metamorphic rocks is also located north of the Ipel'ské Predmostie. The rocks cropping out here are phyllites, quartzites and metadiabases of the Hron Complex of Young Palaeozoic age.

The rocks of this zone are characterised by schistose decomposition. The uniaxial compressive strength reaches 20 to 70 MPa, therefore we assign these rocks into classes R3 and R4 according to the Standard STN 73 1001 and according to the Standard STN 73 3050 into the mineability classes 5 and 6.

The permeability is of fracture type and the springs have a low capacity.

The weathering and the outwash erosion are the dominant geodynamic features in the area. The weathering takes place mainly in cuts and excavations. Due to the low resistance to weathering the use of these rocks as construction material is limited.

The zone is predominantly afforested, but meadows do also occur. The zone has suitable conditions to set up buildings, but it is only conditionally suitable to dispose waste.

#### *Zone of high metamorphosed rocks (Mv)*

The zone of highly metamorphic rocks is represented by meso- and catazonally metamorphosed rocks in the Malé Karpaty area. NW of Pezinok and in the S of

Bratislava the fine- to coarse-grained amphibolites and biotitic schistose gneisses to paragneisses has the greatest areal extent. They belong mainly to Harmónia series of the Malé Karpaty Mts. The migmatites occur at a small scale west of Svät' Jur. In contrast to the schistose gneisses, which are characterised by schistose structure, the amphibolites are massive. Minor occurrences in Austria are in the vicinity of the villages of Berg and Jois.

The tectonical decomposition of rocks is responsible for the engineering geological characteristics, such as the uniaxial compressive strength and weathering resistance. The uniaxial compressive strength of amphibolite reaches 200 MPa. According to the Standard STN 73 1001 we assign these rocks into class R1, the schists and schistose gneisses into the classes R1 and R2, and according to the Standard STN 73 3050 into the mineability classes 6 and 7.

The schistose rocks have a limited resistance to weathering. The morphology of the zone is characterised by medium to high-angle slopes, and by local rock walls.

There is fracture permeability. The springs have low capacity. The water has carbonate type aggressiveness.

The amphibolites were considered a good quality construction stone in the past. Ores occur in the area of Pezinok. The zone is predominantly afforested. It offers suitable conditions to set up building constructions. The high tectonically disrupted zones have to be taken into account while designing tunnel constructions. In this zone only restricted areas are suitable for waste disposal.

#### *Zone of residual sediments (R)*

This zone was shown in one locality — in the Leitha Hills. The weathering zone on top of the micaschists reaches up to 15 m there. It contains sporadic quartz veins.

The geotechnical characteristics are similar to the zone of deluvial —slope— sediments. The material has a character of fine soils. The plasticity is high. The textural prints of the mother rock are partly preserved.

It is suitable for waste disposal.

## References

- ANON. 1976: Engineering geological maps. A guide to their preparation. — The Unesco Press. Paris, 79 p.
- BRIX F. & PASCHER G. 1994: Geologische Karte der Republik Österreich, 1:50 000, Blatt 77 Eisenstadt. — Wien, Geol. B.-A.
- FODOR T-NÉ (ed.) 1971: Irányelvek a 10 000-es méretarányú mérnökgeológiai térképezéshez és térképszerkesztéshez. (Guidelines to 1:10 000 scale engineering-geological mapping and map plotting. — Manuscript, Geol. Inst. of Hungary.
- FODOR T-NÉ & KLEB B., 1986: Magyarország mérnökgeológiai áttekintése. (An engineering-geological overview of Hungary.) — MÁFI publ., p. 199. Budapest.
- FUCHS, W. 1965: Geologie des Ruster Berglandes (Bgld.). — *Jb. Geol. B.-A.* **108**,
- FUCHS, W. 1984: Geologische Karte von Wien und Umgebung 1:200 000. — Geol. B.-A., Wien.
- FUCHS, W. 1985: Geologische Karte der Republik Österreich, 1:50 000, Blatt 59 Wien. — Geol. B.-A., Wien.
- FUCHS, W. 1985: Geologische Karte der Republik Österreich, 1:50 000, Blatt 60 Bruck a.d. Leitha. — Wien, Geol. B.-A.
- FUCHS, W. 1985: Geologische Karte der Republik Österreich, 1:50 000, Blatt 61–62 Hainburg–Pressburg. — Geol. B.-A., Wien.
- FUCHS, W. 1985: Geologische Karte der Republik Österreich, 1:50 000, Blatt 79 Neusiedl am See, 80 Ungarisch-Altenburg. — Geol. B.-A., Wien.
- GRILL, R. 1968: Abgedeckte geologische Karte des Weinviertels 1:50 000. — In: Erläuterungen zur geologischen Karte des nordöstlichen Weinviertels, 1:75 000 und zu Blatt Gänserndorf 1:75 000. — Geol. B.-A., Wien.
- HERRMANN P., PASCHER G. & PISTOTNIK J. 1993: Geologische Karte der Republik Österreich 1:50 000, ÖK 78, Bl. Rust. — Geol. B.-A., Wien.
- HOCH, F. & HACKER, H. 1971: Kartierungsbereich Schwechat, Niederösterreich 1:25 000. — Österr. Bodenkartierung: Bodenkarte 1:25 000, KB 1. — Wien: Lw. chem. BVA. Bodenkart. u. Bodenvirt.

- HUBER, K. 1978: Marchfeld/Kleinformen, 1:100 000. — *Mitt. Komm. Quartärforschung* AW, 3, Taf. 11, Wien.
- JELEM, H. 1975: Marchauen in Niedersterreich. — *Mitt. forstl. Bundesversuchsanst.* **113**, 1–93, 2 Beil., 1 Standortkte. 1:10 000. Wien.
- KAPOUNEK, S. & FINK, I. 1978: Marchfeld/Tektonik, Karte des Tertiärsockels 1:100.000. — *Mitt. Komm. Quartärforschung* AW, 3, Taf. III, Wien.
- KÜMMEL F., FINK J., LECHNER K. & RUTTNER A. 1957: Geologische Karte der Republik Österreich 1:50 000, ÖK 107 u. 108 Mattersburg. — Deutschkreuz, mit Erl. 67S, 12 Taf., Geol. B.-A., Wien.
- Österreichisches Normungsinstitut 1979: Belastungsannahme im Bauwesen an nicht schwingungsfähigen Bauwerken. — NORM B 4015 Teil 1/1979, Wien.
- MATULA, M. et al. 1979: Classification of rocks and soils for engineering-geological mapping. Part I: Rock and soil materials — *Bulletin of Int. Ass. of Eng. Geol.* **19**, 364–371.
- MATULA, M. (ed.) 1981: Rock and soil description and classification for engineering-geological mapping. Report by the IAEG Commission on Engineering geological mapping — *Bulletin of Int. Ass. of Eng. Geol.* **24**, 235–274.
- MATULA, M. et al. 1989: Atlas inžinierskogeologick'ch máp SSR 1:200 000. (Atlas of engineering-geological maps of Slovakia.) — Katerda inžinierskej geológie PFUK, 55 p., Bratislava.
- NEMÉOK, A. 1982: Zosuvy v Slovensk'ch Karpatoch. (Landslides in the Slovak Carpathians.) — Veda, vydavateľstvo Slovenskej akadémie vied. 319 p. Bratislava.
- OVH 1984: Vízgazdálkodási Keretterv. (Base Plan of Water Management.) — Budapest.
- RAINCSÁKNÉ KOSÁRY Zs. (ed.) 1984: Budapest területének földtani, vízföldtani, építésalkalmassági térképei. (Geological, hydrogeological and construction suitability maps of Budapest.) — MÁFI publ., Budapest.
- SCHAREK P. (ed.) 1990: Magyarázó a Győr-Dél jelű térképlaphoz. A Kisalföld földtani térképsorozata 1:100 000. (Explanatory note to sheet Győr-South of the 1:100 000 scale geological map series of the Little Hungarian Plain.) — MÁFI publ., 30 p. Budapest.
- SCHAREK P. (ed.) 1991a: Magyarázó a Győr-Észak jelű térképlaphoz. A Kisalföld földtani térképsorozata 1:100 000. (Explanatory note to sheet Győr-North of the 1:100 000 scale geological map series of the Little Hungarian Plain.) — MÁFI publ., 31 p. Budapest.
- SCHAREK P. (ed.) 1991b: Magyarázó a Mosonmagyaróvár jelű térképlaphoz. A Kisalföld földtani térképsorozata 1:100 000. (Explanatory note to sheet Mosonmagyaróvár of the 1:100 000 scale geological map series of the Little Hungarian Plain.) — MÁFI publ., 35 p., Budapest.
- SCHAREK P. (ed.) 1993: A Kisalföld Földtani Térképsorozata, Kapuvár. (Geological map series of the Little Hungarian Plain, sheet Kapuvár.) — MÁFI Inventory n°: 4990., p. 39. Budapest.
- SCHAREK P. & TULLNER T. (ed.) 1992: A Kisalföld Földtani Térképsorozata Komárom–Tatabánya. (Geological map series of the Little Hungarian Plain, sheet Komárom–Tatabánya.) — Manuscript, Archives of the Little Plain Project, MÁFI, Budapest.
- SCHAREK P. & TULLNER T. (ed.) 1993: A Kisalföld Földtani Térképsorozata Sopron–Kőszeg. (Geological map series of the Little Hungarian Plain, sheet Sopron–Kőszeg.) — Manuscript, Archives of the Little Plain Project, MÁFI, Budapest.
- SCHMID, H. 1970: Regionaler Überblick der erdgeschichtlichen Entwicklung des Mattersburger Beckens. — *Burgenländische Heimatblätter* **32/2**, 49–55.
- SCHMID, H. 1972: Die geologischen Verhältnisse der weiteren Umgebung von Schützen am Leithagebirge (Burgenland). — *Wiss. Arbeiten Bgld.* **48**, 57–67.
- ~AJGALÍK, J. & MODLITBA, I. 1983: Sprae Podunajskej nížiny a ich vlastnosti. (Loesses of the Danube Lowland and their strength.) — Veda, vydavateľstvo Slovenskej akadémie vied. 204 p. Bratislava.
- VETTERS, H. 1910: Die Geologischen Verhältnisse der weiteren Umgebung Wiens und Erl. zur tektonischen Übersichtskarte des Wiener Beckens und seiner Randgebirge im Maastab 1:100 000. — Wien (Verl. österr. Lehrmittelanst.).
- WESSELY, G. 1968: Geomorphologische Karte der Hainburger Berge und angrenzender Gebiete 1:25 000. — Verh. Geol. B.-A., Wien.



# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Jahrbuch der Geologischen Bundesanstalt](#)

Jahr/Year: 2000

Band/Volume: [142](#)

Autor(en)/Author(s): Liscak Pavol, Scharek Peter, Tullner Tibor, Schäffer Gerhard, Kovacik Milos

Artikel/Article: [Engineering Geological Map 505-520](#)