

Linzer biol. Beitr.	42/2	1379-1384	19.12.2010
---------------------	------	-----------	------------

## A Contribution to the Ichneumon Wasps (Hymenoptera: Ichneumonidae) from Golestan National Park and Vicinity, Northeastern Iran

H. GHAHARI & R. JUSSILA

**A b s t r a c t :** The fauna of Ichneumonidae (Hymenoptera) from Golestan National Park and Vicinity (Northeastern Iran) is studied in this paper. In a total of 14 species from 14 genera (*Pristomerus*, *Temelucha*, *Aclastus*, *Gambus*, *Barytarbes*, *Diadromus*, *Rhexidermus*, *Megastylus*, *Phaeostacoenitus*, *Endromopoda*, *Rhyssa*, *Scambus*, *Strongylopis*, *Lathrolestes*), and 8 subfamilies (Cremastinae, Cryptinae, Ctenopelmatinae, Ichneumoninae, Orthocentrinae, Phuridinae, Pimplinae, Ctenopelmatinae) were collected from this large natural ecosystem.

**K e y w o r d s :** Ichneumonidae, Parasitoid, Fauna, New record, Golestan National Park, Iran.

### Introduction

Conservation of natural resources today is a respected and well-merited doctrine at both national and international levels. National parks possessing diverse examples of biotic communication and ecosystems, serving as scientific land use model and representing outstanding values of natural heritage, are one of the means of expressing this doctrine. The distinguishing characteristic of national parks is the non consumptive use of natural resources. They also have many scientific, cultural, educational and recreational inherent values. The Golestan National Park was the first area in Iran to be designated as a national park. It is located at 37,16° to 37,36° north latitude and 55,44° to 56,17° east longitude and has an area of about 91.000 hectares. The terrain is mountainous with altitude varying between 380 and 2819 meters. The park contains a rich diversity of flora and fauna, unique in many respects (HASSAN ZADEH et al. 1993).

Family Ichneumonidae (Hymenoptera) is one of the most species-rich families of all organisms with an estimated 60,000 species in the world, and included beneficial species which parasitize several pests (TOWNES 1969, GUPTA 1987). The Ichneumonidae, along with other groups of parasitic Hymenoptera, are purported to be no more species rich in the tropics than in the Northern Hemisphere temperate regions (OWEN & OWEN 1974, JANZEN 1981), although a number of hymenopteran families, for example the Chalcididae (HESPENHEIDE 1979) and Encyrtidae (NOYES 1989) exhibit an increase in species richness with a decrease in latitude. Other hymenopteran taxa such as sawflies (Symphyta), gall-forming Cynipidae, and bees (Apoidea) peak in species richness at mid- or high

latitudes (NOYES 1989, KOUKI et al. 1994). Considerable debate has centered on the apparent species richness anomaly exhibited by a number of hymenopteran parasitoid taxa in the tropics (e.g., MORRISON et al. 1978, GAULD 1991). Although the fauna of Iranian Ichneumonidae was studied rather well (KOLAROV & GHAHARI 2005, 2006, 2007, 2008, GHAHARI et al. 2010) but the fauna of these powerful parasitoids was not studied in Golestan National Park so far. In this paper we present the result of a faunistic survey in this region of northeastern Iran.

## Materials and Methods

The materials were collected by sweep netting and malaise traps from different regions of Golestan National Park and vicinity through four sampling trips in 2005-2007 by the first author and some of his colleagues and students. Through the faunistic survey, in addition to the collecting of adult Ichneumonidae, many immature insects (especially Lepidoptera and Hymenoptera) were collected and reared in optimum condition ( $26\pm2^{\circ}\text{C}$ ,  $65\pm5\%$  RH, 14:10 L:D) in incubator for emergence of inside parasitoids. Classification, nomenclature and distributional data of Ichneumonidae suggested by KASPARYAN (1981), YU & HORSTMANN (1997) and YU et al. (2005) have been followed.

## Results

Totally, fourteen species in fourteen genera and eight subfamilies were collected in this faunistic survey. The list of species is given below with host for some species.

### Subfamily Cremastinae

#### Genus *Pristomerus* CURTIS 1836

##### *Pristomerus horribilis* NAROLSKY 1987

Material: Golestan province: National Park, 1919 m, 1♂, September 2006.

#### Genus *Temelucha* FÖRSTER 1869

##### *Temelucha decorate* (GRAVENHORST 1829)

Material: Golestan province: Armodlu, 1550 m, 1♀, August 2007.

### Subfamily Cryptinae

#### Genus *Aclastus* FÖRSTER 1869

##### *Aclastus solutus* (THOMSON 1884)

Material: Golestan province: Sulgard, 1565 m, 1♀, September 2006.

**Genus *Gambus* FÖRSTER 1869**

***Gambus carnifex* (GRAVENHORST 1829)**

M a t e r i a l : Golestan province: Yaghtiklan, 1920 m, 1 ♀, August 2007.

**Subfamily Ctenopelmatinae**

**Genus *Barytarbes* FÖRSTER 1869**

***Barytarbes superbus* (SCHMIEDEKNECHT 1914)**

M a t e r i a l : Golestan province: Ghareh-Ghashli, 1825 m, 1 ♀, September 2006.

**Subfamily Ichneumoninae**

**Genus *Diadromus* WESMAEL 1845**

***Diadromus quadriguttatus* GRAVENHORST 1889**

M a t e r i a l : Golestan province: Dasht-e-Mirzabaylu, 1575 m, 2 ♀ ♀, August 2007, reared from *Yponomeuta* sp. (Lepidoptera: Yponomeutidae).

**Genus *Rhexidermus* FÖRSTER 1869**

***Rhexidermus truncator* FÖRSTER 1889**

M a t e r i a l : Golestan province: National Park, 2157 m, 2 ♀ ♀, 1 ♂, June 2006, reared from Lasiocampidae (Lepidoptera). Golestan province: Cheshmeh-khan, 1586 m, 1 ♀, June 2006, larval parasitoid of *Laspeyresiya* sp. (Lepidoptera: Tortricidae).

**Subfamily Orthocentrinae**

**Genus *Megastylus* SCHIODTE 1839**

***Megastylus flavopictus* (GRAVENHORST 1829)**

M a t e r i a l : Golestan province: National Park, 1967 m, 3 ♀ ♀, October 2005.

**Subfamily Phuridinae**

**Genus *Phaestacoenitus* SMITS VAN BURGST 1913**

***Phaestacoenitus niger nitidus* KASPARYAN 1983**

M a t e r i a l : Golestan province: Ghareh-Ghashli, 1825 m, 3 ♀ ♀, September 2006.

### Subfamily P i m p l i n a e

#### Genus *Endromopoda* HELLÉN 1939

##### *Endromopoda detrita* (HOLMGREN 1860)

M a t e r i a l : Golestan province: Dasht-e-Mirzabaylu, 1575 m, 1♀, August 2007.

#### Genus *Rhyssa* GRAVENHORST 1829

##### *Rhyssa persuasoria* (LINNAEUS 1758)

M a t e r i a l : Golestan province: Ghareh-Ghashli, 1825 m, 2♀♀, September 2006.

#### Genus *Scambus* HARTIG 1838

##### *Scambus (Endromopoda) detritus* (HOLMGREN 1860)

M a t e r i a l : Golestan province: Ghoosh-Cheshmeh, 1612 m, 1♀, September 2006.

#### Genus *Strongylopsis* BRAUNS 1896

##### *Strongylopsis belua* KUZIN 1950

M a t e r i a l : Golestan province: Yaghtiklan, 1920 m, 3♂♂, August 2007.

### Subfamily C t e n o p e l m a t i n a e

#### Genus *Lathrolestes* FÖRSTER 1869

##### *Lathrolestes ensator* (BRAUNS)

M a t e r i a l : Golestan province: National Park, 1881 m, 4♀♀, June 2006, larval parasitoid of *Hoplocampa testudinea* (KLUG) (Hymenoptera: Tenthredinidae).

### Discussion

The result of the present research indicates that the Golestan National Park included a diverse and interesting fauna of Ichneumonidae. Since there is a diverse flora especially several forest trees (e.g. the dominant plant species *Acanthophyllum* spp., *Acer* spp., *Artemesia* spp., *Berberis* spp., *Carpinus* spp., *Crateagus* pp., *Festuca* spp., *Quercus* spp., *Rhamnus* spp., *Rosa* spp., *Rubus* spp.) in this natural park (HASSAN ZADEH et al. 1993), therefore several insect pests of different orders (especially Lepidoptera and Coleoptera) are injurious to these plants. As we mentioned in introduction, Golestan National Park is a vast region incorporating various geographical regions and diverse ecosystems; therefore we expect that a large number of ichneumonid species (new records and probably new species) remain to be discovered. To find new species and distributional records, more studies should be conducted in this large natural ecosystem. We suggest strongly that other Iranian researchers especially the taxonomists start to faunistic surveys in different regions of Iran for discovering several unknown data about the fauna of Ichneumonidae and other parasitoids and also their hosts.

### Acknowledgments

The authors are indebted to Dr. J. Kolarov of Bulgaria, Dr. D.R. Kasparyan of Russian Academy of Sciences, Dr. M. Schwarz of Austria and Dr. T. Finlayson of Canada for invaluable helps in progress of the project. We are also thanks to Dr. H. Sakenin and Z. Karimian for invaluable helps and loaning some specimens. The research was supported by Islamic Azad University and Zoological Museum of Turku University.

### Zusammenfassung

Vorliegende Arbeit behandelt die Ichneumonidenfauna des Golestan National Parks und seiner Umgebung im Nordosten des Irans. Insgesamt gelang der Nachweis von 14 Arten aus 14 Gattungen (*Pristomerus*, *Temelucha*, *Aclastus*, *Gambus*, *Barytarbes*, *Diadromus*, *Rhexidermus*, *Megastylus*, *Phaestacoenitus*, *Endromopoda*, *Rhyssa*, *Scambus*, *Strongylopis*, *Lathrolestes*) aus den folgenden 8 Unterfamilien: Cremastinae, Cryptinae, Ctenopelmatinae, Ichneumoninae, Orthocentrinae, Phuridinae, Pimplinae und Ctenopelmatinae.

### References

- GAULD I.D. (1991): The Ichneumonidae of Costa Rica, 1. Introductions, keys to subfamilies, and keys to the species of the lower pimpliform subfamilies Rhyssinae, Pimplinae, Poemeniinae, Acaenitinae and Cyllocerinae. — Mem. Amer. Entomol. Inst. **47**: 1-577.
- GHAHARI H., JUSSILA R., KOLAROV J. & J. SEDIVY (2010): A contribution to the ichneumon wasps (Hymenoptera: Ichneumonidae) from the forests of northern Iran. — Munis Entomology & Zoology **5**: 85-89.
- GUPTA V.K. (1987): The Ichneumonidae of the Indo-Australian area (Hymenoptera). — Mem. Amer. Entomol. Institute **41**: 1-1210.
- HASSAN ZADEH B., ZEHZAD B., FARHANG B., MAJNOUNIAN H. & H. GOSHTASB (1993): Golestan National Park. — Department of Environment, Fardin Publication, 203 pp.
- HESPENHEIDE H.A. (1979): Are there fewer parasitoids in the tropics? — Amer. Natur. **113**: 766-769.
- JANZEN D.H. (1981): The peak in North American ichneumonid richness lies between 38° and 42°N. — Ecology **62**: 532-537.
- KASPARYAN D.R. (1981): Opredelitel Nasekomich Europeiskoy Casti U.S.S.R. III. Part. Pereponchato-krylye 3. Opredelitel Fauny SSSR. Nauka, Moscow-Leningrad, pp. 1-688. [In Russian].
- KOLAROV J. & H. GHAHARI (2005): A catalogue of Ichneumonidae (Hymenoptera) from Iran. — Linzer Biologische Beiträge **37** (1): 503-532.
- KOLAROV J. & H. GHAHARI (2006): A study of the Iranian Ichneumonidae (Hymenoptera): I. Pimplinae and Tryphoninae. — Zoology in the Middle East **38**: 69-72.
- KOLAROV J. & H. GHAHARI (2007): A study of the Iranian Ichneumonidae (Hymenoptera): II. Brachycyrtinae and Cryptinae. — Zoology in the Middle East **42**: 79-82.
- KOLAROV J. & H. GHAHARI (2008): A study of the Iranian Ichneumonidae (Hymenoptera). III. Ichneumoninae. — Acta Entomologica Serbica **13**: 61-76.
- KOUKI J., NIEMELÄ P. & M. VIITASAARI (1994): Reversed latitudinal gradient in species richness of sawflies (Hymenoptera, Symphyta). — Ann. Zool. Fenn. **31**: 83-88.
- MORRISON G., AUERBACH M. & E.D. MCCOY (1978): Anomalous diversity of tropical parasitoids: a general phenomenon? — Amer. Natur. **114**: 303-307.

- NOYES J.S. (1989): The diversity of Hymenoptera in the tropics with special reference to Parasitica in Sulawesi. — *Ecol. Entomol.* **14**: 197-207.
- OWEN D.F. & J. OWEN (1974): Species diversity in temperate and tropical Ichneumonidae. — *Nature* **249**: 583-584.
- TOWNES H.K. (1969): The genera of Ichneumonidae, part 1. — *Mem. Amer. Entomol. Institute* **11**: 1-300.
- YU D.S. & K. HORSTMANN (1997): A catalogue of world Ichneumonidae (Hymenoptera). — *Memoirs of the American Entomological Institute* **58** (1-2): 1558 pp.
- YU D.S., ACHTERBERG C. VAN & K. HORSTMANN (2005): Biological and taxonomical information: Ichneumonoidea 2004. — Taxapad Interactive Catalogue, Vancouver.

Author's addresses:

Hassan GHAHARI  
Department of Entomology  
Islamic Azad University  
Science & Research Branch  
Tehran, Iran  
E-mail: hghahari@yahoo.com

Reijo JUSSILA  
Zoological Museum, Section of Biodiversity and Environmental Sciences, Department of Biology,  
FI-20014 University of Turku Finland  
E-mail: reijo.jussila@utu.fi

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2010

Band/Volume: [0042\\_2](#)

Autor(en)/Author(s): Ghahari Hassan, Jussila Reijo

Artikel/Article: [A Contribution to the Ichneumon Wasps \(Hymenoptera: Ichneumonidae\) from Golestan National Park and Vicinity, Northeastern Iran 1379-1384](#)