

Linzer biol. Beitr.	46/2	1517-1524	19.12.2014
---------------------	------	-----------	------------

Ichneumonidae (Hymenoptera) from Anatolia. I.

J. KOLAROV, İ. ÇORUH & S. ÇORUH

A b s t r a c t : Faunistic data of 12 Ctenopelmatinae species from Turkey are reported. *Euryproctus arbustorum* HOLMGREN, *Euryproctus luteicornis* (GRAVENHORST), *Hadrodactylus larvatus* KRIECHBAUMER, *Lathrolestes (Lathrolestes) buccinator* (HOLMGREN) and *Phobetes atomator* (MÜLLER) marked in the text by asterisk are new records for Turkish fauna. A zoogeographic characterization for each species is proposed.

K e y w o r d s : Ichneumonidae, Ctenopelmatinae, fauna, Turkey, new records, zoogeography.

Introduction

Ichneumonidae is the biggest hymenopteran family with 51 generally recognized subfamilies, 1579 genera and 24281 described species (YU et al. 2012). TOWNES (1969), estimated that there could be about 60 000 Ichneumonidae species in the world, but because of the poor knowledge of the tropical faunas the present investigators estimate that the size of the family could be higher than 100 000 (GAULD 1997). The known number of Ichneumonidae species increases rapidly in the world. Ichneumonids are parasitoids of immature holometabolous insects from orders such as Coleoptera, Diptera, Hymenoptera, Lepidoptera, Raphidiopelta, Trichoptera and also non-insect (Chelicera: Araneae) (TOWNES 1969, 1970; WAHL 1993).

The Ichneumonidae fauna of Turkey is not well studied. In the catalogue of KOLAROV (1995) 383 species were listed from Turkey. Later many authors (KOLAROV et al. 1997a,b; KOLAROV et al. 1999; KOLAROV et al. 2002; ÇORUH et al. 2002; KOLAROV & GÜRBÜZ 2004; GÜRBÜZ 2005; YURTCAN & BEYARSLAN 2005, 2006; ÇORUH & ÖZBEK 2008; RIEDEL et al. 2010; KOLAROV & ÇALMAŞUR 2011; etc.) have made contributions to the Turkish fauna. ÇORUH & GÜCLÜ (2013) reported 983 species in 282 genera for Turkey Ichneumonidae fauna.

Not so far the Ctenopelmatinae has been mentioned as Scolobatinae (TOWNES 1969; WAHL 1993). The subfamily is easily recognized by the following characters: Body small to large (fore wing 2.9-22 mm long). Clypeus fairly flat, usually wide and short, usually separated from face by groove, the apical margin often blunt or rounded; mandible long and weakly narrowed; apex of protibia with tooth on dorsal margin; sternaulus of mesopleuron absent or short; metasomal segment 1 slender to very stout, with or without glymma, and with spiracle before or at the middle; metasoma usually

cylindrical or depressed, sometimes compressed; ovipositor barely extending beyond metasomal apex, the dorsal subapical notch present except when ovipositor is needle-like.

Ctenopelmatinae are koinobiont endoparasitoids of Symphyta and, rarely, Lepidoptera; oviposition is into the egg or larva, with emergence after the host cocoon is spun. Most species have a Holarctic distribution (WAHL 1993).

KOLAROV (1995) reported 26 species in 17 genera for Turkey Ctenopelmatinae fauna. With the below mentioned contributions (KOLAROV et al. 1997a; ÖZBEK et al. 2000; ÇORUH et al. 2002; GÜRBÜZ & AKSOYLAR 2005; RESHCHIKOV 2011; ÇORUH & ÖZBEK 2011; GÜRBÜZ et al. 2011) the numbers of Ctenopelmatinae fauna of Turkey reached to 36 species.

This study is based upon material of family Ichneumonidae collected from different localities of Anatolia in Turkey in 2013.

Material and methods

The collecting of insects was conducted in two different regions - the Black Sea Region (Rize, Trabzon and Ordu) and the Eastern Anatolia Region (Erzurum) (Fig. 1). "Rize, Trabzon and Ordu" are situated in Middle and Eastern Black Sea Region of Turkey. These areas have rainy every season. The highest rainfall in autumn, at least rain fall in spring these areas. While annual rainfall is 1000-1500 mm in Eastern Black Sea Region (Rize and Trabzon), annual rain fall is 1000-1200 mm in Middle Black Sea Region (Ordu). The average of snowy days is 18 days in the region and the average July temperature is 21-23°C. The natural vegetation is forest. The vegetation is broad-leaved forest in the lowlands. The vegetation is changed as altitude increases and seen mixed deciduous forests. Coniferous forests and alpine meadows seen very high places (2100-2300 m) (http 1). In this region, predominant vegetations are as follows:

Heracleum platytaenium BOISS., *Conyza canadensis* (L.) CRONQUIST., *Erigeron acer* L., *Erigeron annuus* (L.) PERS., *Equisetum palustre* L., *Geranium asphodeloides* BURM. FIL., *Geranium ibericum* CAV., *Daucus carota* L., *Geranium sylvaticum* L., *Pteridium aquilinum* (L.) KUHN, *Prunella vulgaris* L., *Salvia forskahlei* L., *Salvia verticillata* L., *Linum hypericifolium* SALISB., *Epilobium parviflorum* SCHREBER, *Plantago media* L., *Alopecurus myosuroides* HUDSON, *Cynodon dactylon* (L.) PERS., *Lolium temulentum* L., *Lolium perenne* L., *Sorghum halepense* (L.) PERS., *Digitaria sanguinalis* (L.) SCOP., *Paspalum dilatatum* POIRÉT, *Paspalum paspalodes* (MICHX.) SCRIBNER, *Setaria glauca* (L.) P. BEAUV., *Poa annua* L., *Poa trivialis* L., *Festuca pratensis* HUDSON, *Phleum phyleoides* (L.) KARSTEN, *Polygonum persicaria* L., *Anagallis arvensis* L., *Delphinium formosum* BOISS. & HUET, *Fragaria vesca* L., *Rubus discolor* WEIHE & NEES, *Rubus hirtus* WALDST. et KIT., *Galium verum* L., *Rhinanthus angustifolius* C.C. GMELIN, *Pedicularis comosa* L., *Corylus avellana* L., *Veronica gentianoides* VAHL. and *Urtica dioica* L.

Three species (*Euryproctus arbustorum*, *Euryproctus luteicornis* and *Phobetes atomator*) which are new records for the Turkish fauna were collected from Ikizdere (Rize). Ikizdere is situated in Eastern Black Sea Region of Turkey (40°42' N, 40° 36' E), covers an area from sea level to 570 m. a. s. l. The climate is typical of the Black Sea region.

Rain is falls all the time. There is snow on the high peaks and lots of glacial formation. This is hilly countryside inland from the Black Sea, forest-covered steep mountainside separated by narrow valleys, with areas of high pasture. There is broad-leaf forest at lower elevations and pine higher up. Ikizdere has Anzer pasture and Ovit mountain (http 2).

"Erzurum" is situated in Eastern Anatolia Region of Turkey. The altitude of this region is higher than Black Sea region. The majority of this region has high altitude. Most plateaus are around 2000 m from sea level, and the mountainous regions beyond the plateaus are 3000 m and higher. Depression plains are located between the mountains and plateaus. The southern mountain ranges of Erzurum city, Palandoken Mountains, with the altitudes of 2200-3176 m. The topographic and climatic structures province has the opportunity of host rich and diverse fauna and flora (YILDIRIM & STRUMIA 2000). Climate in this region is terrestrial. That is, winters are long and hard; summer is very short and warm. Place in Kars and Erzurum is the number of days with snow is about 90 days. The average July temperature is 17-19°C. Annual rainfall is 500-600 mm. The natural vegetation is usually steppe looks like. While forests consist of yellow pines and coniferous trees in the northern part of the region, in other sections, consist of oak. Where the high-end segments of the forest, starts alpine meadows are covered with grassland (http 3). In this region, predominant vegetations are as follows:

Acer monspessulanum L., *Achillea biebersteinii* AFAN., *A. millefolium* L., *Antemis cretica* L., *A. tinctoria* L., *Arabis caucasica* WILLD., *Astragalus christianus* L., *Carum carvi* L., *Cirsium arvense* (L.) SCOP., *Coronilla orientalis* MILL., *Cotinus coggyria* SCOP., *Daucus carota* L., *Equisetum ramosissimum* DESF., *Ephedra major* HOST, *Eryngium billardieri* DELAR, *Euphorbia virgata* WALDST. & KIT., *Ferula communis* L., *F. orientalis* L., *Galium incanum* SM., *Gypsophila bicolor* (FREYN & SINT.), *Hypericum hyssopifolium* CHAIX, *H. scabrum* L., *Juniperus communis* L., *Linum mucronatum* BERTOL., *Papaver orientale* L., *Pimpinella corymbosa* BOISS., *P. tragium* VILL., *Ranunculus cuneatus* BOISS., *Rhus coriaria* L., *Seseli libanotis* (L.) W. KOCH, *Trifolium ambiguum* M. BIEB., *T. hybridum* L., *Sisymbrium elatum* K. KOCH, *Tanacetum punctatum* (BOISS. & NOE), *Veronica orientalis* MILLER, *Ziziphora clinopodioides* LAM. and *Zosima absinthifolia* (WENT.).

Two species (*Hadrodactylus larvatus* and *Lathrolestes (Lathrolestes) buccinator*) which are new records for the Turkish fauna were collected from Gelinkaya (Erzurum). Gelinkaya which is situated in Eastern Anatolia Region of Turkey (40°02' N, 40°91' E), covers an area from sea level to 1790 m.a.s.l. Vegetation is steppe and the topographic structure is mountainous. The climate is terrestrial. However, Black Sea climatic features are also seen. That is, winters are milder due to the geographical location (http 3).

Materials were collected by sweeping on flowering plants in the different localities Rize, Trabzon and Ordu (Black Sea Region) and Erzurum (Eastern Anatolia Region) in Turkey, during 2013. Collected samples were transferred into a handmade aspirator and were killed with ethyl acetate. Conventional standard method (ÇORUH & ÖZBEK, 2008) was used for preparation of the samples. Material is preserved in Collection of Plovdiv University (Bulgaria). Plant specimens were collected by hand and were pressed and they were deposited at the Herbarium of Plant Protection Department (Erzurum). In the present paper unpubl. data for 12 species from subfamily Ctenopelmatinae from Anatolia are listed. New records of species are marked by an asterisk (*). General distributions of the species were taken from YU et al. (2012).

Results

List of the species

In total 12 species in 9 genera of Ctenopelmatinae were determined. The species list is given below.

****Euryproctus arbustorum* HOLMGREN, 1857**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 2♂♂.

D i s t r i b u t i o n : Norway, Sweden, Finland, Poland, Austria, Ukraine, Belarus and Russia – Pskov oblast.

****Euryproctus luteicornis* (GRAVENHORST, 1829)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♀.

D i s t r i b u t i o n : Sweden, Finland, Ireland, United Kingdom, France, Germany, Switzerland, Poland, late Czechoslovakia, Austria, Hungary, Lithuania and Georgia.

***Euryproctus nemoralis* (GEOFFROY, 1785)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♂.

D i s t r i b u t i o n : Europe, Azerbaijan, Turkey and Russia (European part and Siberia).

****Hadrodactylus larvatus* KRIECHBAUMER, 1891**

M a t e r i a l e x a m i n e d : Erzurum, Gelinkaya, 19.06.2013, 1♀.

D i s t r i b u t i o n : Spain, Finland, Luxemburg, Italy, Germany, Austria, late Czechoslovakia, Hungary, Poland, Lithuania, Ukraine, Belarus, Azerbaijan, European Russia and Siberia.

****Lathrolestes (Lathrolestes) buccinator* (HOLMGREN, 1857)**

M a t e r i a l e x a m i n e d : Erzurum, Gelinkaya, 19.06.2013, 1♀.

D i s t r i b u t i o n : Europe, Azerbaijan, Armenia, Afghanistan, Siberia and Pacific coast of Russia.

***Mesoleptidea cingulata* (GRAVENHORST, 1829)**

M a t e r i a l e x a m i n e d : Trabzon, Çaykara, Uzungöl, 21.06.2013, 1♂.

D i s t r i b u t i o n : Europe, Turkey and China – Liaoning.

***Mesoleptidea* sp.**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♀.

***Perilissus spilonotus* (STEPHENS, 1835)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♂.

D i s t r i b u t i o n : Europe and Turkey.

***Perilissus variator* (MÜLLER, 1776)**

M a t e r i a l e x a m i n e d : Erzurum, Gelinkaya, 19.06.2013, 1♂.

D i s t r i b u t i o n : Palaearctic region.

****Phobetes atomator* (MÜLLER, 1776)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♀.

D i s t r i b u t i o n : Europe and Siberia - Yakutia.

***Phobetes leptocerus* (GRAVENHORST, 1820)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♂.

D i s t r i b u t i o n : Europe, Turkey and Siberia – Chita oblast of Russia.

***Rhorus* sp.**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 17♂♂.

***Scolobates auriculatus* (FABRICIUS, 1804)**

M a t e r i a l e x a m i n e d : Ordu, Gülyali, 22.06.2013, 1♂.

D i s t r i b u t i o n : Holarctic and Oriental region.

***Xenoschesis (Polycinetis) ustulata* (DESVIGNES, 1856)**

M a t e r i a l e x a m i n e d : Rize, Ikizdere, 24.06.2013, 1♀.

D i s t r i b u t i o n : Holarctic region.

Zoogeographic characterization

The zoogeographic characterisation follows mainly the chorotype classification of the Near East fauna, proposed by TAGLIANTI et al. (1999). After investigation of the recent geographic distribution of the species listed above, they can be divided into the following groups:

- (1) Species with range in Holarctic and Oriental region: *Scolobates auriculatus*.
- (2) Holarctic chorotypes: *Xenoschesis (Polycinetis) ustulata*.
- (3) Palaearctic ranges: *Mesoleptidea cingulata*, *Lathrolestes* (*Lathrolestes buccinators*) and *Perilissus variator*.
- (4) Sibero-European chorotypes: *Euryproctus nemoralis*, *Hadrodactylus larvatus*, *Phobetes atomator* and *Phobetes leptocerus*.

- (5) European chorotypes: *Euryproctus arbustorum*, *Euryproctus luteicornis* and *Perilissus spilonotus*.

Acknowledgements

The preparation of this article was supported by the Project Nr: BAP- 2012/234 and we are very thankful for it. We are indebted to Halil Çoruh (Erzurum) for collecting a few specimens.

Zusammenfassung

Vorliegende Arbeit berichtet über faunistische Daten von 12 Ctenopelmatinae-Arten (Hymenoptera, Ichneumonidae) aus der Türkei. Die Spezies *Euryproctus arbustorum* HOLMGREN, *Euryproctus luteicornis* (GRAVENHORST), *Hadrodactylus larvatus* KRIECHBAUMER, *Lathrolestes (Lathrolestes) buccinator* (HOLMGREN) und *Phobetes atomator* (MÜLLER), im Text mit * markiert, sind Erstnachweise für die Türkei. Jede angeführte Art wurde einer zoogeografischen Bewertung unterzogen.

References

- ÇORUH S. & C. GÜCLÜ (2013): Contribution to the Ichneumonidae (Hymenoptera) fauna of Turkey with some ecological notes. — Journal of the Entomological Research Society (in press).
- ÇORUH S., ÖZBEK H. & J. KOLAROV (2002): New and rare taxa of Ichneumonidae (Hymenoptera) from Turkey. — Journal of the Entomological Research Society **4** (1): 1-4.
- ÇORUH S., ÖZBEK H. & M. RIEDEL (2001): An additional contribution to the Ichneumoninae (Hymenoptera: Ichneumonidae) fauna of Turkey. — Turkish Journal of Entomology **35** (4): 603-613.
- DAVIS P.H. (1965-1988): Flora of Turkey and the East Aegean Island. — At the University Press, Edinburg, Vol. **1-10**.
- GÜRBÜZ M.F. (2005): A survey of the Ichneumonidae (Hymenoptera) of Isparta in Turkey. — Linzer biologische Beiträge **37** (2): 1809-1817.
- GÜRBÜZ M.F. & M.Y. AKSOYLAR (2005): New records of Ichneumonidae (Hymenoptera) species from Turkey. — Phytoparasitica **33** (2): 121-122.
- GÜRBÜZ M.F., KOLAROV J., ÖZDAN A. & M.A. TABUR (2011): Ichneumonidae (Hymenoptera) fauna of natural protection areas in East Mediterranean Region of Turkey. Part 1. — Journal of the Entomological Research Society **13** (1): 23-39.
- <http://tr.wikipedia.org> (accessed 04-Dec-2013).
- <http://tr.wikipedia.org/wiki/%C4%B0kizdere> (20-Sep-2013).
- <http://www.msxlabs.org/forum/soru-cevap/343506-dogu-anadolu-bolgesinin-bitki-ortusu-nedir.html#ixzz2mPw6Twkb> (accessed 04-Dec-2013).
- KOLAROV J. (1995): A catalogue of the Turkish Ichneumonidae (Hymenoptera). — Entomofauna **16** (7): 137-188.
- KOLAROV J. & M.F. GÜRBÜZ (2004): A study of the Turkish Ichneumonidae (Hymenoptera) I. Pimplinae. — Linzer biologische Beiträge **36** (2): 841-845.
- KOLAROV J. & Ö. ÇALMAŞUR (2011): A study of Ichneumonidae (Hymenoptera) from North Eastern Turkey. — Linzer biologische Beiträge **43** (1): 777-782.

- KOLAROV J., YURTCAN M. & A. BEYARSLAN (1997a): New and rare Ichneumonidae (Hymenoptera) from Turkey. I. Pimplinae, Tryphoninae, Phygaedeontinae, Banchinae and Ctenopelmatinae. — *Acta Entomologica Bulgarica* **3** (4): 9-12.
- KOLAROV J., YURTCAN M. & A. BEYARSLAN (1997b): Ichneumonidae (Hymenoptera) from the Gökçeada and Bozcaada Island, Turkey. — *Acta Entomologica Bulgarica* **3** (4): 13-16.
- KOLAROV J., ÖZBEK H. & E. YILDIRIM (1999): New distributional data of the Turkish Ichneumonidae (Hymenoptera). I. Pimplinae and Tryphoninae. — *Journal of the Entomological Research Society* **1** (2): 9-15.
- KOLAROV J., YURTCAN M. & A. BEYARSLAN (2002): Ichneumonidae species of the Turkish Aegean region. — In: MELİKA G. & C. THUROczy (eds), Parasitic wasps: evolution, systematics, biodiversity and biological control. International Symposium: "Parasitic Hymenoptera: Taxonomy and Biological Control" (14-17 May 2001, Koszeg, Hungary): 299-305. Agroinform Kiado & Nyomda Kft., Budapest.
- ÖZBEK H., PEKEL S. & J. KOLAROV (2000): New distributional data of the Turkish Ichneumonidae (Hymenoptera) II. Ctenopelmatinae and Campopleginae. — *Journal of the Entomological Research Society* **2** (1): 17-24.
- RESCHCHIKOV A. (2011): *Lathrolestes* (Hymenoptera, Ichneumonidae) from Turkey with description of three new species and new synonymy. — *Journal of the Entomological Research Society* **13** (1): 83-89.
- RIEDEL M., ÇORUH S. & H. ÖZBEK (2010): Contribution to the Ichneumoninae (Hymenoptera, Ichneumonidae) fauna of Turkey, with description of three new species. — *Turkish Journal of Entomology* **34** (2): 133-156.
- TOWNES H. (1969): The Genera of Ichneumonidae, Part I. — *Memoirs of the American Entomological Institute* **11**: 1-300.
- TOWNES H. (1970): The genera of Ichneumonidae, Part. III. — *Memoirs of the American Entomological Institute* **13**: 1-307.
- TAGLIANTI, V., AUDISIO A., MAURICIO P.A., BOLOGNA B., CARPANETO M.A., BIASE M.G., FATTORINI A.De., PIATELLA S., SINDACO E., VENCHI R. & A. ZAPPAROLI (1999): A proposal for a chorotype classification of the Near East fauna, in framework of the Western Palaearctic region. — *Biogeographia* **20**: 31-59.
- WAHL D.B. (1993): Family Ichneumonidae. — In: GOULET H. & J. T. HUBER (eds), Hymenoptera of the World: an identification guide to families. Research Branch. Monograph No. 1894c. Agriculture Canada Publication. Ottawa, Ontario, Canada. 668 pp.
- YURTCAN M. & A. BEYARSLAN (2005): Polysphinctini and Pimplini (Hymenoptera: Ichneumonidae: Pimplinae) from the Thrace region of Turkey. — *Fragmata Faunistica* **48** (1): 63-72.
- YURTCAN M. & A. BEYARSLAN (200): Six new Ichneumonidae species from Turkey with special reference to the rare species *Zabraclypus tenuiabdominalis* (UCHIDA, 1941) (Hymenoptera: Ichneumonidae). — *Entomological News* **117** (5): 540-544.
- YU D., Van ACHTERBERG C. & K. HORSTMANN (2012): Taxapad 2012, Ichneumonoidea 2011.— Database on flash-drive. www.taxapad.com, Ottawa, Ontario, Canada.
- YILDIRIM E. & F. STRUMIA (2000): Contribution to the knowledge of Chrysidae fauna of Turkey. Part 1: Cleptinae (Hymenoptera, Chrysidae). — *Frustula entomologica* **23**: 161-166.

Authors' addresses:

Dr. Janko KOLAROV
University of Plovdiv, Faculty of Pedagogy
24 Tsar Assen Str.
4000 Plovdiv, Bulgaria
E-mail: jkolarov@uni-plovdiv.bg

Dr. İrfan ÇORUH and Dr. Saliha ÇORUH
Atatürk University, Faculty of Agriculture
Department of Plant Protection
25240 Erzurum, Turkey
E-mail: spekel@atauni.edu.tr



Fig. 1: Map of the region investigated.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2014

Band/Volume: [0046_2](#)

Autor(en)/Author(s): Kolarov Janko Angelov, Coruh Irfan, Coruh Saliha

Artikel/Article: [Ichneumonidae \(Hymenoptera\) from Anatolia. I. 1517-1524](#)