

Linzer biol. Beitr.	46/2	1599-1606	19.12.2014
---------------------	------	-----------	------------

Mites of the family Laelapidae (Acari: Mesostigmata) associated with olive orchards in Guilan Province Iran

M. MAHJOORI, J. HAJIZADEH & M.R. ABBASII MOZHDEHI

A b s t r a c t : A faunistic study of family Laelapidae (Acari: Mesostigmata) associated with olive orchards in Guilan province, Northern Iran was carried out during 2012-2013. Soil and plant remains samples were taken from olive orchards. The mites were extracted by Berlese funnel and cleared in Nesbitt's fluids; then were mounted in Hoyer's medium on microscopic slides. During this study 18 species belonging to 9 genera were collected and identified. All species are new for olive orchards mite's fauna of Iran.

K e y w o r d s : Fauna, Mesostigmata, Laelapidae, Olive, Iran.

Introduction

Olive, *Olea europaea* L. is one the most important fruit in the world, including Iran. Archaeological findings revealed that olive cultivation in Iran dates back to 2000 years ago (SADEGHI 1992). At present olive cultivars are cultivated mainly in the North of Iran, which is characterized by Mediterranean climatic condition. In the last ten years, olive plantation has grown in Iran and currently, 95.000 hectares of olive orchards produce about 6.500 tons of olive oil annually (SAMAE et al. 2003; HOSSEINI-MAZINANI et al. 2004; NOORMOHAMMADI et al. 2007; OMRANI-SABBAGHI et al. 2007; SHEIDAI et al. 2007). Guilan province has 13118 hectares olive orchards with 28125 tons annual production (MINISTRY of Jihad AGRICULTURE of IRAN 2013). Guilan Province is one of the 31 provinces of Iran. It covers an area of 14.042 km² and located in the North of Iran between the latitude 37°16'38.64"N and the longitude 49°35'20.4"E. It lies along the Caspian Sea and has a humid subtropical climate with a large margin the heaviest rainfall in Iran: reaching as high as 1,900 millimeters (75 in).

The mite family Laelapidae is ecologically diverse, including obligate and facultative parasites of vertebrates, insect paraphages, and free-living predators that inhabit soil-litter habitats and the nests of vertebrates and arthropods (EVANS & TILL 1966; STRONG & HALLIDAY 1994; LINDQUIST et al. 2009). Members of the subfamily Hypoaspidinae (Hypoaspididae of KARG 1993) often are collected in litter or soil substrates (the genera *Pseudoparasitus*, *Ololaelaps* and those often included under *Hypoaspis*, such as *Gaeolaelaps*, *Cosmolaelaps*, and *Stratiolaelaps*, are examples), while species of these and other genera are routinely or occasionally encountered in the nests of mammals or arthropods or directly associated with insects (EVANS & TILL 1966, 1979; KARG 1978,

1993). Some species are ectoparasites of adult bees or their brood, like *Tropilaelaps*, and some of them are also known to feed facultatively on nesting host animals by chewing or punching a hole in the skin (RADOVSKY 1967; LINDQUIST et al. 2009). Some species have established close phoretic relationships with other arthropods and less commonly, with vertebrates (KRANTZ 1998; LINDQUIST et al. 2009). Those are common predators in different kinds of habitats such as: freshwater and marine habitats, the littoral-intertidal-estuarine milieu, organic matter (COSTA 1962, 1968, 1974; LINDQUIST et al. 2009).

The fauna of mites associated with olive groves in Iran is very poorly known. Due to the predatory importance of laelapid mites and their important role in natural control of injurious edaphic pests such as arthropods and nematodes, a faunistic study was carried out for identification of laelapid mites associated with olive orchards in Guilan Province Iran. The present paper is part of a master thesis project of first author on mesostigmatic mites associated with olive orchards in Guilan Province Iran.

Materials and methods

A faunal study on family Laelapidae associated with olive orchards was carried out in Guilan Province, Northern Iran during 2012-2013. Soil and Plant remains samples were collected from olive orchards of Guilan Province. Mites were extracted from collected samples by placing them on Berlese funnel. Specimens preserved in 75% ethanol, cleared in Nesbitt's fluids and mounted on microscopic slides using Hoyer's medium. The slides were placed in at 45°C for two weeks. Specimens were identified by the relevant taxonomic keys and papers. The setations pattern is widely used as a taxonomic criterion in the Mesostigmata, and the system followed in the present work is that of LINDQUIST & EVANS (1965) and LINDQUIST (1994) for dorsal and ventral setations respectively. The voucher material which comprises slide mounted specimens is deposited in the Department of Plant Protection at University of Guilan, Rasht, Iran.

Results

In the current study 18 species belonging to 8 genera from family Laelapidae were collected and identified, in association with olive orchards in Guilan province Iran. All species are new for olive orchards mite fauna in Iran. The list of identified species is as follow.

L a e l a p i d a e BERLESE, 1892

***Haemolaelaps* BERLESE, 1910**

***Haemolaelaps shealsi* (COSTA, 1968)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, October 2012; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, July 2013; Nesfi, 455m, 36°50'44.85''N, 49°30'04.60''E, August 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran, Israel (COSTA 1968; HADDAD IRANI-NEJAD et al. 2003).

***Gaeolaelaps* EVANS & TILL, 1966**

***Gaeolaelaps aculeifer* (CANESTRINI, 1884)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, July 2013; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, July 2013; August 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Canada, England, Germany, Iran, Italy, Netherlands, North American and Sweden, (HUGHES 1976; BALOOCH SHAHRYARI et al. 2012).

***Gaeolaelaps kargi* COSTA, 1968**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, October 2012; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, July 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : America, Bashkiria (in the floodlands of the Belaya river), Dagestan, Europe, Georgia, Iran, Israel, Lithuania, Moldavia, Ukraine and USSR (COSTA 1968; GILYAROV et al. 1977; KARG 1982; BALOOCH SHAHRYARI et al. 2012).

***Gaeolaelaps nolli* KARG, 1962**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, October 2012. Collected by M. Mahjoori.

D i s t r i b u t i o n : Asia, Europe and Iran (KARG 1962; 1982; BALOOCH SHAHRYARI et al. 2012).

***Gaeolaelaps queenslandica* (WOMERSLEY, 1956)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, October 2012 & July 2013 & August 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : America, Australia, Iran (WORMERSLEY 1954; TENORIO 1982; BALOOCH SHAHRYARI et al. 2012).

***Gaeolaelaps praesternalis* (WILLMANN, 1949)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, October 2012. Collected by M. Mahjoori.

D i s t r i b u t i o n : Africa, Europe, Iran and Russia (EVANS & TILL 1966; VAN ASWEGEN & LOOTS 1970; GILYAROV et al. 1977; SOLEIMANI et al. 2011).

***Laelaspisella* MARAIS & LOOTS, 1969**

***Laelaspisella canestrinii* (BERLESE, 1903)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, July 2013; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, July 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran, Israel, Italy, Stavropol Territory and USSR (GILYAROV et al. 1977; CHERAGHALI et al. 2012).

***Gymnolaelaps* BERLESE 1916**

***Gymnolaelaps myrmecophila* (BERLESE, 1892)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Ganjeh, 227m, 36°51'23.26''N, 49°28'10.14''E, September 2012; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, July 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Western Europe, Finland and Iran (GILYAROV et al. 1977; HUHTA & KARG 2010; JOHARCHI & HALLIDAY 2013).

***Cosmolaelaps* BERLESE, 1903**

***Cosmolaelaps claviger* (BERLESE, 1882)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, September 2012; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, July 2013. Collected by M. Mahjoori

D i s t r i b u t i o n : Croatia, Europe, Iran, Latvia, Ukraine and USSR (EVANS & TILL 1966; GILYAROV et al. 1977; SALMANE 2000; KACZMAREK et al. 2009; BAROOZEH et al. 2013).

***Cosmolaelaps luteigiensis* SHCHERBEK, 1971**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, September 2012; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, September 2012; July 2013; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, July 2013; Nesfi, 455m, 36°50'44.85''N, 49°30'04.60''E, August 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran and Ukraine (SHCHERBAK 1971; HAJIZADEH et al. 2010).

***Cosmolaelaps vacua* (MICHAEL, 1981)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, October 2012; June 2013; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, June 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : America, Asia, Europe, Cis-Carpathia and Trans- Carpathia, Crimea, Iran, Lithuania, Ukraine, Uzbekistan and USSR (EVANS & TILL 1966; GILYAROV et al. 1977; KARG 2000; GWIAZDOWIEZ 2004; HAJIZADEH et al. 2010).

***Euanderolaelaps* BREGETOVA, 1977**

***Euanderolaelaps karawajewi* (BERLESE, 1903)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil and plant remains: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, June 2013; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, June 2013; Nesfi, 455m, 36°50'44.85''N, 49°30'04.60''E, July 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Asia, Europe, Iran, Israel, Ryazan oblast, Tataria, Trans-Carpathia, Trans- Caucasus, Ukraine, USSR, Western Kazakhstan and Western Siberia, (COSTA 1968; GILYAROV et al. 1977; KARG 1982; HAJIZADEH et al. 2010).

***Euanderolaelaps sardoa* (BERLESE, 1911)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Ganjeh, 227m, 36°51'23.26''N, 49°28'10.14''E, October 2012; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, October 2012; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, June 2013 & October 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Europe, Iran, and USSR (EVANS & TILL 1966; GILYAROV et al. 1977; DEHGHAN et al. 2011).

***Laelaspis* BERLESE, 1903**

***Laelaspis astronomica* (KOCHE, 1839)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, September 2012. Collected by M. Mahjoori.

D i s t r i b u t i o n : Armenia, Britain, Georgia, Hungary, Iran, Ukraine, USSR, Western Europe, (EVANS & TILL 1966; GILYAROV et al. 1977; KONTSCHÁN 2007; BALOOC SHAHRYARI et al. 2012).

***Laelaspis dariusi* (JOHARCHI & JALAEIAN, 2012)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil and plant remains: Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, September 2012; Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, June 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran (JOHARCHI et al. 2012a).

***Laelaspis pennatus* (JOHARCHI & HALLIDAY, 2012)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, October 2012, Taklim, 535m, 36°50'50.28''N, 49°24'01.12''E, August 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran (JOHARCHI et al. 2012b).

***Ololaelaps* BRELESE, 1903**

***Ololaelaps ussuriensis* BREGETOVA & KOROLEVA, 1964**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Roodbar, 237m, 36°49'26.86''N, 49°25'25.42''E, September 2012; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, July 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : Iran, Primorye Territory, USSR (GILYAROV et al. 1977; RAMROODI et al. 2013).

***Pseudoparasitus* OUDEMANS, 1902**

***Pseudoparasitus dentatus* (HALBERT, 1920)**

M a t e r i a l e x a m i n e d : Adult, Iran, Guilan province, collected from olive soil: Rostamabad, 170m, 36°53'54.00''N, 49°29'26.00''E, October 2013; Manjil, 366m, 36°44'31.80''N, 49°24'57.14''E, September 2013. Collected by M. Mahjoori.

D i s t r i b u t i o n : England; France, Iran, Italy, Northern and Central Europe; and USA (HUNTER 1966; GILYAROV et al. 1977; JOHARCHI et al. 2011).

Zusammenfassung

Im Zeitraum 2012 bis 2013 wurde in der Provinz Guilan im Norden des Iran eine faunistische Untersuchung der Familie Laelapidae (Acari: Mesostigmata) an Olivenplantagen durchgeführt. Nach Entnahme von Laub- und Bodenproben wurden die Milben in einem Berlese-Trichter extrahiert und auf Objekträgern montiert. Vorliegende Studie ergab Milben in 18 Arten und 9 Gattungen die allesamt Neunachweise für die Olivengärten Irans sind.

Literature

- BAROOZEH S., AHADIAT A., JOHARCHI O. & A.M. SARAFRAZI (2013): Species composition of mites of the family Laelapidae (Acari: Mesostigmata) in the main climate zones of Tehran Province, Iran. — Program & Abstract book of the Second International Persian Congress of Acarology: 7.
- BALOOCH SHAHRYARI N., HAJIZADEH J. & M. ASADI (2012): Introduction and identification key for laelapid mite (Mesostigmata: Laelapidae) of Jiroft Township. — Plant Pest Research **2** (2): 21-32.
- BERLESE A. (1882): Acari, Myriapoda, et Scorpiones hucusque in Italia reperta. — Padova **1**: 1-101.
- BERLESE A. (1892): Acari, Myriapoda, et Scorpiones hucusque in Italia reperta. Ordo Mesostigmata (Gamasidae). *Tipografia del Seminario*. — Padova **11**: 1-143.
- BERLESE A. (1903): Diagnosi di alcuni nuove specie di Acari italiani, mirmecofili e liberi. — Zoologischer Anzeiger **27**: 12-28.
- BERLESE A. (1910): Lista de nuovi specie e nuovi generi di Acari. — Redia **6**: 242-271.
- BERLESE A. (1911): Acarorum species novae quindecim. — Redia **7**: 429-435.
- BERLESE A. (1916): Centuria prima di Acari nuovi. — Redia **12**: 19-338.
- CHERAGHALI Z., JOHARCHI O., RASTEGAR J. & T. BOZORG (2012): Mites of the family Laelapidae (Acari: Mesostigmata) in Rudbar-ghasran region (Shemiranat), Tehran Province, Iran. — Abstract Book of the 20th Iranian Plant Protection Congress, Shiraz, Iran, p. 413.
- COSTA M. (1962): Mites from the nests of the mole-rat (*Spalax ehrenbergi*) in Israel. — Annals and Magazine of Natural History **13** (4): 481-503.
- COSTA M. (1968): Little known and new litter inhabiting Laelapine mites (Acari: Mesostigmata) from Israel. — Israel Journal of Zoology **17**: 1-30.
- COSTA M. (1974): Mesostigmatic mites (Acari: Mesostigmata) from the Mediterranean shores of Israel. I, The genus *Hypoaspis* CANESTRINI, 1884. — Israel Journal of Entomology **9**: 219-228.
- DEHGHAN Z., JOHARCHI O. & A. KHANI (2011): Mites of the family Laelapidae (Acari: Mesostigmata) in Yazd province, Iran. — Abstract and proceeding book of the First Persian Congress of Acarology, p. 16.
- EVANS G.O. & W.M. TILL (1966): Studies on the British Dermanyssidae (Acari: Mesostigmata). Part II. — Classification. — Bulletin of the British Museum (Natural History) Zoology **14** (5): 109-370.
- EVANS G.O. & W.M. TILL (1979): Mesostigmatic mites of Britain and Ireland (Chelicera: Acari-Parasitiformes). An introduction to their external morphology and classification. — The Transactions of the Zoological Society of London **35** (2): 139-270.
- GILYAROV M.S., BREGETOVA N.G., WAINSTEIN B.A., KADITE B.A., KOROLEVA E.V., PETROVA A.D., TIKHOMIROV S.I. & G.I. SHCHERBAK (1977): Manual of edaphic mites (Mesostigmata). — Akademiya Nauk SSSR. "Nauka" Publishing House. Leningrad Russia, 718 pp.

- HOSSEINI-MAZINANI S.M., SAMAEI S.M., SADEGHİ H. & J.M. CABALLERO (2004): Evaluation of olive germplasm in Iran on the basis of morphological traits: Assessment of Zard and Rowghani cultivars. — *Acta Horticulturae* **634**: 145-151.
- HADDAD IRANI-NEJAD K., HAJIGANBAR H.R. & P. TALEBI CHAICHI (2003): Introduction of some mesostigmatic mites of sugarbeet fields in Miandoab Plain. — *Journal of Agriculture Sciences and Natural Resources* **10** (2): 147-157.
- HAJIZADEH J., FARAJI F. & M. RAFATIFARD (2010): Laelapidae of Guilan Province, four new species records for the Iranian mite fauna and the identification key for Guilan laelapids. — *Journal of Plant Protection* **24** (2): 196-209.
- HUHTA V. & W. KARG (2010): Ten new species in genera *Hypoaspis* (s.lat.) Canestrini, 1884, *Dendrolaelaps* (s. lat.) Halbert, 1915, and *Ameroseius* Berlese, 1903 (Acari: Gamasina) from Finland. — *Soil Organisms* **82**: 325-349.
- HUNTER P.E. (1966): Some mites of the genus *Pseudoparasitus* Oudemans, 1902 (Acarina: Laelaptidae). — *Journal of the Georgia Entomological Society* **1**: 1-20.
- JOHARCHI O., JALAEIAN M., PAKTINAT-SAEIJ S. & A. GHAFARIAN (2012a): A new species and new records of *Laelaspis* BERLESE (Acari, Laelapidae) from Iran. — *ZooKeys* **208**: 17-25.
- JOHARCHI O., HALLIDAY B. & A. SABOORI (2012b): Three new species of *Laelaspis* BERLESE from Iran (Acari: Laelapidae), with a review of the species occurring in the Western Palaearctic Region. — *Journal of Natural History* **46** (31-32): 1999-2018.
- JOHARCHI O., HALLIDAY B., SABOORI A. & H. OSTOVAN (2011): Identification of Laelapidae (Acari: Mesostigmata) associated with insects and soil in Iran. — Abstract Book of Annual Zoological Congress of "Grigore Antipa" Museum, Bucharest, Romania: 131-132.
- JOHARCHI O. & B. HALLIDAY (2013): A new species and new records of *Gymnolaelaps* Berlese from Iran (Acari: Laelapidae), with a review of the species occurring in the Western Palaearctic Region. — *Zootaxa* **3646** (1): 039-050.
- KACZMAREK S., MARQUARDT T. & K. FALENCZYK-KOZIRÓG (2009): Checklist of soil Mesostigmata (Acari) of Central Croatia (Dalmatia) with some microenvironmental remarks. — *Polish Journal of Entomology* **78**: 177-184.
- KARG W. (1962): Zur Systematik und postembryonalen Entwicklung der Gamasiden (Acarina, Parasitiformes) landwirtschaftlich genutzter Boden. — *Mitteilungen aus dem Zoologischen Museum in Berlin* **38**: 23-119.
- KARG W. (1978): Die Gattung *Pseudoparasitus* OUDEMANS, 1902. — *Mitteilungen aus dem Zoologischen Museum in Berlin* **54**: 203-212.
- KARG W. (1982): Zur Kenntnis der Raubmilben *Hypoaspis* CANESTRINI, 1884 (Acarina, Parasitiformes). — *Mitteilungen aus dem Zoologischen Museum in Berlin* **58**: 233-256.
- KARG W. (1993): Rallbmilben: Acari (Acarina), Milben Parasitiformes (Allactinochaeta) Cohors Gamasina Leach. — *Tierwelt Deutsch.* **59**, Jena: Gustav Fischer.
- KARG W. (2000): Zur Systematik der Raubmilbenfamilien Hypoaspidae v. Vitzthum, 1941 und Rhodacaridae Oudemans, 1902 (Acarina, Parasitiformes) mit neuen Arten aus Süd- und Mittelamerika. — *Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe* **76**: 243-262.
- KRANTZ G.W. (1998): A new genus and two new species of hypoaspidine mites (Acari: Laelapidae) associated with old world carpenter bees of the tribe Xylocopini (Hymenoptera: Apidae: Xylocopa). — *International Journal of Acarology* **24**: 291-300.
- KACZMAREK S., MARQUARDT T. & K.F. KOZIRQG (2009): Checklist of soil Mesostigmata (Acari) of Central Croatia (Dalmatia) with some microenvironmental remarks. — *Polish Journal of Entomology* **78**: 177-184.
- LINDQUIST E.E. & G.O. EVANS (1965): Taxonomic concepts in the Ascidae, with a modified setal nomenclature for the idiosoma of the Gamasina (Acarina; Mesostigmata). — *Memoirs of the Entomological Society of Canada* **47**: 1-64.

- LINDQUIST E.E. (1994): Some observations on the chaetotaxy of the caudal body region of gamasine mites (Acaria: Mesostigmata), with a modified notation for some ventrolateral body setae. — *Acarologia* **35**: 323-326.
- LINDQUIST E.E., KRANTZ G.W. & D.E. WALTER (2009): Order Mesostigmata. — In: KRANTZ G.W. & D.E. WALTER (eds), *A Manual of Acarology*. Third Edition, Texas Tech University Press: 124-232.
- MINISTRY OF JIHAD AGRICULTURE of IRAN, Office of Statistics and Information Technology (2013): Crops and livestock statistics letter. Retrieved December 7 2013. Available from: <http://www.maj.ir/portal/Home/Default.aspx?CategoryID=20ad5e49-c727-4bc9-9254-de648a5f4d52>.
- NOORMOHAMMADI Z., HOSSEINI-MAZINANI M., BELAJ A., TRUJILLO I., RALLO L. & M. SADEGHIZADE (2007): Identification and classification of main Iranian olive cultivars using microsatellite markers. — *Hortscience* **42** (7): 1545-1550.
- OMRANI-SABBAGHI A., SHAHRIARI M., FALAHATI-ANBARAN M., MOHAMMADI S.A., NANKALI A., MARDI M. & B. GHAREYAZIE (2007): Microsatellite markers based assessment of genetic diversity in Iranian olive (*Olea europaea* L.) collections. — *Scientia Horticulturae* **112** (4): 439-447.
- RADOVSKY F.J. (1967): The Macronyssidae and Laelapidae (Acarina: Mesostigmata) parasitic on bats. — University of California Publications in Entomology **46** (8): 1-288.
- RAMROODI S., HAJIZADEH J. & O. JOHARCHI (2013): New records of the family Laelapidae (Acaria: Mesostigmata) from Guilan Province, and a new record for Iran. — Program & Abstract book of the Second International Persian Congress of Acarology: 33.
- SADEGHI H. (1992): Cultivation, preservation and harvesting of olive. Eds. — The publication of Ministry of Agriculture, Tehran, Iran.
- SHCHERBAK G.I. (1971): New species of gamasid from the genus *Hypoaspis* CANESTRINI Acarina, Gamasoidea. — *Vestnik Zoologii, Ukraine* **5**: 76-79.
- SHEIDAI M., VAEZI-JOZE S., SHAHREIYARI Z.H., NOORMOHAMMADI Z., PARSIAN H. & F. FARAHANEI (2007): Study of genetic diversity in some olive (*Olea europaea* L.) cultivars by using RAPD markers. — *Pakistan Journal of Biological Science* **10** (17): 2972-2975.
- SOLEIMANI M., OSTOVAN H. & O. JOHARCHI (2011): Mesostigmatic mites (Acaria: Mesostigmata) in Marvdasht region, Fars province, Iran. — Abstract and proceeding book of the First Persian Congress of Acarology: 17.
- STRONG K.L. & R.B. HALLIDAY (1994): Three new species of *Hypoaspis* CANESTRINI (Acarina: Laelapidae) associated with large Australian cockroaches. — *Journal of the Australian Entomological Society* **33**: 87-96.
- TENORIO J.M. (1982): Hypoaspidae (Acaria: Gamasida: Laelapidae) of the Hawaiian Islands. — *Pacific Insects* **24**: 259-274.
- VAN ASWEGEN P.I.M. & G.C. LOOTS (1970): A taxonomic study of the genus *Hypoaspis* CANESTRINI sens. lat. (Acaria: Laelapinae) in the Ethiopian region. — *Publicacoes Culturais Companhia de Diamantes de Angola Servicos Culturais* **82**: 167-213.

Authors' addresses:

Mojtaba MAHJOORI
 Jalil HAJIZADEH (corresponding author)
 Mohammad Reza Abbasi MOZHDEHI
 Department of Plant Protection, Faculty of Agricultural Sciences
 University of Guilan, Rasht, Iran
 E-mail: hajizadeh@guilan.ac.ir
 Mojtaba.Mahjoori91@yahoo.com
 Mr.mojdehi@areo.ir

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Linzer biologische Beiträge](#)

Jahr/Year: 2014

Band/Volume: [0046_2](#)

Autor(en)/Author(s): Mahjoori Mojtaba, Hajizadeh Jalil, Abbasii Mozhdehi
Mohammad Reza

Artikel/Article: [Mites of the family Laelapidae \(Acari: Mesostigmata\) associated with olive orchards in Guilan Province Iran 1599-1606](#)