

ciis latis fuscis pulcherrime ornata, epidermide cutacea viridi lutescente induta; spira depressa, subgradata, valde erosa. Anfr. superst. $4\frac{1}{2}$ convexi, supra planati sed vix angulati, ult. multo major, rotundatus, circa umbilicum vix levissime compressus. Apert. oblonga, supra et infra angustata, haud dilatata, faucibus fuscescentibus pulcherrime fasciatis; peristoma tenue, acutum, haud incrassatum, marginibus callo tenuissimo junctis. — Operculum normale, annulis prominentibus 3—4 insigne. — Diam. 35, alt. 35, alt. apert. obl. 25, lat. 16 mm.

Meladomus boettgeri Kobelt, in: Mart. Chemn. ed. II t. 30 f. 6, 7.

Hab. Gabun.

Diagnosen neuer Cyclotiden.

Von

Dr. W. Kobelt.

1. *Cyclotus (Opisthoporus) bialatus* (Moellendorff mss.).

T. parva, late umbilicata, depressa, solidula, subtiliter striatula, epidermide tenui, lutescente, adhaerente induta, strigis fulminulatis supra castaneis, infra peripheriam pallidioribus dein evanescentibus ornata. Spira breviter conica, prominula, summo unicolor castaneo, apice acutissimo, sutura impressa. Anfr. $4\frac{1}{2}$ regulariter accrescentes, inferi ad suturam planati, dein convexi, ultimus circa umbilicum compressus, antice valde descendens, 3 mm pone aperturam tubulo erecto retrorsum spectante insignis. Apertura peroliqua, circularis, plano irregulari; peristoma fusco-albidum, subduplex: internum continuum, haud porrectum, externum tenuissimum, extus late dilatatum et reflexum, supra et ad basin alato-productum, ad columellam subite evanescens. — Operculum normale.

Diam. maj. 10, min. 8, alt 7 mm.

Opisthoporus bialatus Moellendorff in coll.

Hab. — ?

2. *Cyclotus (Pseudocyclophorus) papuanus* (Moellendorff mss.).

T. late umbilicata, umbilico $\frac{3}{10}$ diametri aequante, depressa, solidula, sat ruditer striata, hic illic obsolete costulata, albida, epidermide lutescente induta, unicolor vel castaneo strigata et fasciata. Spira deppressa, apice breviter mucronato, sutura perprofunda. Anfractus $4\frac{1}{2}$ convexi, celeriter accrescentes, ultimus dilatatus, rotundatus, antice celeriter descendens. Apertura parum obliqua, circularis, peristomate expanso, super parietem aperturalem continuo, marginibus bene arcuatis, externo dupli. — Operculum planiusculum, lamellis 2 sulco profundo divisis compositum, externa majore ad margines prominente, anf. 7—8 ad suturam prominentibus, interna laevis, concava.

Diam. maj. 20, min. 14, alt. 11, diam. apert. 10 mm.

Cyclotus (Pseudocyclophorus) papuanus Moellendorff in coll. — Kobelt, in: Martini & Chemnitz Conch Cab ed. II p. 812 t. 121 fig. 6—8.

Hab. Takar, in parte boreali Novae Guineae occidentalis.

Var. *kapaurensis* n., differt a typo testa minore, supra saturati strigata, infra peripheriam fascia latiore sub irregulari castanea ornata. Diam. maj. 17, min. 13, alt. 9,5, diam. apert. 9 mm. (M. Ch. II p. 813 t. 121 fig. 9—11).

Hab. Kapaur in parte meridionali Novae Guineae occidentalis.

3. *Cyclotus (Pseudocyclophorus) rollei* Moellendorff mss.

T. modice sed profunde ac pervie umbilicata, globosoturbinata, solida, subtilissime striatula, sub epidermide fusca adhaerente griseo-albida, infra suturam alternatim strigis brevibus obliquis castaneis et lutescentibus ornata, infra

peripheriam obsoletissime castaneo fasciata. Spira conica, subgradata, apice acuto, castaneo; sutura linearis. Anfr. 5 convexi, leniter accrescentes, ult. rotundatus, antice distincte descendens. Apert. parum obliqua, circularis, fau-
cibus saturate castaneis; peristoma album, labio interno subduplicatum, leviter campanulatum dilatatum. — Operc.
magnum, terminale, testaceum, lamella externa majore,
ultra internam prominente, medio tantum spiram distinctam
exhibente, interna vix concava, ad marginem limbo in-
crassato quasi circumvallata.

Diam. maj. 17, min. 15, alt. 16, diam. apert. 9 mm.

Hab. Toll, Key Ins.

Cyclotus (Pseudocyclophorus) rollei Moellendorff in
sched. — Kobelt, in: Martini & Chemnitz, Conch. Cab
ed II t 128 f. 1—4.

4. *Cyclotus (Opisthoporus) kelantanensis* n. sp.

Testa late et aperte umbilicata, deppressa, tenuiuscula,
translucida, rosaceo-albida, supra pulcherrime castaneo
maculata et strigata, strigis ultra peripheriam productis;
spira fere plana apice parum prominulo, castaneo; sutura
profunda. Anfractus 4 convexi, ad suturam planati, reg-
ulariter crescentes, ultimus antice per profundè descendens,
fere 4 mm pone peristoma tubulo erecto humili munitus.
Apertura perobliqua, magna; peristoma rosaceum minime
incrassatum, duplex: internum continuum, filiforme, externum
complanulatum dilatatum, supra valde productum, ad mar-
ginem columellarem evanescens. — Operculum normale.

Diam. maj. 13, min. 11, alt. 8 mm.

Cyclotus (Opisthoporus) kelantanensis Kobelt, in:
Martini & Chemnitz Conch. Cab. ed II t. 26 fig. 19—21.

Kelantan.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Nachrichtsblatt der Deutschen Malakozoologischen Gesellschaft](#)

Jahr/Year: 1912

Band/Volume: [44](#)

Autor(en)/Author(s): Kobelt Wilhelm

Artikel/Article: [Diagnosen von Caclotiden 8-10](#)