

Three new localities of reptiles in Greece are referred in this note.

Mauremys caspica (GMELIN, 1774). This species was found in Gavdos island, 40 km south of the island of Crete. Four adults and 5-6 juveniles were observed in a small pool, near the church of Agios Pavlos, in May 1987.


Coluber najadum (EICHWALD, 1831). This species was found in Lesvos island near to the village of Moria, in July 1987. A juvenile of about 15 cm was captured.

Lacerta agilis LINNAEUS, 1758. This species was found in Vigla region of Florina state (North Greece). Ten specimens were collected near the ski center building in May 1984.

The determination of the species was based on the guide by ARNOLD & BURTON (1978).

### Bibliography

ARNOLD, E. N. & BURTON, J. A. (1978): A Field Guide to the Reptiles and Amphibians of Europe. London (Collins).



Efstratios Valakos, University of Athens, Department of Biology  
Panepistimioupolis, Ilisia,  
GR - 157 - 71, Athens, Greece

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [ÖGH - Nachrichten](#)

Jahr/Year: 1987

Band/Volume: [12-13 1987](#)

Autor(en)/Author(s): Valakos Efstratios D.

Artikel/Article: [New locality records of three Greek reptiles 70-71](#)