Phyton (Austria) Special issue:	Vol. 39	Fasc. 3	(175)-(180)	30. 11. 1999
Plant Physiology				

# Pollen Grain Bioassay for Environmental Pollution Screening

By

## Jasna PARADIŽ<sup>1)</sup> & Milan LOVKA<sup>1)</sup>

Key words: Radioactive contamination, cytogenetic bioassay, Vicia faba, perennials.

#### Summary

PARADIŽ J. & LOVKA M. 1999. Pollen grain bioassay for environmental pollution screening. - Phyton (Horn, Austria) 39 (3): (175) - (180).

Pollen grain assays in *Vicia faba* and naturally growing plants indicated increased genotoxic effects in habitats with elevated radiation levels. However, severe cytological damage in test plants could not be attributed solely to in situ genotoxic effect of prolonged irradiation, but to ageing of plants and complex clastogenic effects in the environment, including harsh weather conditions. Pollen grain assay in different plant species indicated that perennials could be used as biological indicators of long-term genotoxic exposure.

#### Introduction

Plants have been used for the detection of genotoxic agents in the environment for decades. In short-term testing procedure, traditionally the *Tradescantia* micronucleus assay (GILL & SANDHU 1992), *Tradescantia* stamenhair mutation assay (GICHNER & al. 1994a), *Arabidopsis* assay (GICHNER & al. 1994b), *Vicia faba* root tip assay (GRANT & SALAMONE 1994), *Allium* test (SMAKA-KINCL & al. 1996), and Tobacco mutagenicity assay (GICHNER & PLEWA 1998) are utilised. Test plants have been used for the detection of mutagens in industrial areas (RUIZ & al. 1992, GRANT & al. 1992) as well as for the mutagenicity screening of radioactive contamination after the Chernobyl accident (CEBULSKA-WASILEWSKA 1992). But recently the relevant strategy lies in the use

<sup>&</sup>lt;sup>1)</sup> National Institute of Biology Ljubljana, Večna pot 111, 1000 Ljubljana, Slovenia. E-mail: jasna.paradiz@uni-lj.si

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at (176)

of cytogenetic methods in natural vegetation (TOMKINS & GRANT 1976, SHEVCHENKO & GRINIKH 1990, MIČIETA & MURIN 1996, JABEEN & ABRAHAM 1996, BESSONOVA & al. 1996, KORDYUM & SIDORENKO 1997) for the in situ evaluation from environmental pollution.

We performed pollen grain bioassay in surrounding of the former uranium mine and yellow cake production plant at Žirovski Vrh as well as at the Krško Nuclear Power Plant (NPP). *Vicia faba* as test plants was used for the determination of short-term radiation effects as well as naturally growing plants as biomonitors of long-term exposure effects to assess in situ genotoxic hazards.

#### Material and Methods

In the surroundings of Žirovski Vrh, three experimental sites were chosen, where radionuclide levels were constantly measured (Table 1). Site I was on the slope of the hill called Boršt in a drainage ditch about 20 m down the dry-tailings pile. Site II was beyond the valley, opposite to site I and supposedly contaminated by radioactive particles from the Boršt deposit. Site III was close to the former yellow-cake production plant in Todraž. The Botanical Garden of the University of Ljubljana was the control location with no anthropogenic radiation sources.

Routine discharges from the Krško NPP are known to contribute only local environmental radioactivity, and according to measured activities in the environment, the radiological impact is of minor significance (BREZNIK 1996, MIKLAVŽIČ & al. 1996). Inside the Krško NPP area, one experimental site was chosen; the one was outside the NPP area. The habitats of wild experimental plants were about 1 km SE from the NPP.

Table 1. Radionuclides (Bq/kg dry sample) in soil and absorbed dose rate in air ( $\mu$ Gy/h) at experimental sites in the area of the Žirovski Vrh.

	U-238	Ra-226	Th-232	K-40	Cs-137	Dose rate
Site I	1770	1900	41	480	. 12	0.35
Site II	51	55	62	670	367	0.13
Site III	5600	840	65	560	22	0.18

Ten soaked seeds of *Vicia faba* were germinated in soil taken from each experimental site as well as in the normal garden soil. When about 20 cm high, potted seedlings were transferred to experimental sites. At each site, two experimental series of plants, those potted in the soil from the site and those in garden soil, were exposed to naturally occuring conditions. Additionally, all were grown in the Botanical Garden. After 3, 6, and 8 weeks of exposure, the flower buds were sampled.

From naturally growing plants, flower buds were taken from *Berberis communis*, Brassica oleracea, Cornus sanguinea, Crataegus monogyna, Plantago lanceolata, Plantago media, Polygonatum multiflorum, Ranunculus acris, Ranunculus lanuginosus, Sambucus nigra, and Symphytum officinale. For the controls, flower buds from the same plant species were sampled on sites with no anthropogenic radiation sources.

The flower buds were fixed in 3:1 ethanol-acetic acid, then kept at -20°C. The slides were prepared by squashing anthers in aceto-carmine. In fully developed anthers, aberrant shapes and sizes of pollen grains as well as empty pollen grains were scored (Figs. 1-3). In each sample, 500 pollen grains were observed.

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at
(177)

### Results and Discussion

Pollen deformation in *Vicia faba* (Figs. 4 and 5) grown in soil from experimental sites was more marked than in plants grown in the garden soil. In the botanical garden, the frequency of pollen deformation in plants grown in the soil from experimental sites was higher than in those grown in the garden soil. With prolonged exposure, the degree of pollen deformation increased in all plants at experimental sites as well as in the botanical garden. An extremely high degree of pollen deformation was predominantly due to numerous sterile flower buds with anthers with no pollen grains.



Figs. 1-3. Deformation of pollen grains in Vicia faba. Bar represents 10 micrometers.

*Vicia faba* pollen grain assay results indicated the impact of radionuclides from soil as well as their possible transfer and accumulation in plant organs, but the influence of environmental conditions on the results could not be excluded.

The average frequency of pollen deformation was 33.7% in *Plantago lanceolata*, 22.9% in *Ranunculus lanuginosus*, and 16.8% in *Crataegus monogyna*; however, similarly high frequencies in the control plants were observed as well (Fig. 6). Variability of the results in all experimental plants was obvious; the

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at (178)

degree of variation likely reflect instability of cytomorphological features and physiological properties in irradiated natural plant populations (POZOLTINA 1996).



Fig. 4. Vicia faba pollen grain assay in the area of Žirovski Vrh. 1 – Plants potted in soil from site I and exposed to site I. 2 – Plants potted in garden soil and exposed to site I. 3 – Plants potted in soil from site I and grown in Botanical Garden. 4 – Plants potted in soil from site II and exposed to site II. 5 – Plants potted in garden soil and exposed to site II. 6 – Plants potted in soil from site II and grown in Botanical Garden. 7 – Plants potted in soil from site III and exposed to site III. 8 – Plants potted in garden soil and exposed to site III. 8 – Plants potted in garden soil and exposed to site III. 9 – Plants potted in soil from site III and grown in Botanical Garden. 10 – Plants potted in garden soil and grown in Botanical Garden.



Fig. 5. Vicia faba pollen grain assay in the area of the Krško NPP.

(179)



Fig. 6. Pollen grain assay in naturally grown plants close to the Krško NPP.

#### References

- BESSONOVA V.P., GRITSAI Z.V. & YUSYPIVA T.I. 1996. The use of cytogenetic methods for the assessment of industrial pollutant mutagenicity. - Tsitologiya i Genetika 30 (5): 70-76.
- BREZNIK B. 1996. Radiation protection programme at Krško Nuclear Power Plant. Symposium on radiation protection in neighbouring countries in central Europe - 1995, Portorož, Slovenia, Proceedings Session VI: 385-389.
- CEBULSKA-WASILEWSKA A. 1992. *Tradescantia* stamen-hair mutation bioassay on the mutagenicity of radioisotope-contaminated air following the Chernobyl nuclear accident and one year later. Mutat. Res. 270: 23-29.
- GILL B.S. & SANDHU S.S. 1992. Application of the *Tradescantia* micronucleus assay for the genetic evaluation of chemical mixtures in soil and aqeous media. Mutat. Res. 270: 65-70.
- GICHNER T. & PLEWA M.J. 1998. Induction of somatic DNA damage as measured by single cell gel electrophoresis and point mutation in leaves of tobacco plants. - Mutat. Res. 401: 143-152.
  - , CABRERA LOPEZ G., WAGNER E.D. & PLEWA M.J. 1994 a. Induction of somatic mutations in *Tradescantia* clone 4430 by three phenylenediamine isomers and the antimutagenic mechanisms of diethyldithiocarbamate and ammonium meta-vanadate. - Mutat. Res. 306: 165-172.
  - BADAYEV S.A., DEMCHENKO S.I., RELICHOVA J., SANDHU S.S., USMANOV P.D., USMANOVA O. & VELEMINSKY J. 1994 b. Arabidopsis assay for mutagenicity. - Mutat. Res. 310: 249-256.
- GRANT W.F. & SALAMONE M.F. 1994. Comparative mutagenicity of chemicals selected for test in the International program on chemical safety's collaborative study on plants systems for the detection of environmental mutagens. - Mutat. Res. 310: 187-209.
- ---, LEE H.G., LOGAN D.M. & SALAMONE M.F. 1992. The use of *Tradescantia* and *Vicia faba* bioassays for the in situ detection of mutagens in an aquatic environment. Mutat. Res. 270: 53-64.
- JABEEN C. & ABRAHAM S. 1996. Impact of gaseous pollutants from the kerala minerals and metals Ltd. on meiosis in *Datura metel* L. - Caryologia 49 (2): 193-197.
- KORDYUM E.L. & SIDORENKO P.G. 1997. The results of cytogenetic monitoring in angiosperm species of plants in the zone of radionuclide contamination after the Chernobyl NPP accident. - Tsitologiya i Genetika 31 (3): 39-46.

©Verlag Ferdinand Berger & Söhne Ges.m.b.H., Horn, Austria, download unter www.biologiezentrum.at (180)

- MIČIETA K. & MURIN G. 1996. Microspore analysis for genotoxicity of a polluted environment. -Environ. Exp. Bot. 36 (1): 21-27.
- MIKLAVŽIČ U., MARTINČIČ R., KANDUČ M., LULIĆ S., KOVAČ J. & BREZNIK B. 1996. Radioactivity monitoring programme of Krško Nuclear Power Plant. - In: Symposium on radiation protection in neighbouring countries in central Europe - 1995, Portorož, Slovenia, Proceedings Session VI: 379-384.
- POZOLTINA V.N. 1996. Adaptation processes of plants under the influence of radiation. Russ. J. Ecology 27 (2): 107-112.
- RUIZ E.F., RABAGO V.M.E., LECONA S.U., PEREZ A.B. & MA T.H. 1992. Tradescantiamicronucleus (Trad-MNC) bioassay on clastogenicity of wastewater and in situ monitoring. - Mutat. Res. 270: 45-51.
- SHEVCHENKO V.V. & GRINIKH L.I. 1990. Cytogenetic effects in natural Crepis tectorum populations subjected to chronic irradiation within the zone of Chernobyl accident. -Radiobiologija 30 (6): 728-734.
- SMAKA-KINCL V., STEGNAR P., LOVKA M. & TOMAN M.J. 1996. The evaluation of waste, surface and ground water quality using the *Allium* test procedure. - Mutat. Res. 368: 171-179.
- TOMKINS D.J. & GRANT W.F. 1976. Monitoring natural vegetation for herbicide-induced chromosomal aberrations. Mutat. Res. 36: 73-84.

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Phyton, Annales Rei Botanicae, Horn

Jahr/Year: 1999

Band/Volume: 39\_3

Autor(en)/Author(s): Paradiz Jasna, Lovka Milan

Artikel/Article: <u>Pollen Grain Bioassay for Environmental Pollution</u> <u>Screening. 175-180</u>