

---

***Attagenus (s. str.) pseudorobustior n. sp. (Coleoptera: Dermestidae: Attagenini)  
from Namibia***

Author(s): Andreas Herrmann & Jiří Háva

Source: Stuttgarter Beiträge zur Naturkunde A, 9(1):177-179.

Published By: Stuttgart State Museum of Natural History

<https://doi.org/10.18476/sbna.v9.a8>

URL: <http://www.bioone.org/doi/full/10.18476/sbna.v9.a8>

---

BioOne ([www.bioone.org](http://www.bioone.org)) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/page/terms\\_of\\_use](http://www.bioone.org/page/terms_of_use).

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

# *Attagenus* (s. str.) *pseudorobustior* n. sp.

## (Coleoptera: Dermestidae: Attagenini) from Namibia

ANDREAS HERRMANN &amp; JIŘÍ HÁVA

## A b s t r a c t

*Attagenus pseudorobustior* n. sp. from Namibia is described, illustrated and compared with the similar species *Attagenus robustior* (Pic, 1951). A list of all *Attagenus* species so far recorded from Namibia is given.

**K e y w o r d s :** Taxonomy, new species, nomenclature, Coleoptera, Dermestidae, Attagenini, *Attagenus*, Namibia.

## Z u s a m m e n f a s s u n g

*Attagenus pseudorobustior* n. sp. aus Namibia wird beschrieben, abgebildet und mit der sehr ähnlich aussehenden Art *Attagenus robustior* (Pic, 1951) verglichen. Alle bisher aus Namibia gemeldeten Arten dieser Gattung werden aufgelistet.

## C o n t e n t s

1	Introduction .....	177
2	<i>Attagenus pseudorobustior</i> n. sp. ....	177
3	List of the <i>Attagenus</i> species from Namibia .....	179
4	References .....	179

## 1 Introduction

Currently the dermestid genus *Attagenus* Latreille, 1802 contains more than 200 known species and subspecies worldwide (HÁVA 2015), the majority of them belonging into the subgenus *Attagenus* (s. str.). The present paper considers the preceding taxonomical publications on the species of *Attagenus* from the Afrotropical region (e.g. THUNBERG 1795, 1815; REITTER 1881; PÉRINGUEY 1885, 1886; KALÍK 1955; KALÍK & HÁVA 2005; HÁVA 2003a, 2003b; HERRMANN & HÁVA 2014). During the determination of some material from South Africa deposited in the collection of the SMNS the authors found a new species of the genus *Attagenus* which already was recognised by the deceased coleopterist VLADIMÍR KALÍK (1918–2010) as being new to science (see Remarks). This new species is described below.

### Acronyms of type depositories

AHEC	Private collection of ANDREAS HERRMANN, Stade, Germany
JHAC	Private Entomological Laboratory and Collection JIŘÍ HÁVA, Prague, Czech Republic
SMNS	Staatliches Museum für Naturkunde, Stuttgart, Germany

### Abbreviations

EW	elytral width = maximum elytral width
TL	total length = distance from anterior margin of pronotum to apex of elytra

### A c k n o w l e d g e m e n t s

The authors are deeply indebted to JOHANNES REIBNITZ (SMNS) for taking photos of the new species, and to Dr. WOLFGANG SCHAWALLER (SMNS) for the loan of specimens.

## 2 *Attagenus pseudorobustior* n. sp.

(Figs. 1–4)

Nomen nudum: *Attagenus roeri* Kalík: SCHAWALLER 1994: 27 (see Remarks).

**H o l o t y p e** (♂): “SWA / Namibia, Nyanga, 14.–22.I.1985, leg. H. ROER”.

**P a r a t y p e s** (not sexed): 39 specimens, same data as holotype.

All specimens of the newly described species are provided with the following labels: “HOLOTYPE” [or “PARATYPE” respectively], “*Attagenus pseudorobustior* sp. n., A. HERRMANN & J. HÁVA det. 2015”.


The holotype is stored in SMNS, paratypes are deposited in SMNS, AHEC and JHAC.

### Etymology

The name of the new species refers to its similarity to *Attagenus robustior* (Pic, 1951).

### Description

**Holotype** (♂): Body longish ovate, slightly arched. Measurements: TL 3.5 mm, EW 2.1 mm. Head slightly



**Figs. 1–4.** *Attagenus pseudorobustior* n. sp., ♂ holotype (1–3), ♀ paratype (4). – 1. Habitus, dorsal aspect. 2. Antenna. 3. Aedeagus. 4. Habitus, dorsal aspect. – Scales: 1 mm (1, 4), y mm (2, 3).

broader than long, sparsely and finely punctate, densely covered with long grey, decumbent hairs; frons flattened and shiny; palpi light brown; Ocellus present on front. Eyes large with hardly visible microsetae. Antennae 11-segmented; antennal club consisting of 3 antenno-meres, combined shorter than the shaft (Fig. 2); terminal segment nearly as long as the preceding two segments combined; antennal club brown to darkish brown, shaft light brown. Pronotum twice as broad as long, arcuately narrowed anteriorly, finely and sparsely punctate (like head), cuticula shiny, entirely covered by uniform long grey, decumbent hairs; anterior angles rounded, not visible from above; posterior angles acute with rounded edge, middle of the base convexly extended. Scutellum small and triangular. Punctuation and pubescence of the elytra similar to that of head and pronotum, but punctuation somewhat more rough and dense; cuticle reddish brown with several indistinct black spots and fasciae, hardly visible because of the dense pubescence as shown in Fig. 1. Legs brown with long and coarse yellow brown pubescence; tarsi roughly as long as tibiae, the latter with strong spines on the exterior edge arranged in two irregular lines. Mesosternum, metasternum and all visible abdominal sternites with the same kind of punctuation and pubescence as in elytra. Male genitalia as shown in Fig. 3.

**Female:** Habitually quite similar to male, slightly more elongated. The antennal club somewhat smaller (as usual in this genus), the spots and fasciae much more distinct than in male, the hairs located on the black elytral parts are darkened (Fig. 4).

**Variability:** Measurements of paratypes: TL 2.9–3.8 mm, EW 1.8–2.3 mm. Dimension, shape and position of spots and fascia on the elytra are slightly variable.

#### Differential diagnosis

The new species strongly resembles *Attagenus robustior* (Pic, 1951) (figure of habitus see <http://www.dermestidae.com/Attagenusrobustior.html>), but differs in the following characters:

*A. robustior*: Antennal club black, broad; body narrower, pubescence of pronotum with several spots and patches, legs dark brown to black, male genitalia almost circular in dorsal aspect (Congo, Namibia, Senegal, Tanzania).

*A. pseudorobustior* n. sp.: Antennal club brown and more narrow; body broader, pubescence of pronotum uniform without any spots or patches, legs brown, male genitalia elongated in dorsal aspect (Namibia).

The new species differs from all other species of the genus occurring in Namibia in shape and arrangement of the elytral spots.

## Remarks

KALÍK's dermestids are stored in several institutions. Some of them are new to science and already got a name by him and are labelled as types, but they were never published as valid descriptions. This concerns also a couple of beetles labelled as "*Attagenus roeri* sp. n." which are deposited in the collection of the SMNS. This species is now described in the present paper. SCHAWALLER (1994) cited the material from SMNS erroneously under the name *Attagenus roeri* Kalík in his type catalogue of Coleoptera from SMNS.

## 3 List of the *Attagenus* species from Namibia

- Attagenus aurofasciatus* Háva, 2005. – Distribution: Namibia, South Africa.  
*Attagenus cinereus* (Thunberg, 1815). – Distribution: Namibia, South Africa.  
*Attagenus fasciatus* (Thunberg, 1795). – Distribution: cosmopolitan.  
*Attagenus havai* Kadej, 2006. – Distribution: Angola, Namibia.  
*Attagenus heinigi* Herrmann & Háva, 2007. – Distribution: Namibia.  
*Attagenus kaniai* Háva & Kadej, 2008. – Distribution: Namibia, Zimbabwe.  
*Attagenus muehlei* Herrmann & Háva, 2014. – Distribution: Namibia.  
*Attagenus pardus* Arrow, 1915. – Distribution: Botswana, Namibia, South Africa, Tanzania, Zimbabwe.  
*Attagenus pellio* (Linnaeus, 1758). – Distribution: cosmopolitan.  
*Attagenus pseudorobustior* n. sp. – Distribution: Namibia.  
*Attagenus robustior* (Pic, 1951). – Distribution: Congo, Namibia, Senegal, Tanzania.

## 4 References

- HÁVA, J. (2003a): New and interesting Dermestidae from Burkina Faso (Coleoptera). – *Lambillionea* **103**: 500–504.  
HÁVA, J. (2003b): Notes on Dermestidae (Coleoptera) with description of eight new species. – *Annales historico-naturales Musei nationalis hungarici* **95**: 19–35.  
HÁVA, J. (2015): World Catalogue of Insects. Volume 13. Dermestidae (Coleoptera), XXVI + 419 pp.; Leiden, Boston (Brill).  
HERRMANN, A. & HÁVA, J. (2014): Description of two new species of the genus *Attagenus* Latreille, 1802 from the Afrotropical Region (Dermestidae: Attageninae). – *Studies and Reports, Taxonomical Series* **10**: 93–98.  
KALÍK, V. (1955): Dermestidae der aethiopischen Region aus den Sammlungen des Königl. Museums von Belgisch Kongo in Tervuren (Mit Bemerkungen über verwandte Arten). – *Revue de Zoologie et de Botanique Africaines* **52**: 294–322.  
KALÍK, V. & HÁVA, J. (2005): *Attagenus holmi* n. sp. (Coleoptera: Dermestidae) from South Africa, with notes on *A. prescutellaris* Pic and *A. rufiventris* Pic. – *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* **686**: 7 pp.  
REITTER, E. (1881): Die aussereuropäischen Dermestiden meiner Sammlung. Mit 70 Diagnosen neuer Arten. – *Verhandlungen des naturforschenden Vereines in Brünn* **19**: 27–60.  
PÉRINGUEY, L. (1885): First contribution to the South-African coleopterous Fauna. – *Transactions of the South African philosophical Society* **3**: 74–149, 4 pls.  
PÉRINGUEY, L. (1886): Second contribution to the South-African coleopterous Fauna. – *Transactions of the South African philosophical Society* **4**: 67–194, 4 pls.  
SCHAWALLER, W. (1994): Die Käfersammlung am Naturkundemuseum Stuttgart. – *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)* **508**: 40 pp.  
THUNBERG, C. P. (1795): *Dissertatio entomologica sistens insecta Svecica. Pars 9*, pp. 105–113; *Upsaliae* [Uppsala] (Edman).  
THUNBERG, C. P. (1815): *Anthreni monographia*. – *Nova Acta Regiae Societatis Scientiarum Upsaliensis* **7**: 150–156.

## Authors' addresses:

ANDREAS HERRMANN, Bremervörder Straße 123, 21682 Stade, Germany;  
e-mail: [herrmann@coleopterologie.de](mailto:herrmann@coleopterologie.de)

JIŘÍ HÁVA, Department of Forest Protection and Entomology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Kamýcká 1176, 16521 Prague 6 – Suchdol, Czech Republic;  
e-mail: [jh.dermestidae@volny.cz](mailto:jh.dermestidae@volny.cz)

Manuscript received: 11.II.2015, accepted: 29.VIII.2015.

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Stuttgarter Beiträge Naturkunde Serie A \[Biologie\]](#)

Jahr/Year: 2016

Band/Volume: [NS\\_9\\_A](#)

Autor(en)/Author(s): Herrmann Andreas, Hava (Háva) Jiri

Artikel/Article: [Attagenus \(s. str.\) pseudorobustior n. sp. \(Coleoptera: Dermestidae: Attagenini\)  
from Namibia 177-179](#)