

WISSENSCHAFTLICHE KURZMITTEILUNGEN

Population dynamics of the Common opossum, *Didelphis marsupialis* (Mammalia, Marsupialia), in southern Brazil

By N. C. Cáceres and E. L. A. Monteiro-Filho

Departamento de Zoologia, Universidade Federal do Paraná, Curitiba, Brasil

Receipt of Ms. 01. 04. 1997 Acceptance of Ms. 10. 12. 1997

Key words: Didelphis marsupialis, population density, sexual rate, age structure

Nowadays, the majority of the American marsupial species is found distributed in the Neotropical Region and included in three families (KIRSCH 1977). The common opossum, *Didelphis marsupialis* Linnaeus, 1758, one of these marsupials, shows a clear neotropical distribution occurring from Mexico to northern Argentina (HERSHKOVITZ 1969). Despite its abundancy in urban environments (HUSSON 1978), the main studies concerning the population dynamics of the common opossum in Brazil were carried out by DAVIS (1945), CERQUEIRA et al. (1993), and CHEREM et al. (1996). Thus, as there is little demographic data available to understand the general ecology of this opossum, this study aims to verify its sexual rate, age structure, population density, recruitment, and mortality.

This study was carried out in an area of Curitiba City, State of Paraná, southern Brazil ($25^{\circ}25'$ S and $49^{\circ}18'$ W), at 940 m above sea level. The area has five hectares covered with mixed ombrophyllous forest, and it has many roads, houses, and buildings in the vicinity. The mean annual temperature is 16.5 °C (MAACK 1981). It is possible to define two distinct seasons in Curitiba City, the dry season (April to August) and the wet season (September to March), in accordance with climatic data obtained at the Meterorological Station of the Universidade Federal do Paraná, 8 km distant from the study area.

Opossums were captured in 30 live traps $(40 \times 20 \times 20 \text{ cm})$ that were uniformly placed in the study area between February 1995 and January 1996. These live traps were baited during the afternoon and observed for opossum captures the following morning once per week. The bait was ripe banana with peanut butter and codfish liver oil. After capture, animals were sexed, aged (TYNDALE-BISCOE and MACKENZIE 1976), marked by combinations of holes in each ear and released. Sexes of pouch young were determined taking care not to detach them from teats. A binomial distribution test (ZAR 1984) was utilized to verify if there was a significant difference between expected (1:1) and observed sexual rates. For a better understanding of the age structure, the opossum permanence in the area was verified taking into account the time between the first and the latter capture day of each individual. The immigration rate was obtained by taking the proportion of the number of new individuals captured during each period of two months divided per the total of individuals in the area during the last two months. Inversely, the emmigration rate was obtained by taking the individuals that were not recaptured during each period of two months. The density and migration rates were based on the minimum number known to be alive in the study area. The mortality was verified through direct observation in the study area and its surroundings during the field phases.

N. C. Cáceres and E. L. A. Monteiro-Filho

After 12 months of studies, 1761 baited traps resulted in 192 opossum captures. The estimated population size was 18 opossums, with 11 females and seven males (1.6 females: 1.0 male). Utilizing the binomial distribution test (ZAR 1984), the sexual rate observed for adults and subadults did not differ significantly from the expected rate (P > 0.12). The same occurred for pouch young (P > 0.10) with 25 females and 21 males (1.2 females: 1.0 male).

Juveniles and subadults (all less than one year old) were captured during the end of the wet season and the beginning of the dry season, respectively. Some of them (three females) remained in the area until the adult stage (seven to 12 months). Two older adult females were recaptured for a time longer than eight months. Adult males (about one year old) and females with pouch young were captured mainly during the wet season (September to January and August to January, respectively). Males remained in the area for three months at maximum.

The mean population density was 1.4 opossums/ha (range: 0.8 to 2.2 individuals/ha). The immigration rate showed a positive correlation with the density (r = 0.91, P < 0.05). The emmigration rate did not contribute to density fluctuations (r = 0.00, P < 0.05). Based on births, the density was high during August and November (8.2 and 6.8 pouch young/ha, respectively), and was zero from February to July.

There were two common opossum deaths in the area. One death was as a result of human contact. On the other hand, several *D. albiventris* Lund, 1841 were found dead on roads near the study area.

Sexual rates observed here corroborate some other studies on *Didelphis* (Stout and SONENSHINE 1974; ATRAMENTOWICZ 1986; CHEREM et al. 1996). However, there was an untested trend for a greater number of female opossums in the population (see Holmes and SANDERSON 1965; ATRAMENTOWICZ 1986). The higher rate of females in the study area is most probably related to this trend. Studies of DAVIS (1945), ATRAMENTOWICZ (1986), SUNQUIST et al. (1987) and O'CONNELL (1989) also revealed greater female permanence in other regions. The absence of males in the study area during the dry season is probably related to non-oestrous females that possibly avoided them during this time (see MOTTA et al. 1983). The study of RYSER (1992) on *Didelphis virginiana* Kerr, 1792 gives some evidence to support this, with females showing stable home ranges and males increasing theirs during the breeding season while seeking for mates.

The presence of pouch young was only observed during a certain period of the year. This is in agreement with studies of FLEMING (1973), TYNDALE-BISCOE and MACKENZIE (1976), CERQUEIRA et al. (1993), and CHEREM et al. (1996) concerning the same species. Juveniles were also seasonally captured, which is in agreement with the study of FLEMING (1972) in Panamá. The common opossum probably showed this population pattern in response to advantages that their young gain if they are weaned during a season with widely available resources (see FLEMING 1973; JULIAN-LAFERRIERE and ATRAMENTOWICZ 1990). Some cited evidences of seasonality in the population succession were already reported by O'CONNELL (1989) and CERQUEIRA et al. (1993) from other sites in South America.

Our observations in southern Brazil were in accordance with other authors that reported a rapid turnover, every two or three years, for the common opossum in northern South America (ATRAMENTOWICZ 1986; SUNQUIST et al. 1987; SUNQUIST and EISENBERG 1993). Hence, we noticed that the great majority of captured opossums originated from the preceeding breeding season as reported by O'CONNELL (1989) from Venezuela.

Except for the study of August (1984) in the llanos of Venezuela, other estimated densities for the common opossum (FLEMING 1972; CHARLES-DOMINIQUE et al. 1981; ATRAMENTOWICZ 1986; SUNQUIST et al. 1987; O'CONNELL 1989; CERQUEIRA et al. 1993) were similar to the density verified here.

Considering the greater difference between the high densities of pouch young and the

170

low abundance of independent opossums in our field area, it is probable that relatively few individuals survive until maturity, which is in agreement with data of ATRAMENTOWICZ (1986).

The main factor of opossum mortality in the study area was human action. Based on our data, the common opossum must be less often run over by cars than the white-eared opossum in Curitiba City. GARDNER (1983) reported that the common opossum is often run over by cars in Costa Rica. On the other hand, ATRAMENTOWICZ (1986) and SUNQUIST and EISENBERG (1993) reported that the main mortality causes in natural environments outside urban areas are wild predators.

Therefore, the common opossum showed a population dynamic subject to seasonal variations, exhibiting a population density strongly related to births and immigrations. Evidences of sexual demographic differences were observed, with females tending to remain in forested sites, when the species occurs in urban environment. This trend may reflect an attempt to better guarantee reproductive success in these sites.

Acknowledgements

We would like to thank V. M. CáCERES-JÚNIOR for help in the field and J. QUADROS, E. SEEGAR, and S. SMITH for help in translating the manuscript. This study was supported by the Curso de Pós-Graduaçao em Zoologia (UFPR), CAPES and CNPq.

References

- ATRAMENTOWICZ, M. (1986): Dynamique de Population chez Trois Marsupiaux Didelphidés de Guyane. Biotropica 18, 136–149.
- AUGUST, P. V. (1984): Population ecology of small mammals in the llanos of Venezuela. Spec. Publ. Mus. Texas Tech. Univ. 22, 71–104.
- CERQUEIRA, R.; GENTILE, R.; FERNANDEZ, F. A. S.; D'ANDREA, P. S. (1993): A five-year population study of an assemblage of small mammals in southeastern Brazil. Mammalia 57, 507–517.
- CHARLES-DOMINIQUE, P.; ATRAMENTOWICZ, M.; CHARLES-DOMINIQUE, M.; GÉRARD, H.; HLADIK, A.; HLADIK, C. M.; PRÉVOST, M. F. (1981): Les mammiferes frugivores arboricoles nocturnes d'une forest guyanaise: interrelations plantes-animaux. Rev. Ecol. Terre Vie 35, 341–435.
- CHEREM, J. J.; GRAIPEL, M. E.; MENEZES, M. E.; SOLDATELI, M. (1996): Observações sobre a biologia do gambá (*Didelphis marsupialis*) na Ilha de Ratones Grande, Estado de Santa Catarina, Brasil. Biotemas 9, 47–56.
- DAVIS, D. E. (1945): The annual cycle of plants, mosquitoes, birds, and mammals in two Brazilian forests. Ecol. Monogr. 15, 243–295.
- FLEMING, T. H. (1972): Aspects of the population dynamics of three opossums in the Panama Canal Zone. J. Mammalogy **53**, 619–623.
- FLEMING, T. H. (1973): The reproductive cycles of three species of opossums and other mammals in the Panama Canal Zone. J. Mammalogy **54**, 439–455.
- GARDNER, A. L. (1983): *Didelphis marsupialis*. In: Costa Rican Natural History. Ed. by D. H. JANZEN. Chicago: Univ. Press. Pp. 468–469.
- HERSHKOVITZ, P. (1969): The evolution of mammals on southern continents. VI. The recent mammals of the neotropical region: a zoogeographic and ecological review. Q. Rev. Biol. 44, 1–70.
- HOLMES, A. C. V.; SANDERSON, G. C. (1965): Populations and movements of opossums in east central Illinois. J. Wildl. Manage. 20, 287–295.
- HUSSON, A. M. (1978): The Mammals of Suriname. Leiden: E. J. Brill.
- JULIAN-LAFERRIERE, D.; ATRAMENTOWICZ, M. (1990): Feeding and reproduction of three didelphid marsupials in two neotropical forests (French Guiana). Biotropica 22, 404–415.
- KIRSCH, J. A. W. (1977): The classification of marsupials. In: The biology of marsupials. Ed. by D. HUNSAKER II. New York: Academic Press. Pp. 1–50.
- MAACK, R. (1981): Geografia Fisica do Estado do Paraná. 2nd Ed. Rio de Janeiro: J. Olympio.

- MOTTA, M. F. D.; CARREIRA, J. C. A.; FRANCO, A. M. R. (1983): A note on reproduction of *Didelphis marsupialis* in captivity. Mem. Inst. Oswaldo Cruz 78, 507–509.
- O'CONNELL, M. A. (1989): Population dynamics of neotropical small mammals in seasonal habitats. J. Mammalogy **70**, 532–548.
- RYSER, J. (1992): The mating system and male mating success of the Virginia Opossum (*Didelphis virginiana*) in Florida. J. Zool. (London) **228**, 127–139.
- STOUT, I. J.; SONENSHINE, D. E. (1974): Ecology of an opossum population in Virginia, 1963–69. Acta theriol. 19, 235–245.
- SUNQUIST, M. E.; AUSTAD, S. N.; SUNQUIST, F. (1987): Movements patterns and home range in the common opossum (*Didelphis marsupialis*). J. Mammalogy 68, 173–176.
- SUNQUIST, M. E.; EISENBERG, J. F. (1993): Reproductive strategies of female *Didelphis*. Bull. Florida Mus. Nat. Hist., Biol. Sci. 36, 109–140.
- TYNDALE-BISCOE, C. H.; MACKENZIE, R. B. (1976): Reproduction in Didelphis marsupialis and D. albiventris in Colombia. J. Mammalogy 57, 249–265.
- ZAR, J. H. (1984): Biostatistical Analysis. 2nd Ed. Englewood Cliffs: Prentice-Hall.
- Authors' address: NILTON C. CÁCERES and EMYGDIO L. A. MONTEIRO-FILHO, Departamento de Zoologia, Universidade Federal do Paraná, Cx. P. 19020, Curitiba-PR, 81531-990, Brasil

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Mammalian Biology (früher Zeitschrift für Säugetierkunde)

Jahr/Year: 1998

Band/Volume: 63

Autor(en)/Author(s): Cáceres Nilton C., Monteiro-Filho Emygdio L. A.

Artikel/Article: Population dynamics of the Common opossum, Didelphis marsupialis (Mammalia, Marsupialia), in southern Brazil Population dynamics of the Common opossum, Didelphis marsupialis (Mammalia, Marsupialia), in southern Brazil 169-172