Zitteliana

An International Journal of Palaeontology and Geobiology

Series A/Reihe A Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Geologie





München 2010

Zitteliana

An International Journal of Palaeontology and Geobiology

Series A/Reihe A

Mitteilungen der Bayerischen Staatssammlung für Paläontologie und Geologie

50

CONTENTS/INHALT

BABA SENOWBARI-DARYAN & MICHAELA BERNECKER Amblysiphonella agahensis nov. sp., and Musandamia omanica nov. gen., nov. sp. (Porifera) from the Upper Triassic of Oman	3
ALEXANDER NÜTZEL A review of the Triassic gastropod genus <i>Kittliconcha</i> BONARELLI, 1927 – implications for the phylogeny of Caenogastropoda	9
Andrzej Kaim & Maria Alessandra Conti A problematic zygopleuroid gastropod <i>Acanthostrophia</i> revisited	21
GERNOT ARP Ammonitenfauna und Stratigraphie des Grenzbereichs Jurensismergel/Opalinuston- Formation bei Neumarkt i.d. Opf. (oberstes Toarcium, Fränkische Alb)	25
VOLKER DIETZE Über <i>Ammonites Humphriesianus umbilicus</i> QUENSTEDT, 1886 an seiner Typus-Lokalität (östliche Schwäbische Alb, Südwestdeutschland)	55
Volker Dietze, Günter Schweigert, Gerd Dietl, Wolfgang Auer, Wolfgang Dangelmaier, Roger Furze, Stefan Gräbenstein, Michael Kutz, Elmar Neisser, Erich Schneider & Dietmar Schreiber	
Rare Middle Jurassic ammonites of the families Erycitidae, Otoitidae and Stephanoceratidae from southern Germany	71
WOLFGANG WITT Late Miocene non-marine ostracods from the Lake Küçükçekmece region, Thrace (Turkey)	89
Jérôme Prieto Note on the morphological variability of <i>Keramidomys thaleri</i> (Eomyidae, Mammalia) from Puttenhausen (North Alpine Foreland Basin, Germany)	103
MARTIN PICKFORD Additions to the DEHM collection of Siwalik hominoids, Pakistan: descriptions and interpretations	111
MICHAEL KRINGS, NORA DOTZLER, THOMAS N. TAYLOR & JEAN GALTIER Microfungi from the upper Visean (Mississippian) of central France: Structure and development of the sporocarp <i>Mycocarpon cinctum</i> nov. sp.	127
ZLATKO KVAČEK & JOSEF BOGNER <i>Aracistrobus</i> , an enigmatic non-araceous fossil from the Eurasian Oligocene and Miocene	137
Instructions for authors/Hinweise für Autoren	143

Γ	Zitteliana	A 50	143 Seiten	München, 30.06.2010	ISSN 1612-412X

Editors-in-Chief/Herausgeber: Gert Wörheide, Michael Krings Production and Layout/Bildbearbeitung und Layout: Martine Focke, Manuela Schellenberger Bayerische Staatssammlung für Paläontologie und Geologie

Editorial Board

A. Altenbach, München B.J. Axsmith, Mobile, AL F.T. Fürsich, Erlangen K. Heißig, München H. Kerp, Münster J. Kriwet, Stuttgart J.H. Lipps, Berkeley, CA T. Litt, Bonn A. Nützel, München O.W.M. Rauhut, München B. Reichenbacher, München J.W. Schopf, Los Angeles, CA G. Schweigert, Stuttgart F. Steininger, Eggenburg

Bayerische Staatssammlung für Paläontologie und Geologie Richard-Wagner-Str. 10, D-80333 München, Deutschland http://www.palmuc.de email: zitteliana@lrz.uni-muenchen.de

Für den Inhalt der Arbeiten sind die Autoren allein verantwortlich. Authors are solely responsible for the contents of their articles.

Copyright © 2010 Bayerische Staassammlung für Paläontologie und Geologie, München

Die in der Zitteliana veröffentlichten Arbeiten sind urheberrechtlich geschützt. Nachdruck, Vervielfältigungen auf photomechanischem, elektronischem oder anderem Wege sowie die Anfertigung von Übersetzungen oder die Nutzung in Vorträgen, für Funk und Fernsehen oder im Internet bleiben – auch auszugsweise – vorbehalten und bedürfen der schriftlichen Genehmigung durch die Bayerische Staatssammlung für Paläontologie und Geologie, München.

ISSN 1612-412X

Druck: Gebr. Geiselberger GmbH, Altötting

Cover illustrations: (from left to right) Shell of the gastropod *Loxonema regium* DE KONINCK from the Carboniferous of Belgium (redrawn from DE KONINCK 1881); Solitary coral *Caninia* sp. from the Carboniferous of England (redrawn from RAMSBOTTOM in MCKERROW 1978); Tooth of the rare ruminant *Orygotherium escheri* VON MEYER from the Miocene of Germany (after RÖSSNER & MÖRS 2001). **Back cover:** Atrium of the Munich Palaeontological Museum, view from the main entrance.

Umschlagbilder: (von links nach rechts) Gehäuse der Schnecke *Loxonema regium* DE KONINCK aus dem Karbon von Belgien (neu gezeichnet nach DE KONINCK 1881); Solitärkoralle *Caninia* sp. aus dem Karbon von England (neu gezeichnet nach RAMSBOTTOM in MCKERROW 1978); Zahn des seltenen Wiederkäuers *Orygotherium escheri* von MEYER aus dem Miozän von Deutschland (nach Rössner & Mörs 2001). **Rückseite:** Lichthof des Paläontologischen Museums München, Blick vom Haupteingang.

Zitteliana	A50	9 - 24	1 Pl., 5 Textfigs	München, 30.06.2010	ISSN 1612 - 412X
------------	-----	--------	-------------------	---------------------	------------------

A review of the Triassic gastropod genus *Kittliconcha* BONARELLI, 1927 – implications for the phylogeny of Caenogastropoda

By

Alexander Nützel*

Bayerische Staatssammlung für Paläontologie und Geologie; Department of Earth- and Environmental Sciences, Palaeontology & Geobiology and GeoBioCenter LMU, Ludwig-Maximilians Universität München, Richard-Wagner Str. 10, 80333 München, Germany

Manuscript received March 15, 2010; revised manuscript accepted April 05, 2010

Abstract

Eight Triassic species and one variety have been included in the genus Kittliconcha BONARELLI, 1927. In this composition, Kittliconcha is polyphyletic. The holotypes of Kittliconcha cassiana BONARELLI, 1927 (type species of Kittliconcha) and Settsassia obliquecostata (MÜNSTER, 1841) (type species of Settsassia), as well as the type material of "Loxonema" walmstedti (KLIPSTEIN, 1843) are studied and illustrated (all are from the Cassian Formation). All three species are considered to be synonymous here. The former diagnosis of Kittliconcha encompassed the presence of a varix and reduction of axial ribs on the last whorl. However, these characters resulted from a repaired shell fracture in the holotype of Kittliconcha cassiana. This specimen represents actually Melania obliquecostata MÜNSTER, 1841, which is the type species of Settsassia BANDEL, 1992. Therefore, Settsassia BANDEL, 1992 is a synonym of Kittliconcha BONARELLI, 1927. Kittliconcha alias its junior synonym Settsassia is probably a representative of the cerithimorph stem line as is indicated by its small, smooth larval shell which resembles that of Campaniloidea. A relationship with Littorinimorpha or other basal caenogastropods is also possible. Only a single additional nominate Triassic species may actually represent Kittliconcha: K. obesa from the Middle Triassic of China. Two species from the Late Triassic of South America have been previously assigned to Kittliconcha. They are based on poorly preserved and insufficiently documented material so that any generic assignment is basically speculative. These species are K. dissimilis HAAS, 1953, and K. peruviana HAAS, 1953. K. doelloi BONARELLI, 1927 from Argentina is also based on poorly preserved material which is of unknown age. The material from the Late Triassic Cassian Formation which was assigned to Kittliconcha by ZARDINI (1978) does not belong here. It represents the new sinistral genus Triadosinister and two species (including one new species) which are tentatively assigned to the genus Katosira. These species are Katosira? cortinensis n. sp. and Katosira? giauensis ZARDINI, 1978. The holotype of the type species of the sinistral genus Allostrophia from the Cassian Formation is illustrated. Due to its poor preservation this genus is considered here to represent a nomen dubium. The early Triassic species *Kittliconcha sciaphostera* (BATTEN & STOKES, 1986) also does not represent *Kittliconcha*. It is transferred to the genus *Pseudotritonium* which is assigned to the family Maturifusidae. A lectotype of the type species of *Pseudotritonium* (*P. venustum* from the Cassian Formation) is designated and illustrated. The cancellate ornament, shape, aperture, and the protoconch morphology of *Kittliconcha sciaphostera* suggest that this species is one of the earliest members of an evolutionary line leading to the Mesozoic family Maturifusidae, which could be the stem-group of Neogastropoda.

Key words: Triassic; Gastropoda; Systematics; New Taxa; Maturifusidae; Cassian Formation.

Zusammenfassung

Der Gattung Kittliconcha BONARELLI, 1927 wurden bislang acht triassische Gastropodenarten zugeordnet. In dieser bisherigen Zusammensetzung ist Kittliconcha polyphyletisch. Die Holotypen von Kittliconcha cassiana BONARELLI, 1927 (Typusart von Kittliconcha) und von Settsassia obliquecostata (MÜNSTER, 1841) (Typusart von Settsassia) sowie das Typusmaterial von "Loxonema" walmstedti (KLIPSTEIN, 1843) werden nachuntersucht und abgebildet (alle stammen aus der Cassian Formation). Sie drei Artnamen sind Synonyme. Die bisherige Diagnose von Kittliconcha beinhaltete das Vorhandensein einer Varix und die Reduktion von Axialrippen auf der letzten Windung. Diese Merkmale sind jedoch das Resultat einer reparierten Schalenfraktur beim Holotyp von Kittliconcha cassiana. Dieses Exemplar gehört der Art Melania obliquecostata MÜNSTER, 1841 an, die Typusart von Settsassia BANDEL, 1992 ist. Also ist Settsassia BANDEL, 1992 ein Synonym von Kittliconcha BONARELLI, 1927. Kittliconcha alias ihrem jüngeren Synonym Settsassia ist vermutlich ein Vertreter der Stammlinie der Cerithimorpha, worauf die kleine, glatte, heliciforme Larvalschale hinweist, die in ähnlicher Weise bei den Campaniloidea vorkommt. Eine engere verwandtschaftliche

^{*}E-mail: a.nuetzel@lrz.uni-muenchen.de.

Beziehung ist auch zu den Littorinimorpha oder anderen basalen Caenogastropoden möglich. Nur eine weitere nominelle triassische Art steht vermutlich zu Recht in Kittliconcha: K. obesa aus der mittleren Trias von China. Drei Arten aus der oberen Trias von Südamerika wurden der Gattung Kittliconcha zugeordnet. Sie basieren auf schlecht erhaltenem und unzureichend dokumentiertem Material, so dass jegliche Gattungszuweisung spekulativ erscheint. Es handelt sich dabei um die Arten K. dissimilis HAAS, 1953 und K. peruviana HAAS, 1953. K. doelloi BONARELLI, 1927 aus Argentinien basiert ebenfalls auf ungenügend bekanntem Material und ist überdies unsicheren Alters. Das Material aus der obertriassischen Cassian Formation, das ZARDINI (1978) Kittliconcha zuordnete, gehört dieser Gattung nicht an. Es repräsentiert die neue sinistrale Gattung Triadosinister und zwei Arten (davon eine neue), die vorläufig der Gattung Katosira zugerechnet werden. Es sind dies die Arten Katosira? cortinensis n. sp. und Katosira? giauensis ZAR-DINI, 1978. Der Holotyp der Typusart der sinistralen Gattung Allostrophia aus der Cassian Formation wird abgebildet. Das Stück ist so schlecht erhalten, dass die Gattung als Nomen dubium anzusehen ist. Die untertriassische Art Kittliconcha sciaphostera gehört ebenfalls nicht der Gattung Kittliconcha an. Sie wird der Gattung Pseudotritonium zugeordnet und diese der Familie Maturifusidae. Für die Typusart von Pseudotritonium (P. venustum aus der Cassian Formation) wird ein Lectotyp festgelegt und abgebildet. Das gitterartige Ornament, die Gestalt, Mündung und die Protoconchmorphologie legen nahe, dass es sich bei Kittliconcha sciaphostera um einen der frühesten Vertreter der evolutionären Linie handelt, die zur mesozoischen Familie Maturifusidae führt, die ihrerseits die Stammgruppe der Neogastropoda sein könnte.

Schlüsselwörter: Trias; Gastropoda; Systematik; Neue Taxa; Maturifusidae; Cassian Formation.

1. Introduction

The subclass Caenogastropoda is richly diversified in the Triassic and the famous fauna from the Cassian Formation greatly contributes to this diversity. There are probably as many as 200 described Triassic caenogastropod genera. Many of them are well characterized whereas others are obscure because they are based on poorly preserved or insufficiently documented material and many of them have never been revised since their original publication. One of the more obscure genera is *Kittliconcha* BONARELLI, 1927 which has a type species from the Cassian Formation. It was originally erected as a subgenus of *Zygopleura* KOKEN, 1892 but later promoted to genus level (KNIGHT et al. 1960; NÜTZEL 1998). The restudy of the type specimen in question shows that at least two diagnostic features, a special type of varix-like axial rib and the reduction of the ornament on the last part of the last whorl, are results of a healed shell fracture. The type species is here re-studied and a new diagnosis is given. Based on this, the species which have been placed in *Kittliconcha* are re-evaluated.

Seven Triassic species and one variety as well as one species of uncertain age have to date been assigned to the genus *Kittliconcha*:

- K. cassiana BONARELLI, 1927, Cassian Formation, Carnian
- K. contraria ZARDINI, 1978, Cassian Formation, Carnian
- K. dissimilis HAAS, 1953, Pucara Formation, Peru, Late Triassic
- K. doelloi BONARELLI, 1927 (age unclear, Argentina)
- Zygopleura (K.) cassiana n. f. fascicostata ZARDINI, 1978, Cassian Formation, Carnian
- K. giauensis ZARDINI, 1978, Cassian Formation, Carnian,
- K.? obesa Yu, PAN & WANG, 1974, Qingyan Fm., S China
- K. peruviana HAAS, 1953, Pucara Formation, Peru, Late Triassic
- K. sciaphostera BATTEN & STOKES, 1986, Moenkopi Formation, Utah

These species are discussed below, showing that *Kittliconcha* in this composition is clearly polyphyletic.

2. Repositories

Geologische Bundesanstalt Wien; BSPG: Bayerische Staatssammlung für Geologie und Paläontologie München; MPRZ: Museo Paleontologico Rinaldo Zardini, Cortina d'Ampezzo; NHMW: Naturhistorisches Museum Wien; NHM: Natural History Museum London.; AMNH: American Museum of Natural History, New York.

Plate 1:

Kittlicon	icha (obli	quecostata	(Bron	v in	Münster	, 1841),	Late	Tr	iassic	Cassian	Fo	ormation.
-----------	--------	------	------------	-------	------	---------	----------	------	----	--------	---------	----	-----------

- Fig. 1: Holotype of *Kittliconcha cassiana* BONARELLI, 1927 (synonym of *Kittliconcha obliquecostata*), KITTL's (1894: pl. 8, fig. 5) original; Geologische Bundesanstalt, Wien.
- Fig. 1a-d: Lateral views at different angles, height of specimen 18.4 mm; repaired shell fracture on last whorl; axial ribs reduced on repaired shell after fracture.
- Fig. 1e: Oblique lateral view to show varix-like, irregular rib on last whorl; height 10 mm.

Fig. 6: *Kittliconcha obliquecostata*; juvenile specimen showing smooth protoconch and smooth early teleoconch; NHMW 1997/0040/0089; 6a: height 0.9 mm; from NUTZEL (1998: pl. 27Q); 6b: small, smooth protoconch of about 2.5 whorls; larval shell terminating at sinusigera; height 0.34 mm; from NUTZEL (1998: pl. 27R).

Fig. 2: Lectotype of *Melania obliquecostata* selected from MÜNSTER's type collection, type species of *Kittliconcha* and its junior synonym *Settsassia*; 17 mm high, BSPG AS VII 506.

Fig. 3: *Turritella walmstedti* Klipstein from the type series of Klipstein, Natural History Museum, London (NHM 35692C); this taxon represents a junior synonym of *Kittliconcha obliquecostata*; height 13.4 mm.

Fig. 4: Kittliconcha obliquecostata; height 6 mm; from NÜTZEL (1998: pl. 27N), NHMW 1997/0040/0087.

Fig. 5: *Kittliconcha obliquecostata*; juvenile specimen showing smooth protoconch, smooth early teleoconch, and transition to teleoconch with axial ribs; height 4.4 mm; from NÜTZEL (1998, pl. 27P), NHMW 1997/0040/0088.



3. The genus Kittliconcha

Genus Kittliconcha BONARELLI, 1927

Type species: Zygopleura (Kittliconcha) cassiana BO-NARELLI, 1927 pro Loxonema walmstedti (KLIPSTEIN, 1843) sensu KITTL (1894: pl. 8, fig. 5) = "Gehäuse mit abnormer (?) Schlußwindung" ("Shell with abnormal (?) final whorl"); early Carnian, Cassian Formation.

Other species originally included: *Kittliconcha doelloi* BONARELLI, 1927, Argentina (age unknown).

Remarks: BONARELLI (1927) designated Zygopleura (Kittliconcha) cassiana BONARELLI, 1927 as the type species. He based this species on a specimen which was illustrated by KITTL (1894: pl. 8, fig. 5) under the name Loxonema walmstedti (KLIPSTEIN, 1843). KITTL noted that this specimen has an "abnormal (?) final whorl". When BONARELLI (1927) erected the type species Zygopleura (Kittliconcha) cassiana, he just referred to this specimen which therefore is the holotype. This type specimen is housed in the collection of the Geologische Bundesanstalt in Vienna (Austria). Obviously, BONARELLI never studied this specimen or topotypical material from the Cassian Formation. The holotype is re-described and illustrated below. BONARELLI (1927) gave no formal diagnosis for Kittliconcha but his description and comparisons to other taxa warrants the validity of the new subgenus. However, BONARELLI (1927) had no material from the Cassian Formation at hand and his interpretation of KITTL's (1894) illustration is obviously flawed, therefore, his taxonomical considerations are largely irrelevant. A short diagnosis for Kittliconcha was given by KNIGHT et al. (1960): "Broader and less acute than in most genera of the family [Zygopleuridae]; whorls strongly convex; collabral riblets strong on spire whorls, where only their opisthocyrt upper part is exposed, obsolete on last whorl except for a varix; no spiral ornament." In the following the holotypes of the type species of Kittliconcha and Settsassia are re-described and illustrated. It becomes clear that both taxa are synonyms and consequently Settsassia Bandel, 1992 is a junior synonym of *Kittliconcha* BONARELLI, 1927.

3.1 Re-description of the holotype of *Kittliconcha cassiana*

Kittliconcha cassiana BONARELLI, 1927 Pl. 1, Fig. 1

1894 *Loxonema walmstedti* (KLIPSTEIN, 1843) – KITTL: 169, pl. 8, fig. 5.

Material: The holotype of Zygopleura (Kittliconcha) cassiana, KITTL's original specimen (1894, pl. 8, fig. 5) from the Stuores Meadows, Cassian Formation, N Italy, Early Carnian; housed in the Geologische Bundesanstalt Wien.

Description: Shell high-spired; about 8.5 whorls, 18.4 mm high, 7.4 mm wide; apical angle 26°; whorls convex, ornamented with 12 to 14 axial ribs per whorl; ribs round, orthocline to slightly opisthocyrt, little narrower than space between ribs; ribs do not continue onto base; base smooth, convex, without ledge when joining whorl face; suture incised; last whorl with pronounced deep, healed shell fracture; shell after fracture without regular ribs, but with one irregular varix and a varix-like aperture.

Remarks: It is clear that the presence of a varix and the reduction of the ribs on the last whorl which have been considered diagnostic for *Kittliconcha* by KNIGHT et al. (1960) resulted from a healed shell fracture. The animal survived a rather deep shell fracture but was unable to build a shell with regular axial ribs afterwards. Shell fractures are common in the Cassian gastropods as can be seen in ZARDINI's (1978) monograph and as was discussed by VERMEIJ et al. (1982). KITTL (1894) interpreted this varix as a not resorbed apertural varix but this is obviously not the case.

3.2 Re-description of *Kittliconcha obliquecostata* including designation of lectotype

Kittliconcha obliquecostata (BRONN in MÜNSTER, 1841) Pl. 1, Figs 2–6

- * 1841 Melania obliquecostata n. sp. BRONN in MÜNSTER: 97, pl.
 9, fig. 43.
 - 1843 Turritella walmstedti n. sp. KLIPSTEIN: 179, pl. 11, figs 28, 29.
 - 1843 Melania haueri n. sp. KLIPSTEIN: 190, pl. 12, fig. 30.
 - 1978 Zygopleura walmstedti (Klipstein, 1843) Zardini: 40, pl. 24, figs 13–16.
 - 1978 Zygopleura obliquecostata (BRONN in MÜNSTER, 1841) ZARDINI: 40, pl. 24, figs 17–22.
 - 1978 Zygopleura haueri (KLIPSTEIN, 1843) ZARDINI: 40, pl. 25, figs 8, 9.
 - 1992 Settsassia obliquecostata (BRONN in MÜNSTER, 1841) BAN-DEL: 64, pl. 11, figs 3, 4, non fig. 7, pl. 12, figs 1, 2.
 - 1993 Settsassia obliquecostata (BRONN in Münster, 1841) BANDEL: 13, pl. 4, fig. 2.
 - 1994 Settsassia obliquecostata (Bronn in Münster, 1841) BANDEL: 145, pl. 3, fig. 1.
 - 1998 Settsassia obliquecostata (Bronn in Münster, 1841) NÜTZEL: 167, pl. 27 figs L– R.

For additional synonyms, refer to DIENER (1926) and KU-TASSY (1940).

Lectotype: BSPG AS VII 506, from the type series, specimen illustrated by Bronn in Münster (1841: pl. 9, fig. 43).

Description: Shell high-spired, conical, straight-sided; lectotype comprising 10 teleoconch whorls (apical whorls missing), 17 mm high, 7.1 mm wide; whorls regularly expanding, evenly convex; suture distinct; first two to three teleoconch whorls smooth; later whorl with 12 to 15 axial ribs per whorl; ribs wave-like, round, approximately as wide as interspaces between them; ribs almost straight, orthocline, slightly opisthocyrt or somewhat prosocline; ribs of about equal strength over the whorl but submedian somewhat stronger in some



Textfigure 1: *Triadosinister contrarius* (Zardini, 1978), Late Triassic Cassian Formation. (1.1) Paralectotype, height 17.5 mm; from the location Rumerlo, from ZARDINI (1978: pl. 26, fig. 5), MPRZ 1492. (1.2) lectotype height 21 mm; from the location Alpe di Specie, from ZARDINI (1978: pl. 26, fig. 6), MPRZ 1493. (1.3) height 21 mm; from the location Misurina Skilift, BSPG 2010 III 1; 1.3a, b: height 6 mm; 1.3c: apex detail, height 2.3 mm; 1.3d, e: last whorl detail, height 3 mm.

specimens; ribs terminate somewhat above the suture; they do not continue onto base; base flatly convex, smooth; last whorl of lectotype with very faint, obscure spiral lineation which is absent in earlier whorls; protoconch smooth, heliciform, consisting of 2.5 to 3 whorls, up to 0.34 mm high and 0.25 mm wide; diameter of initial whorl 0.10 to 0.13 mm; protoconch whorls convex, terminating at a distinct sinusigera.

Remarks: As outlined above, *Kittliconcha cassiana* represents a synonym of "*Melania*" *obliquecostata*. Exhaustive descriptions of *Kittliconcha obliquecostata* are available from KITTL (1894) and BANDEL (1992). The small smooth heliciform larval shell of the planktotrophic type and the smooth early teleoconch separate it from similar, convergent species of the Zygopleuridae (BANDEL 1992; NÜTZEL 1998). BANDEL (1992) reported that the last whorl is detached and becomes phane-

romphalous. This could not be observed in the present material. KITTL (1994) noted that a faint spiral striation can be present as is the case in the lectotype figured here (Pl. 1, Fig. 2). KITTL (1894) and BANDEL (1992) found that there are morphological transitions to *Turritella walmstedti* KLIPSTEIN, 1843 and *Melania haueri* KLIPSTEIN, 1843 from the Cassian Formation. The three taxa differ in minor details of the axial ornament and in the spiral angle. These differences can be interpreted as intraspecific variation. Thus, NUTZEL (1998) treated both KLIPSTEIN taxa as synonyms of *Kittliconcha obliquecostata*. For instance, in Plate 1, Fig. 3, a specimen of *Turritella walmstedti* from KLIPSTEIN's type collection (Natural History Museum London) is illustrated. This teleoconch fragment shows no differences to *M. obliquecostata* and its synonym *K. cassiana*.

Kittliconcha obliquecostata is abundant at the Stuores Meadows localities (Prati di Stuores). ZARDINI (1978) reported this species also from the area near Cortina d'Ampezzo, especially at the locations Costalaresc and Alpe di Specie.

3.3 Emended diagnosis for *Kittliconcha*, largely based on the diagnoses of *Settsassia* as given by BANDEL (1992) and NÜTZEL (1998)

Shell high-spired, conical, straight-sided, up to about 2 cm high; apical angle 25° to somewhat more than 30°; whorls regularly expanding, evenly convex; first two to three teleoconch whorls smooth; later whorl with axial ribs; ribs broad, round; almost straight, orthocline to slightly opisthocyrt or somewhat prosocline; base flatly convex, smooth; spiral ornament basically absent but an obscure spiral lineation may be present; protoconch small, smooth, heliciform, terminating at a distinct sinusigera.

3.4 Discussion of the systematic placement of *Kittliconcha*

It is now clear that the main diagnostic features of Kittliconcha as indicated by KNIGHT et al. (1960), rib reduction and varix on the last whorl, are results of a repaired shell fracture. This raises the question about the validity of the genus Kittliconcha and K. cassiana. A comparison of the holotypes of K. cassiana and Melania obliquecostata BRONN in MÜNSTER, 1841 from the Cassian Formation (see Pl. 1, Figs 1 and 2) shows that both specimens are very similar and are probably conspecific. Therefore, K. cassiana is a junior subjective synonym of Melania obliquecostata. Melania obliquecostata and its synonyms have long been attributed to the genus Zygopleura KOKEN, 1892. However, BANDEL (1992, 1993) reported a small smooth, heliciform larval shell which is untypical for Zygopleura (see also NÜTZEL 1998). Therefore, BANDEL (1992) introduced the genus Settsassia with the type species Melania obliquecostata. Thus, Kittliconcha and Settsassia have the same type species and therefore, Settsassia BANDEL, 1992 is a junior synonym of Kitt*liconcha* BONARELLI, 1927. BANDEL (1992) also introduced the family Settsassiidae and this name remains valid according to the ICZN. Therefore, the following arrangement is suggested:

Family Settsassiidae BANDEL, 1992

Type genus: Settsassia BANDEL, 1992, a junior synonym of Kittliconcha BONARELLI, 1927.

Genus: Kittliconcha BONARELLI, 1927

Type species: *Melania obliquecostata* BRONN in MÜN-STER, 1841 (synonyms *Turritella walmstedti* KLIPSTEIN, 1843, *Melania haueri* KLIPSTEIN, 1843, and *Zygopleura* (*Kittliconcha*) *cassiana* BONARELLI, 1927.

Kittliconcha obliquecostata is certainly a caenogastropod and is probably related to the Cerithioidea (BANDEL 1992, 1993) or to the Campaniloidea. A close relationship to Littorinimorpha is also possible. The small, smooth heliciform larval shell of *Kittliconcha* strongly resembles that of the Mesozoic basal cerithimorph genera *Campanile*, *Metacerithium*, and *Dzikella* (the latter is probably a synonym of *Metacerithium*) (see KIEL et al. 2000; KAIM 2004; KIEL 2006). The larval shells of the mentioned campanilids have a sinusigera which is heavily strengthened by a varix. However, this is not the case in *Kittliconcha*. Nevertheless, it seems to be possible that Campanilidae and Settsassiidae are closely related to each other. NUTZEL (1998) supposed that Settsassiidae are descendants of the Palaeozoic Palaeostyloidea which encompass high-spired, axially ribbed gastropods with smooth or poorly ornamented larval shells.

4. Re-assignment and discussion of other species that have been included in *Kittliconcha*

As mentioned above, besides Kittliconcha obliquecostata alias K. cassiana, seven other species and one variety have been assigned to the genus Kittliconcha: two species and one variety from the Early Carnian Cassian Formation (K. contraria ZARDINI, 1978, K. giauensis ZARDINI, 1978, and K. cassiana n. f. fascicostata ZARDINI, 1978), two species from the Late Triassic of the Pucara Formation, Peru (K. dissimilis HAAS, 1953 and K. peruviana HAAS, 1953), K. doelloi BONARELLI, 1927 from Argentina (age uncertain), K.? obesa YU, PAN & WANG, 1974 from the Middle Triassic of South China, and K. sciaphostera BATTEN & STOKES, 1986 from the Early Triassic Moenkopi Formation of Utah, USA. The new information on the type species of Kittliconcha suggests that most of aforementioned species do not belong to this genus. These species share a zygopleuroid habitus i. e., they are more or less high-spired with a teleoconch ornament of axial ribs. Most are probably caenogastropods, however, the protoconch is known only from K. sciaphostera and thus, the higher systematic placement of the other species is tentative. Many of the taxa assigned to Kittliconcha have an additional ornament of distinct spiral lirae or a spiral micro-ornament. Such features have not been observed in the type species of Kittliconcha (see above and Pl. 1). Kittliconcha obliquecostata may show only a very faint spiral lineation which is obviously result of intraspecific variation and cannot be considered diagnostic. BONARELLI (1927: 60) excluded a spiral ornament in his discussion of Kittliconcha: "... el ejemplar de referencia no presenta estrías espirales y la falta absoluta de este tipo de ornamentación acompañando a las cóstulas, permite excluir del todo su pertenencia al género Katosira" (translation: "... the specimen in question [holotype of Kittliconcha cassiana] shows no spiral striae and the complete absence of such an ornament in connection with the ribs facilitate to exclude an assignment to the genus Katosira entirely.").

In the following, the species which have been assigned to *Kittliconcha* are discussed and re-assigned whenever possible.

4.1 Re-assignment of *Kittliconcha contraria* ZARDINI, 1978

Kittliconcha contraria does not belong to *Kittliconcha* and it is designated here as a type species of the new genus *Triadosinister*.

Triadosinister n. gen.

Etymology: For the Triassic period and sinistrality.

Diagnosis: High-spired sinistral shells with prominent axial ribs and a reticulate micro-ornament of fine spiral grooves and threads intersected by strengthened growth lines; whorls convex, pendant; relatively few axial ribs per whorl; axial ribs not reaching the adapical suture; aperture seemingly with wide siphonal opening.

Type species: *Kittliconcha contraria* ZARDINI, 1978 (Carnian, Cassian Formation).

Remarks: Triadosinister differs from Kittliconcha in being left coiled and in having a reticulate micro-ornament. Allostrophia KITTL, 1894 with the type species Allostrophia perversa (MÜNSTER, 1841) from the Cassian Formation is another high spired sinistral genus with axial ribs on the teleoconch whorls. MÜNSTER's type specimen (BSPG AS VII 1681) is so poorly preserved (encrusted and abraded) (Textfig. 2) that the species and consequently also the genus are considered here to be Nomina dubia. According to KITTL (1894) and KNIGHT et al. (1960), Allostrophia has less convex whorls, continuous axial ribs and lacks a reticulate micro-ornament. The relatively well-preserved specimens illustrated as A. perversa by ZARDI-NI (1978: pl. 26, figs 2-4) show straight axial ribs, the whorls are not pendant (i.e., they do not have their maximum width below mid-whorl) and obviously lack a micro-ornament. A siphonal notch which is present in Triadosinister is obviously absent in Allostrophia (MARSHALL 1990). The protoconchs of Allostrophia and Triadosinister are unknown so far. BANDEL (1994: 138) mentioned that he observed that the protoconch of Allostrophia it is not typical for Zygopleuridae. The Early Jurassic high-spired sinistral genus Virgella DEGREGORIO, 1930 has numerous fine axial ribs.

Triadosinister contrarius (ZARDINI, 1978) Textfig. 1

* 1978 *Kittliconcha contraria* n. sp. – ZARDINI: 42, pl. 26, figs 5, 6 (?fig. 7).

Lectotype: Here designated, specimen illustrated in ZAR-DINI (1978: pl. 26, fig. 6a, b, MPRZ 1493) Cassian Formation, Early Carnian, from Alpe di Specie, Seelandalpe; ZARDINI's illustration is reproduced here (Textfig. 1.2).

Paralectotype: The specimen illustrated in ZARDINI (1978: pl. 26 fig. 5a, b, MPRZ 1492) from Rumerlo; ZARDINI's illustration is reproduced here (Textfig. 1.1).

Additional material: One specimen from the Misurina Skilift location (see NÜTZEL et al. 2010), BSPG 2010 III 1 (Textfig. 1.3).

Description: Shell high-spired, slender, sinistral; lectotype is a teleoconch fragment of about five whorls, 21 mm high, 8 mm wide; whorls distinctly convex; whorl profile somewhat pendant; whorls somewhat adpressed; sutures distinct; whorls



Textfigure 2: The holotype of *Turritella perversa* MÜNSTER, 1841, type species of *Allostrophia* KITTL, 1894; the preservation of the type specimen (illustrated in MÜNSTER 1841: pl. 9, fig. 41) is so poor that this taxon is considered to represent a nomen dubium; BSPG AS VII 1681; 2.1a: 16.1 mm high, 5.9 mm wide; 2.1b: detail of earliest preserved whorls showing faint, eroded axial ribs; 8.8 mm high.

ornamented with prominent axial ribs; number of ribs increasing during ontogeny from 8–9 to about 15 per whorl; ribs reduced on last whorl; ribs slightly opisthocline somewhat curving opisthocyrt; ribs much narrower than interspaces between ribs; ribs weak or reduced below adapical suture and absent on base; whorls with reticulate micro-ornament of fine spiral grooves and threads intersected by strengthened parasigmoidal growth lines; growth lines especially strong below adapical suture; aperture seemingly with wide siphonal opening; base flatly convex, covered with reticulate microornament; protoconch unknowon.

Remarks: The differences to other species and genera are discussed above. ZARDINI (1978: pl. 26, figs, 5-7) illustrated three specimens which he assigned to Kittliconcha contraria. The specimen in ZARDINI (1978: pl. 26, fig. 7) is poorly preserved and its identity is questionable. The shell figured here in Textfigure 1.3 is the first known from the location Misurina Skilift near Cortina d'Ampezzo. It shows a well-preserved micro-ornament. The three specimens of Triadosinister contrarius illustrated here seem to exhibit a considerable intraspecific variability in number and strength of the ribs. However, these specimens represent different growth stages and the pronounced ontogenetic change of the ornament of T. contrarius (increase of the number of ribs per whorl and reduction of ribs on mature whorls) is suggestive of this apparent variability. However, there is no doubt that these specimens represent a single species.

4.2 Re-assignment and discussion of Kittliconcha cassiana sensu ZARDINI (1978) and Kittliconcha giauensis

ZARDINI (1978: pl. 28, figs 3–4) illustrated two specimens from the early Carnian Cassian Formation and assigned them to *Kittliconcha cassiana* BONARELLI, 1927. These specimens represent a new species and probably a new genus. This new species is described herein and tentatively assigned to *Katosira*. *Kittliconcha giauensis* ZARDINI, 1978 also does not belong to *Kittliconcha* and is tentatively transferred to *Katosira* herein.

Genus Katosira Koken, 1892

Type species: *Katosira periniana* (ORBIGNY, 1851), Early Jurassic, France.

Katosira? cortinensis n. sp. Textfig. 3.1, 3.2

*1978 Zygopleura (*Kittliconcha*) cassiana BONARELLI, 1927 – ZAR-DINI: 42, pl. 28, figs 3, 4.

?1978 Zygopleura (Kittliconcha) cassiana n. f. fascicostata – ZARDINI: 42, pl. 28, fig. 2.

Diagnosis: Shell broad, relatively low-spired, last whorl somewhat higher than spire; whorls convex, somewhat adpressed; whorls ornamented with narrow, sharp, fold like axial ribs; axial ribs curving, opisthocyrt to opisthocline; early teleoconch whorls with few ribs, last whorl with numerous ribs; whorls with additional spiral micro-ornament; base convex.

Holotype: The specimen illustrated by ZARDINI (1978) in pl. 28, fig. 3a, b, reproduced here in Textfigure 3.1a, b, MPRZ 3706.

Paratype: The specimen illustrated by ZARDINI (1978) in pl. 28, fig. 4a, b, reproduced here in Textfigure 3.2a, b, MPRZ 3707.

Etymology: Atte Cortina d'Ampezzo (type locality)

Locus typicus: The holotype comes from Alpe di Specie



Textfigure 3: Reproduction of ZARDINI's (1978) illustrations of specimens which have been assigned to *Kittliconcha*; Late Triassic Cassian Formation. (3.1, 3.2): *Katosira? cortinensis* n. sp. (3.1): height 13 mm; from the location Alpe di Specie, from ZARDINI (1978: pl. 28, fig. 3a, b), MPRZ 3706. (3.2): height 9 mm; from the location Costalaresc, from ZARDINI (1978: pl. 28, fig. 4a, b), MPRZ 3707. (3.3): "*Zygopleura (Kittliconcha) cassiana* n. f. *fascicostata*", height 9.6 mm; from the location Sasso di Stria, from ZARDINI (1978: pl. 28, fig. 2), MPRZ 3705. (3.4): *Katosira? giauensis* (ZARDINI, 1978), height 9.8 mm; from the location Forcella Giau, from ZARDINI (1978: pl. 28, fig. 6a, b), MPRZ 3709.

near Cortina d'Ampezzo (South Tyrol, N Italy).

Stratum typicum: Early Carnian, Cassian Formation.

Description: Shell broad, relatively low-spired, last whorl somewhat higher than spire; holotype comprises about three whorls (apex missing), 13 mm high, 7.2 mm wide; whorls convex, somewhat adpressed; whorls ornamented with narrow, sharp, fold like axial ribs; axial ribs curving, opisthocyrt to opisthocline; early teleoconch whorls with about 9 ribs per whorl, last whorl of holotype with more than 15 ribs; distance between ribs much wider than width of ribs, especially in early whorls; ribs fade on base; whorls ornamented with additional spiral micro-ornament; base convex, not demarcated from whorl face, minutely phaneromphalous.

Remarks: The broad, stout shape, the sharp ribs and the spiral micro-ornament described by ZARDINI (1978) strongly suggest that these specimens do not belong to Kittliconcha obliquecostata alias K. cassiana. The generic assignment to Katosira is tentative until more information about the new species become available. Katosira encompasses generally more high-spired and slender species with axial ribs crossed by finer spiral threads. The genus is based on the K. periniana from the Early Jurassic of France. This species is based on a poorly preserved specimen (FISCHER & WEBER 1997). The variety Zygopleura (Kittliconcha) cassiana n. f. fascicostata described by ZARDINI (1978) could also belong to Katosira? cortinensis n. sp. but the single teleoconch fragment illustrated by ZARDINI (1978: pl, 28, fig. 2) differs in shape and ornament (Textfig. 3.3). It does not provide enough information for a meaningful systematic placement. Kittliconcha giauensis ZARDINI, 1978 from the Cassian Formation has broad, round axial ribs crossed by several distinct spiral ribs. The holotype, a fragment of two teleoconch whorls, is the only known shell (re-illustrated here in Textfig. 3.4). The strong spiral ribs suggest that it does not belong to Kittliconcha. It is tentatively transferred to Katosira but could also represent a cerithioid.

4.3 Discussion of *Kittliconcha obesa* from the Middle Triassic of South China

Kittliconcha obesa YU, PAN & WANG, 1974 from the Middle Triassic Qingyan Formation of South China as described and illustrated by YIN HONG-FU & YOCHELSON (1983) seems to be similar to the type species of *Kittliconcha*. It is a relatively broad shell with prominent axial ribs. It lacks any spiral ornament. As in *K. obliquecostata*, the earliest teleoconch whorls of *K. obesa* are smooth. The protoconch of *K. obesa* is unknown. Nevertheless, *K. obesa* can be assigned to *Kittliconcha* according to the current state of knowledge.

4.4 Discussion of *Kittliconcha doelloi*, *K. peruviana*, and *K. dissimilis* from the Upper Triassic of South America

Kittliconcha doelloi BONARELLI, 1927 is based on a single specimen from the Pertrolifera Formation, Carbajal, Argen-

tina. This material is probably not of Triassic age but much younger (Mariel Ferrari, written communication 2010). This teleoconch fragment is poorly preserved and poorly illustrated. It is considered here as nomen dubium until the holotype is re-studied and topotypical material is found and described.

Kittliconcha dissimilis HAAS, 1953 and *K. peruviana* HAAS, 1953, both from the Late Triassic Pucara Formation, Peru are also problematic. Judging from HAAS's illustrations of these species, the type material is poorly preserved. The zygopleuroid shells are relatively stout and do resemble *Kittliconcha*. However, HAAS mentioned a spiral striation of the whorls which does not become obvious from the illustrations. This character would not suggest a placement in *Kittliconcha*. BANDEL (1994: 138) mentioned that *Kittliconcha dissimilis* and *K. peruviana* have a small and simple protoconch. I suggest leaving both species tentatively in *Kittliconcha* until more information becomes available.

4.5 Re-assignment of *Kittliconcha sciaphostera* from the Lower Triassic of Utah

Kittliconcha sciaphostera BATTEN & STOKES, 1986 (Early Triassic, Utah) does not belong to *Kittliconcha* and it is included here in the genus *Pseudotritonium* WENZ, 1940.

Family Maturifusidae Gründel, 2001 (= Pseudotritoniinae Golikov & Starobogatov, 1987?)

Remarks: BANDEL (1994: 139) and BOUCHET et al. (2005) assigned *Pseudotritonium* (replacement name for *Palaeotriton* KITTL, 1894) to the family Purpurinidae. GOLIKOV & STARO-BOGATOV (1987) erected the subfamily Pseudotritoniiae based on this genus. Here, *Pseudotritonium* is preliminarily placed in the Maturifusidae. If *Pseudotritonium* and *Maturifusus* are really confamilial, then Maturifusidae GRÜNDEL would fall into the synonymy of Pseudotritoniidae. However, synonymization should await a better knowledge of the type species of *Pseudotritonium*, *Scalaria venusta* from the Late Triassic Cassian Formation. So far, the protoconch of this species is insufficiently known (see below) and it would be important to document it better in order to clarify the relationship between *Pseudotritonium* and *Maturifusus*.

Genus *Pseudotritonium* WENZ, 1940 (replacement name for *Palaeotriton* KITTL, 1894)

Type species: *Scalaria venusta* MÜNSTER, 1841, Early Carnian, Cassian Formation; designation by Wenz (1940) (see Textfig. 4).

A lectotype for the type species *Scalaria venusta* MÜNSTER, 1841 from the Cassian Formation is selected here (BSPG AS VII 1871) (Textfig. 4). It is the specimen which was illustrated by MÜNSTER (1841: pl. 10, fig. 28) and KITTL (1894: pl. 11, fig. 3). *Pseudotritonium venustum* has a fusiform shell shape and a wide aperture. The aperture has a straight inner and a D-shaped outer lip. It seems to be siphonostomous. The teleoconch ornament is cancellate with oblique, opisthocline axial ribs which are crossed by numerous finer spiral lirae. The intersections of spiral lirae and axial ribs are not nodular at intersections





which separates it from species of *Maturifusus* or *Astandes*. The Early Triassic *Pseudotritonium sciaphosterum* (BATTEN & STOKES), 1986 resembles *P. venustum* in general shape and in the cancellate teleoconch ornament.

Pseudotritonium sciaphosterum (BATTEN & STOKES, 1986) Textfig. 5

* 1986 *Kittliconcha sciaphostera* n. sp. BATTEN & STOKES: 19, figs 26–28.

Material: A paratype still fixed on a SEM-stub (AMNH 46314) (Textfig. 5.1) and a topotypical specimen with well preserved protoconch (BSPG 2010 IV 1) (Textfig. 5.2).

Description: *Pseudotritonium sciaphosterum* is a highspired gastropod with an ornament of strong axial ribs crossed by finer spiral threads. The axial ribs are oblique, especially near the adapical suture; on later whorls, they may be almost orthocline to slightly opisthocyrt.

BATTEN & STOKES (1986) described the aperture as holostomous. However, the specimen from the type collection figured here (Textfig. 5.1b, c) shows an elongated aperture which seemingly has a siphonal notch. The protoconch was described as smooth and evenly inflated and is illustrated in BATTEN & STOKES (1986: fig. 28). The topotypical juvenile specimen illustrated here (Textfig. 5.2) is better preserved and shows the protoconch in detail. It consists of almost three whorls and is 0.44 mm high and 0.27 mm wide; the initial whorl has a diameter of 0.15 mm. The larval whorls are convex and somewhat adpressed so that a spiral band is present in a subsutural position. The larval shell is densely ornamented with faint, oblique threads.

Remarks: BATTEN & STOKES (1986) correctly pointed out that the teleoconch ornament of Pseudotritonium sciaphosterum closely resembles that of the Late Palaeozoic genus Plocezyga und Gamizyga (Pseudozygopleuridae) (see HOARE & STURGEON 1980; NÜTZEL 1998). They even synonymised these genera with Kittliconcha. However, Plocezyga and Gamizyga have a larval shell with strong parasigmoidal, collabral axial ribs which are typical for Pseudozygopleuridae. In contrast, Kittliconcha has a smooth larval shell (Pl. 1) and Pseudotritonium sciaphosterum has a non-collabral micro-ornament of oblique threads (Textfig. 5.2). The reticulate teleoconch ornament and the overall shape of Pseudotritonium sciaphosterum resemble that of the Jurassic/Cretaceous family Maturifusidae (e.g., KAIM 2004). The protoconchs of most Maturifusus species are largely smooth but have angulations accompanied by spiral ribs which gives them a step-like whorl profile. However, A. Kaim (written communication) showed me unpublished micrographs of a maturifusid larval shell with relatively strong non-collabral ornament. The protoconch of the species from the Cassian Formation which have been assigned to the genus Pseudotritonium (including that of the type species) are insufficiently known. According to KITTL (1894: pl. 11, figs 5, 9) the protoconch is heliciform and has an oblique axis to the teleoconch axis (but not heterostrophic). Such oblique protoconchs have been documented for some caenogastropods e.g., for the Jurassic species Plocezyga gruendeli KAIM, 2004. The teleoconch morphology of members of the Mesozoic caenogastropod family Purpurinidae resembles that of Pseudotritonium sciaphosterum. However, Purpurinidae normally have a subsutural ramp or shelf. The Hettangian type species of the cerithioid genus Paracerithium also resembles that of Pseudotritonium sciaphosterum (see GRÜNDEL 1997; HIKUROA & KAIM 2007). However, the axial ribs of Paracerithium are much stronger and form almost spines.



Textfigure 5: *Pseudotritonium sciaphosterum* (BATTEN & STOKES, 1986) formerly assigned to *Kittliconcha*, Sinbad Limestone, Moenkopi Formation, San Rafael Swell Utah, Early Triassic. (5.1): Paratype, AMNH 46314; **5.1a**, **b**: height 3.1 mm; 5.1c: detail last whorl and aperture, height 2.5 mm; 5.1d: apical view, diameter 1.8 mm. (5.2): Topotypical juvenile specimen with well-preserved protoconch, BSPG 2010 IV 1; 5.2a, b: height 0.83 mm; 5.2c: protoconch, height 0.47 mm; 5.2d: detail larval shell with non-collabral micro-ornament, height 0.24 mm.

5. Conclusions

Zygopleuroid caenogastropods play an important role in Mesozoic gastropod faunas. This group is difficult because high-spired teleoconchs with a dominant axial ornament have obviously evolved in various groups as a convergent feature. The re-study of the holotype of the type species of Kittliconcha helps clarify the systematic placement of some involved taxa. Ironically, the characters which were initially used to characterize Kittliconcha resulted from a repaired shell fracture. Only later it turned out that the genus is nevertheless valid because the protoconch morphology of Kittliconcha alias its junior synonym Settsassia is unlike that of other gastropods with zygopleuroid habitus. Kittliconcha is probably a representative of the cerithimorph stem line as is indicated by its small, smooth larval shell (see also BANDEL 1992). Similar protoconchs have been found in the Mesozoic basal cerithimorph genera Campanile, Metacerithium, and Dzikella (the latter is probably a synonym of Metacerithium) (KIEL et al. 2000; KAIM 2004; KIEL 2006). Therefore, a close phylogenetic relationship of Campanilidae and Settsassiidae seems to be possible. A relationship with Littorinimorpha or other basal caenogastropods is also possible. Only a single additional

Triassic species may actually represent Kittliconcha: K. obesa from the Middle Triassic of China. Three South American species which have been assigned to Kittliconcha are based on so poorly preserved and documented material that any generic assignment is basically speculative. The material from the Late Triassic Cassian Formation which was assigned to Kittliconcha by ZARDINI (1978) does not belong here. It represents the new sinistral genus Triadosinister and two species (including one new species) are tentatively assigned to the genus Katosira. The early Triassic species Kittliconcha sciaphostera also does not represent Kittliconcha. It is transferred to the purpurinid genus Pseudotritonium. Its cancellate ornament, shape, aperture, and the protoconch morphology suggest that this is one of the earliest members of an evolutionary line which leads to the Mesozoic family Maturifusidae which could be the stem-group of Neogastropoda (RIEDEL 2000; KAIM 2004; PONDER et al. 2008).

Acknowledgments

Andrzej KAIM is acknowledged for helpful discussion and his constructive review of the manuscript. The Deutsche For-

schungsgemeinschaft is acknowledged for financial support (Project NU 96/11-1).

6. References

- BANDEL, K. (1992): Über Caenogastropoden der Cassianer Schichten (Obertrias) der Dolomiten (Italien) und ihre taxonomische Bewertung. – Mitteilungen aus dem Geologisch-Paläontologischen Institut der Universität Hamburg, 73: 37–97.
- BANDEL, K. (1993): Caenogastropoda during Mesozoic times. Scripta Geologica, special issue, 2: 7–56.
- BANDEL, K. (1994): Comparison of Upper Triassic and Lower Jurassic Gastropods from the Peruvian Andes (Pucará Group) and the Alps (Cassian Formation). – Palaeontographica, Abteilung A, 233: 127–160.
- BATTEN, R. L. & STOKES, W. L. (1986): Early Triassic gastropods from the Sinbad Member of the Moenkopi Formation, San Rafael Swell, Utah. – America
- BONARELLI, G. (1927): Fósiles de la Formación Petrolífera o "Sistema de Salta". Boletín de la Academia Nacional de Ciencas de la república Argentina, **30:** 51–115.
- BOUCHET, P., FRÝDA, J., HAUSDORF, B., PONDER, W. F., VALDÉS, Á & WARÉN, A. (2005): Working classification of the Gastropoda. – In: P. BOUCHET & J.-P. ROCROI (Eds), Classification and Nomenclator of Gastropod Families. – Malacologia, 47: 1–397.
- DE GREGORIO, A. (1930): Fossili Anisici delle Vace di Billiemi presso, Palermo conservati nel mio privato gabinetti geologico. – Annales de Geologie et de Paleontologie, **54**: 1–40.
- DIENER, C. (1926): Fossilium Catalogus, I Animalia, **34**, Glossophora triadica.; Berlin (W. Junk), 242 pp.
- FISCHER, J. C. & WEBER, C. (1997): Révision critique de la Paléontology francaise d'Alcide d'Orbigny 2, Gastropodes Jurassiques; Paris (Muséum national d'Histoire Naturelle Paris), 300 pp.
- GOLIKOV, A. N. & STAROBOGATOV, Y. I. (1987): Systematics of the order Cerithiiformes and its position within the subclass Pectinibranchia, Eighth meeting on the investigation of molluscs. Molluscs results and perspectives of investigation. – Vsesoiuznoe soveshchaniae po izucheniiu molliuskov [Leningrad], 8: 23–28.
- GRÜNDEL, J. (1997): Zur Kenntnis einiger Gastropoden-Gattungen aus dem französischen Jura und allgemeine Bemerkungen zur Gastropodenfauna aus dem Dogger Mittel- und Westeuropas. Berliner geowissenschaftliche Abhandlungen, Reihe E, 25: 69–129.
- GRÜNDEL, J. (2001): Neritimorpha und weitere Caenogastropoda (Gastropoda) aus dem Dogger Norddeutschlands und des nordwestlichen Polens. Berliner geowissenschaftliche Abhandlungen, Reihe E, 36: 45–99.
- HAAS, O. (1953): Mesozoic invertebrate faunas of Peru. Bulletin of the American Museum of Natural History, 101: 1–328.
- HIKUROA, D. S. H. & KAIM, A. (2007): New gastropods from the Jurassic of Orville Coast, eastern Ellsworth Land, Antarctica. – Antarctic Science, **19:** 115–124.
- HOARE, R. D. & STURGEON, M. T. (1980): The Pennsylvanian pseudozygopleurid gastropod genus *Gamizyga* n. gen. from Ohio and West Virginia. Journal of Paleontology, **54**: 159–187.
- KAIM, A. (2004): The evolution of conch ontogeny in Mesozoic open sea gastropods. – Palaeontologia Polonica, 62: 1–183.
- KIEL, S. (2006): New and little-known gastropods from the Albian of the Mahajanga Basin, Northwestern Madagascar. – Journal of Paleontology, 80: 455–476.

- KIEL, S., BANDEL, K., BANJAC, N. & PERILLIAT, M. D. C. (2000): On Cretaceous Campanilidae (Caenogastropoda, Mollusca). Freiberger Forschungshefte, C, 490: 15–26.
- KITTL, E. (1894): Die Gastropoden der Schichten von St. Cassian der südalpinen Trias. II. Theil. – Annalen des Kaiserlich-Königlichen Naturhistorischen Hofmuseums, 9: 143–277.
- KLIPSTEIN, A. VON (1843): Beiträge zur geologischen Kenntnis der östlichen Alpen; Gießen (G.F. Heyers Verlag), 311 pp.
- KNIGHT, J. B.; BATTEN, R. L.; YOCHELSON, E. L. & COX, L. R. (1960): Paleozoic and some Mesozoic Caenogastropoda and Opisthobranchia. – In: R. C. MOORE (Ed.), Treatise on Invertebrate Paleontology, Part I, Mollusca 1, Supplement; Lawrence (Geological Society of America and University of Kansas Press), 1310–1324.
- KOKEN, E. (1892): Ueber die Gastropoden der rothen Schlernschichten nebst Bemerkungen über Verbreitung und Herkunft einiger triassischer Gattungen. – Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, **2**: 25–36.
- KUTASSY, A. (1940): Fossilium Catalogus, I Animalia, 81, Glossophora triadica; Berlin (W. Junk), 243–477.
- MARSHALL, B. A. (1990): An unusual triphorid (Mollusca: Gastropoda) from the Moluccas, Indonesia. – Zoologische Mededelingen (Leiden), **68**: 39–43.
- NÜTZEL, A. (1998): Über die Stammesgeschichte der Ptenoglossa (Gastropoda). – Berliner Geowissenschaftliche Abhandlungen, Reihe E, **26:** 1–229.
- NÜTZEL, A., JOACHIMSKI, M. & LÓPEZ CORREA, M. (2010): Seasonal climatic fluctuations in the Late Triassic tropics – high-resolution oxygen isotope records from aragonitic bivalve shells (Cassian Formation, Northern Italy). – Palaeogeography, Palaeoclimatology, Palaeoecology, 185: 194–204.
- OrbiGNY, A. D' (1851–1860): Paléontologie française. Terrains jurassiques. Tome II, Gastéropodes; Paris (Masson), 621 pp.
- PONDER, W. F., COLGAN, D. J., HEALY, J. M., NÜTZEL, A., SIMONE, L. R. L. & STRONG, E. E. (2008): Caenogastropoda. – In: W. F. PONDER & D. L. LINDBERG (Eds), Phylogeny and evolution of the Mollusca; Berkeley, Los Angeles, London (University of California Press), 331–383.
- RIEDEL, F. (2000): Ursprung und Evolution der "höheren" Caenogastropoda. – Berliner geowissenschaftliche Abhandlungen, Reihe E, **26:** 1–240.
- VERMEIJ, G. J., ZIPSER, E. & ZARDINI, R. (1982): Breakage-induced shell repair in some gastropods from the Upper Triassic of Italy. – Journal of Paleontology, 56: 233–235.
- WENZ, W. (1938–44): Gastropoda, Teil I. In: O. H. SCHINDEWOLF (Ed.), Handbuch der Paläozoologie, Vol. 6.; Berlin (Borntraeger), 1639 pp.
- YIN HONG-FU & YOCHELSON, E. L. (1983) Middle Triassic Gastropoda from Qingyan, Guizhou Province, China: 3 – Euomphalacea and Loxonematacea. – Journal of Paleontology, 57: 1098–1127.
- YU WEN, PAN HUA-ZHANG & WAN HUIJI (1974): Triassic Gastropoda – In: Nanjing Institute of Geology and Palaeontology, Academia Sinica (Ed.), Handbook of Stratigraphy and Palaeontology of Southwestern Region; Beijing (Publishing House of Science), 320–326.
- ZARDINI, R. (1978): Fossili Cassiani (Trias Medio-Superiore). Atlante dei Gasteropodi della formazione di S. Cassiano raccolti nella regione Dolomitica attorno a Cortina D'Ampezzo; Cortina d'Ampezzo (Edizione Ghedina), 58 pp.
- ZITTEL, K. A. (1895): Grundzüge der Palaeozoologie; München (Oldenbourg), 971 pp.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Zitteliana Serie A

Jahr/Year: 2010

Band/Volume: 50

Autor(en)/Author(s): Nützel Alexander

Artikel/Article: <u>A review of the Triassic gastropod genus Kittliconcha Bonarelli, 1927 –</u> <u>implications for the phylogeny of Caenogastropoda 9-24</u>