

Ann. Naturhist. Mus. Wien	104 A	297–305	Wien, Mai 2003
---------------------------	-------	---------	----------------

Carnivores from the Middle Miocene deposits of Grund (Lower Austria)

by Doris NAGEL¹

(With 2 textfigures and 1 plate)

Manuscript submitted on 21 October 2002,
the revised manuscript on 14 January 2002

Abstract

The carnivore remains from the Middle Miocene of Grund (MN5, Lower Austria) are described. An upper carnassial can be assigned to *Semigenetta sansaniensis*, which is the first record of this species in Austria. The scarce postcranial remains can only be identified as Carnivora indet. The caput femuri is probably close to the mustelids while the phalange remain could fit into the morphology of the amphicyonids.

Zusammenfassung


Die Raubtierreste aus der mittel-miozänen Fundstelle Grund (MN5, Niederösterreich) werden beschrieben. Ein oberer vierter Prämolare kann der *Semigenetta sansaniensis* zugeordnet werden und ist damit der erste Nachweis dieser Art in Österreich. Die wenigen postcranialen Reste können nur als Carnivora indet. bestimmt werden. Das caput femori ist mustelidenartig, dagegen könnte die Phalange am ehesten der Amphicyoniden-Morphologie entsprechen.

Key words: Carnivora, Viverridae, *Semigenetta*, Grund, MN5, Middle Miocene, Austria.

1. Introduction

The material was recovered during an excavation of the Institute of Palaeontology Vienna in 1998. ROETZEL et al. (1999) published a description of the excavation and the stratigraphy of the situation in Grund. A detailed interpretation is currently in progress (PERVESLER & ROETZEL, in prep.). After these authors, the evaluation of the profiles (B1, B2) indicate a fast change between erosion and sedimentation. Sedimentation took place in channels. At the base of profil B1 a small vertebrate fauna was found together with pulmonate snails, marine mollusc shell and pelites. Vertebrates suffered of abrasion probably because of high energy transportation. Gudrun Daxner-Höck gave the provisional faunal list in ROETZEL et al. (1999) and also made the age determination into MN5 on the presence of *Cricetodon* (DAXNER-HÖCK, this volume).

¹ Doz. Dr. Doris NAGEL, Institute of Palaeontology, Althanstrasse 14, A-1090 Vienna, Austria. – e-mail: doris.nagel@univie.ac.at



2. Systematic palaeontology

Viverridae GRAY, 1821

Semigenetta HELBING, 1927

Type species: *Semigenetta repelini* HELBING, 1927

Included species: *S. laugnacensis* DE BONIS, 1973

S. elegans DEHM, 1950

S. sansaniensis (LARTET, 1851)

S. grandis CRUSAFONT & GOLPE, 1981

S. ripolli PETTER, 1976

Viverridae are characterized by an elongated auditory bulla, transversal depression at the ectotympanic-entotympanic boundary and the presence of a weak parastyl on P⁴ (WOZENCRAFT 1989, GINSBURG 1999). *Semigenetta* shows a reduced talonide which is smaller than the trigonide. Furthermore, the talonide carries not three cusps as in other Viverridae but one and one cutting crest. M₂ is reduced (see HEIZMANN 1973). The few postcranial remains from *Semigenetta* HELBING 1927 do not differ much from the recent *Genetta*, whereas the latter has still 2 upper molars but *Semigenetta* only one (ROTH 1989).

Three larger species are assumed by HEIZMANN (1973), which probably belong to one lineage: *S. laugnacensis*, *S. elegans*, *S. sansaniensis* whereas DE BONIS (1994) argued for a fourth species – *S. steinheimensis*. In contrast to the summary by GINSBURG (1999) both authors agreed that *S. repelini* (HELBING, 1927) from Captieux belongs to *S. sansaniensis*. I follow HEIZMANN (1973) because the few specimens from Steinheim and Vieux-Collognes show only a rather large size variation, which can easier be explained for the time being by sexual dimorphism than by two different species in one locality.

The even larger *S. grandis* from Castell de Barberà (CRUSAFONT & GOLPE 1981; GOLPE 1981) originates probably from a different lineage. It appears at MN9, parallel to the smaller *S. sansaniensis* from Rudabanya and the even smaller *S. ripolli*. Two smaller forms of *Semigenetta* were described as *S. cadeoti* (ROMAN & VIRET 1934) from La Romieu and *S. ripolli* (PETTER, 1976) from Can Llobateres. *S. cadeoti* may belong to *Herpestes* as was suggested by ROTH (1989) and followed by WERDELIN (1996), but he prefers to keep this genus distinct from the recent one and refers to it as *Leptoplesictis*. Though both species are rarely known till now they surely belong to another lineage than the hitherto mentioned species. This is verified by the following characters: *S. cadeoti* is very small and the M₁ has a well developed metaconide as well as a longer talonid than e.g. *S. elegans*, which is the smallest form in the *S. elegans-sansaniensis* line (DEHM 1950). Although *S. ripolli* (PETTER 1976, Can Llobateres, MN9) is as small as *S. elegans* (MN4), it is chronologically distinct.

Semigenetta sansaniensis (LARTET, 1851)

H o l o t y p e : *S. sansaniensis* LARTET, 1851

M a t e r i a l : GRU-B1-1: left maxillar fragment with broken distal root of P³, P⁴ (10,3 x 5,5 mm) and alveolus of M¹ (NHMW 2002z0155/0001). P⁴ consists of a slender metacone blade, the paracone is about the same length as the metacone blade. The protocone is well developed, antero-lingually detached from the paracone, and at the same proximal distance as the parastyle. The tiny parastyle is connected to the paracone by a small crest. A cingulum is developed at the lingual base of the metastyle. The alveoli of M¹ are preserved and indicate a triple rooted M².

D i s c u s s i o n : *Semigenetta* is quite conservative in its morphology but varies considerably in size through time. HEIZMANN (1973) and DE BONIS (1994) used this variability to distinguish the different species. In Fig. 1 this is done for the upper P⁴ from Grund. Without taking any morphological differences into account, size alone places this tooth into *S. sansaniensis* (fig. 2). *S. laugnacensis* is the smallest representative and reported from MN2b. *S. elegans* is slightly larger than *S. laugnacensis* and only known from MN3 and MN4 (ROTH 1989, GINSBURG 1989, DEHM 1959). Morphological similarities are the very small parastyle and the cingulum developed at the lingual base of the metastyle.




Fig. 2: P₄ scatter diagramm of *Semigenetta*, length versus breadth

Comparison with Austrian specimens: '*Viverra*' *miocenica* was described by PETERS (1869) from Eibiswald/Styria (MN5). PETERS (1869) described the mandible fragment with P₃ and P₄: "similar to *Vzibethoides* and *V. sansaniensis* but because of the P₄ where the posterior accessory cusp is suppressed and the anterior one is projected forward it cannot be assigned as *V. sansaniensis*." A definite determination is not possible so far because no M₁ or P⁴ were found and for the time being it should be referred as Viverridae indet.

From Leoben (MN5) ZDARSKY (1909) reported *Viverra sansaniensis* but he already argued for another possibility: *V. leptoryncha* FILHOL, 1883. HEIZMANN (1973) agreed that this specimen has nothing to do with *S. sansaniensis*. The mandibular fragment with P₄ fits into the size distribution of *Semigenetta* but the posterior accessory cusp is placed very low and not close to the main cusp as known from viverrids. THENIUS (1948) assigned *V. leptoryncha* from Göriach to *Alopecodon*. A similar assignment of the Leoben specimen is probable.

Semigenetta indet. from Götzendorf (MN9, RÖGL et al. 1993) represents the latest occurrence of this genus in Austria. Unfortunately, the remains are too scarce to give a more detailed interpretation.

S. sansaniensis is known from Spain (Hostalets da Pierola, VILLALTA & CRUSAFONT 1943), France (Sansan, LARTET 1851; Vieux-Collogne, MEIN 1958; La Grive Saint-Alban, VIRET 1951; Captieux, HELBING 1927), Switzerland (Rümikon, HELBING 1927; Anwil, ENGESSER 1972), Germany (Steinheim, HEIZMANN 1973, Großlappen, STROMER 1940), Serbia (Sibnica, Mala Miliva; PETRONIJEVIC 1967), Hungary (Rudabánya), Ukraine (Grytsiv, Morlo pers. comm.), and now for the first time from Austria. Most of the localities are from MN5 but it is reported till MN9 (Rudabánya).

Caniformia gen. et spec. indet.

Material: GRU B1-1/ right fourth phalanx (26,7:7,4 mm), probably from a middle-sized amphicyonid; fragment of a cervical vertebra. – Inv.No. NHMW 2002z0156/0001.

Mustelidae gen. et spec. indet.

Material: GRU B1-1/ right caput femoris (10,2:10 mm), broken right under the joint facet. Form and size is mustelin-like. – Inv.No. NHMW 2002z0157/0001

Carnivora indet.

Material: vertebra fragment. – Inv.No. NHMW 2002z0158/0001.

Acknowledgements

I want to thank Gudrun Höck for entrusting me with the material. I further want to thank Michael Morlo for his helpful remarks. This paper is a contribution to the project P-15724 supported by the Austrian Science Fund, project leader Gudrun Daxner-Höck.

References




- BONIS DE, L. (1994): Carnivores viverroïdes (Carnivora, Mammalia) du Miocène inférieur d'Europe occidentale: origine des familles modernes de Feliformia. – Bulletin de la Société Géologique de France, **165**/1: 85-92. – Paris.
- CRUSAFONT, M. & GOLPE, J.M. (1981): Hallazgo de una especie del genero *Semigenetta* del Vindobonensis terminal de Castell de Barberà (Depresion prelitoral catalania; España.). – Boletín de la Real Sociedad Española de Historia Natural, (Geología) **79**: 67-76. – Madrid.
- DAXNER-HÖCK, G. (this volume): *Cricetodon meini* and other rodents from Mühlbach and Grund, Lower Austria (Middle Miocene, late MN5). – Annalen des Naturhistorischen Museums Wien, **104**/A. – Wien (this volume).
- DEHM, R. (1950): Die Raubiere aus dem Mittel-Miocän (Burdigalium) von Wintershof-West bei Eichstätt in Bayern. – Abhandlungen der Bayerischen Akademie der Wissenschaften. Mathematisch-naturwissenschaftliche Klasse, Neue Folge **58**: 1-141. – München.
- ENGESSER, B. (1972): Die obermiocäne Säugetierfauna von Anwil (Baselland). – Tätigkeitsbericht der naturforschenden Gesellschaft Baselland, **28**: 37-363. – Liestal.
- FILHOL, H. (1883): Note sur quelques mammifères de l'époque miocène. II. Observations relatives à divers carnassiers fossils provenant de La Grive-St. Alban (Isère). – Archives du Muséum d'Histoire Naturelle de Lyon, **3**: 56-69. – Lyon.
- GINSBURG, L. (1999): Order Carnivora. – pp 109-148. – In: RÖSSNER, G.E. & HEISSIG, K. (eds): The Miocene Land Mammals of Europe. – München (Dr. Friedrich Pfeil Verlag).
- GOLPE, J.M. (1981): Estudio comparativo entre la mandibula de *Semigenetta grandis* del Vindobonsense de Castell de Barberà (Depresion Prelitoral Catalana, España) y otros ejemplares europeos de *Semigenetta*. – Paleontologia i Evolució, **16**: 15-24. – Sabadell.

- HEIZMANN, E. (1973): Die Carnivoren des Steinheimer Beckens. – *Palaeontographica, Supplement* **8** (5 B): 1-95. – Stuttgart.
- HELBING, H. (1927): Une genette miocène trouvée dans les argiles de Capitieux (Gironde). – *Verhandlungen der naturforschenden Gesellschaft in Basel*, **38**: 305-315. – Basel.
- LARTET, E. (1851): Notice sur la colline de Sansan. – 45pp. – Auch.
- MEIN, P. (1958): Les Mammifères de la faune sidérolithique de Vieux-Collonges. – *Nouvelles Archives du Muséum d'Histoire Naturelle de Lyon*, **5**: 122pp. – Lyon.
- PERVESLER, P. & ROETZEL R. (eds., in prep.): Lithology, Stratigraphy and environmental Paleontology of the Grund Formation (Middle Miocene, Lower Badenium) in the Carpathian Foredeep and the Molasse Basin. – *Geologica Carpathica*.
- PETERS, K.F. (1869). Zur Kenntniss der Wirbelthiere aus den Miocänschichten von Eibiswald in Steiermark. II *Amphicyon, Viverra, Hyotherium*. – *Denkschrift der Akademie der Wissenschaften*, **29**: 189-120. – Wien.
- PETRONIJEVIC, Z. M. (1967): Die mittelmiozäne und untersarmatische (steirische) Säugetierfauna Serbiens. – *Palaeontologica Jugoslavica*, **7**: 1-157. – Zagreb.
- PETTER, G. (1976): Étude d'un nouvel ensemble de petits carnivores du Miocène d'Espagne. – *Géologie méditerranéenne*, **3/2**: 135-154. – Marseille.
- RÖGL, F., ZAPFE, H., BERNOR, R. L., BRZOBOHATY, R. L., DAXNER-HÖCK, G., DRAXLER, I., FEJFAR, O., GAUDANT, J., HERMANN, P., RABEDER, G., SCHULTZ, O. & ZETTER, R. (1993): Die Primatenfundstelle Götzendorf an der Leitha (Obermiozän des Wiener Beckens, Niederösterreich). – *Jahrbuch der Geologischen Bundesanstalt*, **136/2**: 503-526. – Wien.
- ROETZEL, R., PERVESLER, P., DAXNER-HÖCK, G., HARZHAUSER, M., MANDIC, O., ZUSCHIN, M. & CICHA, I. (1999): Grund – Kellergasse. – Arbeitstagung der Geologischen Bundesanstalt 1999 Retz-Hollabrunn: 328-334. – Wien.
- ROTH, C. (1989): Die Raubtierfauna (Carnivora, Mammalia) der untermiozänen Spaltenfüllung von Erkertshofen 2 bei Eichstätt/Bayern. – *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie*, **29**: 163-205. – München.
- ROMAN, F. & VIRET, J. (1934): La faune de mammifères du Burdigalien de la Romieu (Gers). – *Mémoires de la Société Géologique de France, (Nouvelle Série)* **21**: 1-67. – Paris.
- STROMER VON REICHENBACH, E. (1940): Die jungtertiäre Fauna des Flnzes und des Schweissandes von München. – *Abhandlungen der Bayerischen Akademie der Wissenschaften*, **48**: 1-102. – München.
- THENIUS, E. (1948): Was ist *Viverra leptoryncha* Filhol. – *Anzeiger der mathematisch-naturwissenschaftlichen Klasse der Österreichischen Akademie der Wissenschaften*, **11**: 121-123. – Wien.
- VILLALTA, J. F. & CRUSAFONT, M. (1943): Los Vertebrados del Mioceno continental de la cuenca Vallés-Penedés (Provincia de Barcelona). I: Insectívoros. II: Carnívoros. – *Boletín del Instituto geológico y Minero de España*, **56/3**: 145-336. – Madrid.
- VIRET, J. (1951): Catalogue critique de la faune des mammifères miocènes de La Grive Saint-Alban (Isère). I. Chiroptères, Carnivores, Édentés, Pholidotes. – *Nouvelles Archives du Muséum d'Histoire naturelle de Lyon*, **3**: 104 pp. – Lyon.
- WERDELIN, L. (1996): Carnivores, exclusive of Hyaenidae from the Later Miocene of Europe and Western Asia. – In: BERNOR, R.L., FAHLBUSCH, V. & MITTERMAN, H.-W. (eds.): *The Evolution of Western Eurasian Neogene Mammal Faunas*. – 271-289. – New York (Columbia University Press).


- WOZENCRAFT, Ch. (1989): The Phylogeny of the Recent Carnivora. – In: GITTELMAN, J. (ed.): Carnivore Behavior, Ecology, and Evolution. – 495-535. – New York (Cornell University Press).
- ZDARSKY, A. (1909): Die miocäne Säugetierfauna von Leoben. – Jahrbuch der kaiserlich-königlichen geologischen Reichsanstalt, **59**/2: 279-288. – Wien.

Plate 1

- Fig. 1: *Semigenetta sansaniensis* (LARTET, 1851) (NHMW 2002z0155/0001), P⁴
a-lingual, b-buccal, c-occlusal
- Fig. 2: *Caniformia* gen. et spec. indet. (NHMW 2002z0156/0001), Phalange
- Fig. 3: *Mustelidae* indet. (NHMW 2002z0157/0001), caput femori.



1 cm



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Annalen des Naturhistorischen Museums in Wien](#)

Jahr/Year: 2003

Band/Volume: [104A](#)

Autor(en)/Author(s): Nagel Doris

Artikel/Article: [Carnivores from the Middle Miocene deposits of Grund \(Lower Austria\) 297-305](#)