

Ann. Naturhist. Mus. Wien	107 A	59–69	Wien, Mai 2006
---------------------------	-------	-------	----------------

***Alvania erentoezae* – a new Rissoid gastropod species from the Early Tortonian of the Antalya Basin (Western Taurids, SW Turkey)**

by Yeşim İSLAMOĞLU¹

(With 2 text-figures and 1 plate)

Manuscript submitted on 1 April 2005,
the revised manuscript on 7 June 2005

Summary

The Miocene of the Antalya basin is represented mostly by coarse and fine detritic and partly carbonate deposits in the Western Taurids, SW Turkey. One new gastropod species belonging to the Rissoidae family is described from pelites of the Kargı section comprising Early Tortonian deposits of the Aksu formation. *Alvania erentoezae* nov. sp. is slender shaped, the teleoconch is narrow, the aperture is completely round, thickened by a varix, its last whorl makes 50 % of the total shell. On the other hand, *Alvania erentoezae* nov. sp. has paucispiral protoconch which indicates direct development with a very slight angulation, fine spiral threads on the abapical part of the whorland.

Key words: Gastropoda, Rissoidae, *Alvania*, Late Miocene, Tortonian, Mediterranean, Antalya basin, Turkey

Zusammenfassung

Miozäne Ablagerungen des Antalya Beckens (W-Tauriden, Türkei) sind vorwiegend durch grobe bis feine siliziklastika und untergeordnet durch karbonate vertreten. Eine neue rissoidae Gastropodenart wird aus den bei Kargı aufgeschlossenen unterortonen peliten der Aksu Formation beschrieben. *Alvania erentoezae* nov. sp. ist eine schlanke Form mit runder Apertur und kräftiger Varix. Der letzte Umgang macht 50 % der gesamten Schalenhöhe aus. Der paucispirale Protoconch ist Hinweis auf direkte Entwicklung und zeigt eine schwache Angulation und feine Spiralen im abapikalen Bereich der Windung.

Schlüsselwörter: Gastropoden, Rissoidae, *Alvania*, Spätes Miozän, Tortonian, Mediterranean, Antalya Becken, Türkei

Introduction

The Antalya Basin is located in the southwestern part of Turkey within the Antalya-Isparta-Alanya provinces (Fig. 1). It overlies the Precambrian – Early Cainozoic basement rocks having autochthonous or allochthonous units. The autochthonous rock units are Beydagları and Anamas-Akseki autochthones, whereas the allochthonous units are the Antalya, Alanya, Beyşehir-Hoyran-Hadım nappes (BRUNN et al. 1971; DUMONT & KEREY 1975a; MONOD 1977; POISSON 1977; AKBULUT 1977, 1980; WALDRON 1982; AKAY et al. 1985; ŞENEL et al. 1996; ŞENEL 1997a, b).

¹ Dr. Yeşim İSLAMOĞLU, Mineral Research and Exploration General Directorate, Natural History Museum, 06520 Balgat-Ankara, Turkey, e-mail: yesimislamoglu@yahoo.com




Fig. 1: Antalya Miocene basin. AA': Kargı measured section (adapted from İSLAMOĞLU 2001–2002).

The Miocene of the Antalya basin is represented mostly by coarse and fine detritic and partly carbonatic deposits. The main facies in the basin are alluvial–fan deltaic conglomerates sometimes yielding patch reefs and turbiditic detritics (BLUMENTHAL 1951; ŞENEL et al. 1996; FLECKER et al. 1995; ROBERTSON et al. 1996; KARABIYIKOĞLU et al. 1996, 2000).

Within the Miocene units, molluscan faunas were found only in the Oymapınar limestone, the Sevinç conglomerate, the Altinkaya formation and the Aksu formation (İSLAMOĞLU 2001–2002). Based on these faunas, the Miocene stratigraphy of the basin was re-evaluated (İSLAMOĞLU 2001–2002; İSLAMOĞLU & TANER 2004a, 2004b). According to this, the early Miocene Kepez travertine is the lowermost unit and lies unconformable on the

basement in the southeast of the basin. Upsection, the Burdigalian Sevinç conglomerate of alluvial fan/fan delta character overlies this unit. The Upper Burdigalian to Langhian Oymapınar limestone is composed of reefal and massive limestones and conformably overlies the Sevinç conglomerate. The Sevinç conglomerate and Oymapınar limestone are also unconformably overlying the basement rocks. The Oymapınar limestone is overlain by the Geceleme formation (Langhian) and the Upper Burdigalian to Tortonian Karpuzçay formation composed of turbiditic sediments. While the Geceleme formation is overlain by Serravallian-Tortonian units of the Karpuzçay formation along the eastern side of the basin, Upper Burdigalian-Langhian deposits of the Karpuzçay formation laterally pass into the Altinkaya formation in the central part and it is overlain by the Aksu formation at the west side of the basin. The Altinkaya formation is exposed in central and northern parts of the basin (İSLAMOĞLU 2001–2002). It is characterized by brackish water and marine settings and contains Upper Burdigalian to Langhian mollusc faunas. It unconformably overlies the basement rock units and is overlain by the Aksu formation. Laterally, the Altinkaya formation is also transitional with Upper Burdigalian to Langhian parts of the Karpuzçay formation. The Aksu formation crops out widely at western and central parts of the Miocene Antalya basin. Molluscan faunas are confined to its Early Tortonian parts. Based on the thick terrestrial sediments underlying the marine sediments, the overall age-estimation of the formation is accepted as Serravallian – Tortonian (İSLAMOĞLU 2001–2002).

The other units in the studied area are the Early Pliocene Gebiz, Eskiköy and Yenimahalle formations, the Upper Pliocene Alakilise formation, the Pleistocene Belkıs conglomerate and the Quaternary Antalya travertine and alluvions (POISSON 1977; AKAY et al. 1985).

A compressional/extensional type of tectonic regime prevailed during the Miocene in the region. The products of the compressional regime are the emplacement of Lycian nappes from northwest to southeast, NE-SW trending Aksu thrust and Kırkkavak fault which is a right – lateral strikeslip fault with reverse slip component (DUMONT & KEREY 1975b; POISSON 1977; AKAY & UYSAL 1988). This active tectonic regime especially affected the central, western and northern part of the basin and caused rapid change in environmental conditions (ÖZER et al. 1974; AKAY et al. 1985; ŞENEL et al. 1996; FLECKER et al. 1995; ROBERTSON et al. 1996; KARABIYIKOĞLU et al. 1996, 2000). Therefore, a restricted fauna was developed in the short stable periods. Generally, the reefal fauna consisting mainly of corals and coralline algae in the patch reefs (ÖZER et al. 1974; AKAY et al. 1985; AKAY & UYSAL 1988; ŞENEL et al. 1996; TUZCU et al. 1994; FLECKER et al. 1995; KARABIYIKOĞLU et al. 1996, 2000; ATABEY 1998) and molluscs in the lagoons and shallow marine environments was found in central, northern and eastern parts of the basin (İSLAMOĞLU 2001–2002, İSLAMOĞLU & TANER 2003, 2004a, b).

In terms of biogeography, the mollusc fauna was part of the Miocene Proto-Mediterranean-Atlantic region (HARZHAUSER et al. 2002), as documented by taxa such as *Cingula (Peringiella) ventricosella* CERULLI-IRELLI, 1914, *Terebralia subcorrugata* (d'ORBIGNY, 1852), *Cerithium appenninicum dertosulcata* SACCO, 1895, *Cerithium vulgatum miocenicum* VIGNAL, 1910, *Granulolabium (Tiaracerithium) pseudotiarella* (d'ORBIGNY, 1852), *Triphora adversa miocenica* COSSMANN & PEYROT, 1924, *Xenophora infundibulum* (BROCCHI, 1814).




Fig. 2: Kargı section measured from Aksu formation (simplified from İSLAMOĞLU 2001–2002).

In this study, *Alvania erentoezae* nov. sp. is found in the Early Tortonian of the Kargı section in the Aksu formation (Fig. 2). At the Kargı section, the Aksu formation starts with thick alternations of terrestrial conglomerate, sandstone and mudstone. No molluscs were detected in these sediments. Above this, an 11 m thick, massive, light grey corallgal reefal limestone follows (Fig. 2). After that, the grey mudstone with a rich mollusc fauna representing normal marine conditions is found [e.g.: *Cypraea fabagina* (LAMARCK, 1811), *Conus mercati* BROCCHI, 1814]. This unit is overlain by another level of reefal limestone. *Alvania erentoezae* nov. sp. is found in the brown colored mudstone overlying that reefal limestone. In this level, a rich mollusc fauna comprising species such as *Gibbula magus* (LINNEAUS, 1758), *Gibberulina philippi* (BUCQUOY, DAUTZENBERG & DOLLFUS, 1882), *Odostomia (Megastomia) conoidea* (BROCCHI, 1814), *Arcularia (Arcularia) ringicula* (BELLARDI, 1878) indicates an Early Tortonian age and normal marine conditions (İSLAMOĞLU 2001–2002, İSLAMOĞLU & TANER 2003, 2004a, b). The alternations of reefs and sandy mudstones with sublittoral marine mollusc fauna are interpreted as result of rapidly fluctuating environmental conditions during the Early Tortonian (İSLAMOĞLU 2001–2002, İSLAMOĞLU & TANER 2003, 2004a, b). A list of the entire molluscan fauna is given in Figure 2.

Material and Method

The investigated material was collected in the Early Tortonian levels of the Aksu formation. The specimens were extracted from pelitic samples after processing with H₂O₂ drying and screening through sieves. Shells were sputtered with gold and documented by using scanning electron microscopy.

In this study, the protoconch feature is accepted as a main distinctive factor. Besides this, the morphology and the height/width ratio (SH/SW) of the shells are used as factors for emphasizing the distinct character of that new species. Dimensions were measured after the SEM investigations and all figured specimens are deposited in the collection of the NHMW with the collection number 2005z0034/0001-0003 (Natural History Museum Vienna).

Systematic Paleontology

Class Gastropoda CUVIER, 1797

Subclass Caenogastropoda COX, 1959

Order Littorinimorpha GOLIKOV & STAROBOGATOV, 1975

Superfamily Rissooidea GRAY, 1847

Family Rissoidae GRAY, 1847

Subfamily Rissoinae GRAY, 1847

Genus: *Alvania* RISSO, 1826

Type species: *Turbo cimex* LINNEAUS, 1758 from the Mediterranean Sea and NE Atlantic (KOWALKE & HARZHAUSER 2004).

In the recent studies, *Alvania* is regarded as belonging to the Rissoinae subfamily, because of exhibiting similarities with the genus *Rissoa* concerning teleoconch sculpture, operculum, radular characters and protoconch sculptures (KOWALKE & HARZHAUSER 2004). Generally, *Alvania* develops very small, egg shaped and reticulate sculptured shells with rounded whorls and sometimes thickened varix at the aperture (WENZ 1938). In addition, the protoconch morphology and sculpture exhibit a coarser larval sculpture with occasionally occurring zigzag-shaped spiral striae (KOWALKE & HARZHAUSER 2004). Therefore, the new species is included within *Alvania*.

***Alvania erentoezae* nov. sp.**

(Plate 1, Fig. 1–3)

H o l o t y p e: NHMW 2005z0034/0001.

L o c u s t y p i c u s e t s t r a t u m t y p i c u m: 12 km south of Kargı, Aksu formation in Antalya basin, Turkey (Isparta N25b4 X1: 06150 Y1: 27350), pelites within the Kargı section (meter 30 of Fig. 2).

D e r i v a t i o n o m i n i s: This species is dedicated to Dr. Lütfiye ERÜNAL ERENTÖZ (Mineral Research and Exploration, retired) in memory and honour to her contributions to the Neogene Molluscan biostratigraphy in Turkey.

D e s c r i p t i o n: Small-sized, slender consisting of 5 convex whorls including the protoconch separated by a deep suture line. Spire narrow with an angle of 35°. Sculpture consisting of 11–12 axial ribs and 5–6 spirals; body-whorl rather narrow attaining about 50 % of the total shell height. Aperture small, rounded, with thick varix. The protoconch consisting of 1.5 rounded whorls is paucispiral, points to direct development, very slight angulation, fine spiral threads on the abapical part of the whorland.

M e a s u r e m e n t s: Shell height: 1.43–2 mm, shell width: 0.79–1.23 mm, last whorl height: 0.97 mm, aperture height: 0.60 mm, aperture width: 0.60 mm, varix width: 275 micron, protoconch height: 0.2–0.3 mm, protoconch width: 0.25–0.37 mm.

R e m a r k s: Mainly, *Alvania* species are divided into two categories according to their protoconch features as having multispiral indirect development or not (LOZOUET 1998, KOWALKE & HARZHAUSER 2004). While the species with multispiral indirect development have a protoconch with a few whorls or more, the species showing direct developments have a very short protoconch (LOZOUET 1998). *Alvania erentoezae* nov. sp. has a short protoconch indicating direct larval development. In first view, it differs from the ones having multispiral protoconchs such as the Late Oligocene to Middle Miocene species *Alvania (Alvania) oceani* (d'ORBIGNY, 1852) (COSSMANN & PEYROT 1919: 587–588, pl. 16, fig. 11–112; KOWALKE & HARZHAUSER 2004: 120, fig. 7A) and the Miocene species *Alvania (Alvania) venus* (d'ORBIGNY, 1852) (COSSMANN & PEYROT 1919: 585–587, pl. 17, fig. 23–24). The protoconch of *Alvania erentoezae* nov. sp. with its 1.5 rounded whorls allows a separation from *Alvania oceani* which develops a protoconch consisting of 2.5 slightly rounded whorls with a slightly thickened, fractioned sinusigera notch. Similarly, *Alvania venus* displays a considerably longer protoconch with 2.75 whorls (KOWALKE & HARZHAUSER 2004). Among the other *Alvania* groups

having short protoconch which indicate non-planktotrophic development, *Alvania erentoezae* nov. sp. differs from the Late Oligocene and Early Miocene species *Alvania aturensis* LOZOUET, 1998 (LOZOUET 1998: 78, fig. 8 k-l; 11 a-b) and the Late Aquitanian species *Alvania andraldensis* LOZOUET, 1998 (LOZOUET 1998: 76, fig. 8 f-h) in its sculpture. The protoconch of *Alvania erentoezae* nov. sp. shows a very slight angulation and fine spiral threads on the abapical part of the whorland. The new species has compared to the original collection of type species of *Alvania aturensis* LOZOUET, 1998 and *Alvania andraldensis* LOZOUET, 1998 in NHMN of Paris. Thus, *Alvania andraldensis* LOZOUET, 1998 is very similar but differs by its ovoid shape and the fewer number of teleoconch whorls (3 canalaculate whorls). But, the protoconch of *Alvania andraldensis* is carinate with spiral cords; the body whorl attains 64 % of total shell height. The protoconch of *Alvania aturensis* differs by its fine opisthocryt striae. Apart from the protoconch features, the shell measurements and ratios of the similar *Alvania* species to new one is compared. Therefore, it is found that *Alvania erentoezae* nov. sp. has more elongate, slender shape and the rounded aperture and its last whorl makes 50% of the total shell. While SH/SW rates of *Alvania erentoezae* nov. sp. range around 1.94, this ratio is 1.96 for *Alvania andraldensis* LOZOUET, 1998, 1.77 for *Alvania (Alvania) venus* (d'ORBIGNY, 1852), 1.6 for *Alvania aturensis* LOZOUET, 1998 and 1.48 for *Alvania (Alvania) oceani* (d'ORBIGNY, 1852).

D i s t r i b u t i o n: Early Tortonian in the Antalya Miocene basin of SW Turkey.

Conclusion

One new gastropod species belonging to the family Rissoidae is described from the Miocene Antalya Basin. *Alvania erentoezae* nov. sp. is found in the early Tortonian Aksu formation at the Kargı section. It has short protoconch which indicates non-planktotrophic development. It differs from the other species having short protoconch by showing a very slight angulation and fine spiral threads on the abapical part of the whorland. Apart from this, while comparing to the shell measurements and ratios, the shape of it is slender, the teleoconch is narrow, the aperture is completely round with a thickened varix.

On the other hand, while the all Middle Miocene (Badenian) Central Paratethian *Alvania* species have long (multispiral) protoconch pointing to planktotrophic development (KOWALKE & HARZHAUSER 2004), a part of Oligocene – Early Miocene *Alvania* species (LOZOUET 1998) and the Late Miocene *Alvania erentoezae* nov. sp. have a short protoconch indicating direct development.

Acknowledgements

This study was granted by the SYNTHESYS Program. The author expresses sincere thanks to Dr. Mathias HARZHAUSER (Director of the Geological-Paleontological Department of Natural History Museum Vienna) for providing access to rare literature and valuable discussions, to Alice SCHUMACHER (Natural History Museum of Vienna) for electron microscope photographs (SEM) and to Dr. Pierre LOZOUET (Natural History Museum of Paris) for showing the type species and given helpful comments on the manuscript.

References

- AKAY, E. & UYSAL, Ş. (1988): Post-Eocene tectonics of Mid-Taurids. – Bulletin of the Mineral Research and Exploration, **108**: 57–68 (in Turkish).
- , UYSAL, Ş., POISSON, A., CRAVETTE, Y. & MULLER, C. (1985): Stratigraphy of Antalya Neogene basin. – Geology Bulletin of Turkey, **28**: 105–109 [in Turkish].
- AKBULUT, A. (1977): Etude géologique d'une partie du Taurus occidental aud' Eğirdir (Turquie). – These 3eme Cycle Univercite Paris-Sud-Orsay: 203 pp.
- (1980): Geology of Çandır (Sütçüler, Isparta) region in the south of Eğirdir lake, Western Taurids. – Geology Bulletin of Turkey, **23/1**: 1–10 [in Turkish].
- ATABEY, N. (1998): Paleocology and sedimentologic environments of Miocene red algae in the Western Taurids belt. – 51 th. Geological Congress of Turkey, Abstracts: 58 [in Turkish].
- BELLARDI, L. (1878): I molluschi dei terreni terziarii del Piemonte e della Liguria. – Torino.
- BLUMENTHAL, M. (1951): Geological researchs of Alanya region in the Western Taurids. – Bulletin of the Mineral Research and Exploration, (D. Ser.) **5**: 134 p. – Ankara [in Turkish].
- BROCCHI, G. V. (1814): Conchiologia fossile subappennina, con osservazioni sugli Appennini e sul suolo adjacente. – 712 pp. – Milano (Dalla Stamperia Reale).
- BRUNN, J.H., DUMONT, J.F., GRACIANSKY, P.CH. De, GUTNIC, M., JUTEAU, TH., MARCOUX, J., MONOD, O. & POISSON, A. (1971): Outline of the Geology of the Western Taurids. – Geology and History of Turkey, Petroleum Exploration Society of Libya. – Tripoli.
- BUCQUOY, E., DAUTZENBERG & DOLLFUS, G.F. (1882): Mollusques Marins du Roussillon. t. I, Gastropodes.
- CERULLI-IRELLI, S. (1914): Fauna malacologia marina, Parte I-III. – Paleontographica Italica.
- COSSMANN, M. & PEYROT, A. (1919): Conchologie néogénique de l'Aquitaine. – Acta de la Société Linnéenne de Bordeaux, **70/3**: 181–356; **70/4**: 357–491.
- & — (1924): Conchologie Neogénique de l'Aquitaine, **4**. – Bordeaux.
- COX, L. R. (1959): Thoughts on the classification of the Gastropoda. – Proceedings of the Malacological Society of London, **33**: 239–261.
- CUVIER, G. (1797): Tableau élémentaire de l'histoire naturelle des animaux. – 710 pp. – Paris (Baudoin).
- DUMONT, J.F. & KEREY, K. (1975) a. Basement geological study in the south of Eğirdir lake. – Geology Bulletin of Turkey, **18/2**: 169–174 [in Turkish].
- & — (1975) b. Kırkkavak fault: a N-W strike-slip fault in the boundary of Western Taurids and Köprüçay basins. – Geology Bulletin of Turkey, **18/1**: 59–62 [in Turkish].
- FLECKER, R., ROBERTSON, A.H.F., POISSON, A. & MÜLLER, C. (1995): Facies and tectonic significance of two contrasting Miocene basins in south coastal Turkey. – Terra Nova, **7**: 221–232.
- GOLIKOV, A. N. & STAROBOGATOV, Y. A. I. (1975): Systematics of prosobranch gastropods. – Malacologia, **15/1**: 185–232.
- GRAY, J. E. (1847): A list of the genera of Recent Mollusca, their synonyma and types. – Proceedings of the Zoological Society of London, **15**: 129–219.
- HARZHAUSER, M., PILLER, W. E. & STEININGER, F. F. (2002): Circum Mediterranean Oligo - Miocene biogeographic evolution – the gastropods' point of view. – Palaeogeography, Palaeoclimatology, Palaeoecology, **183**: 103–133.
- İSLAMOĞLU, Y. (2001–2002): The molluscan fauna and stratigraphy of Antalya Miocene basin (West-Central Taurids, SW Turkey). – Bulletin of the Mineral Research and Exploration, **123-124**: 27–58.
- & TANER, G. (2003): Paleogeographic and paleoecologic features of Miocene molluscan fauna of Antalya and Kasaba basins. – Bulletin of the Mineral Research and Exploration, **126**: 11–42.
- & — (2004a): Bivalvia and Scaphopoda fauna of Antalya Miocene basin. – Bulletin of the Mineral Research and Exploration, **127**: 1–27 [in Turkish].
- & — (2004b): Gastropoda fauna of Antalya Miocene basin. – Bulletin of the Mineral Research and Exploration, **127**: 29–65 [in Turkish].
- KARABIYIKOĞLU, M., TUZCU, S., ÇUHADAR, Ö., İSLAMOĞLU, Y. & ATABEY, N. (1996): Batı Toroslar Aksu önülke havzası resifal Miyosen çökel dolgununun litofasiyes analizi, çökelme sistemleri ve tektono-sedimenter evrimi. – 49. Geological Congress of Turkey, Abstracts: 25–26.
- , ÇİNER, A., MONOD, O., DEYNAUX, M., TUZCU, S. & ÖRÇEN, S. (2000): Tectonosedimentary evolution of the Miocene Manavgat Basin, Western Taurids, Turkey. – In: BOZKURT, E., WINCHESTER, J.A. & PIPER, J.D.A. (eds.): Tectonics and Magmatism in Turkey and Surrounding Area. – Geological Society of London, special publication, **173**: 271–294.
- KOWALKE, T. & HARZHAUSER, M. (2004): Early ontogeny and palaeoecology of mid-Miocene rissoid gastropods of the Central Paratethys. – Acta Palaontologica Polonica, **49/1**: 111–134.
- LAMARCK, J. B. (1811): De la détermination des mollusques testacés. – Annales du museum, **17**: 54–80.
- LINNAEUS, C. (1758): Systema naturae per regna tria naturae, secundum Classes, Ordines, genera, species, cum characteribus, differentiis, synonymis, locis. – Editio Decima, Reformata [not seen].
- LOZOUET, P. (1998): Nouvelles espèces de Gastéropodes (Mollusca: Gastropoda) de l'Oligocène et du Miocène inférieur de l'Aquitaine (sud-ouest de la France). – Cossmanniana, **5/3-4**: 61–102.
- MONOD, O. (1977): Recherches géologique dans le Taurus occidental au sud de Beyşehir (Turquie). – These. University of Paris-Sud Orsay.
- d'ORBIGNY, A. (1852): Prodrome de Paléontologie Stratigraphique Universelle des Animaux Mollusques et Rayonnés faisant suite au Cours Elémentaire de Paléontologie et de Géologie Stratigraphique, **3**. 196 + 190 pp. – Paris (Masson).
- ÖZER, B., BIJU-DUVAL, P., COURRIER, P. & LETOUZEY, J. (1974): Geology of Antalya – Mut and Adana basins. – Proceedings of Second Petrol Congress of Turkey: 57–84 [in Turkish].
- POISSON, A. (1977): Recherches Géologiques dans les Taurides occidentales (Turquie). – These, l'universite de Paris-Sud (Centre D'Orsay), Tome 1, Paris.
- RISSE, A., 1826. Histoire naturelle des principales productions de l'Europe meridionale et principalement de celles des environs de Nice et des Alpes maritimes. 4. – Paris et Strasbourg.
- ROBERTSON, A.H.F., COLLINS, A., FLECKER, R., GLOVER, C., PICKETT, E., USTAÖMER, T. & DIXON (1996): Tectonic evolution of western Turkey from upper Paleozoic to recent. – 49th Geological Congress of Turkey, Abstracts: 9–14.
- SACCO, F. (1895): I Molluschi dei terreni terziarii del Piemonte e della Liguria. – Memorie Royal Accademia delle Scienze Torino, **23**.

- ŞENEL, M. (1997a): 1:250 000 scaled Turkey Geological map, Antalya section, 3, Mineral Research and Exploration General Directorate, Geological Department. – Ankara [in Turkish].
- (1997b): 1:250 000 scaled Turkey Geological map, Isparta section, 4, Mineral Research and Exploration General Directorate, Geological Department: 47 pp. – Ankara [in Turkish].
- , GEDIK, I., DALKILIÇ, H., SERDAROĞLU, M., BILGIN, A.Z., UĞUZ, M.F., BÖLÜKBAŞI, A.S., KORUCU, M. & ÖZGÜL, N. (1996): Stratigraphy of allochthons and autochthones in the east of Isparta angle (Western Taurids). – Bulletin of the Mineral Research and Exploration, **118**: 111–160 [in Turkish].
- TUZCU, S., KARABIYIKOĞLU, M. & İSLAMOĞLU, Y. (1994): Batı Toroslar Miyosen Mercan resifleri: Bileşimleri, fasiyes özellikleri ve ortamsal konumları. – 47. Geological Congress of Turkey, Abstracts: 16.
- VIGNAL, L. (1910): Cerithiidae du Tertiaire superieur extrait. – Journal de Conchyliologie, **58**. – Paris.
- WALDRON, J.W.F. (1982): Antalya Karmaşığı kuzeydoğu uzanımının Isparta bölgesindeki stratigrafisi ve sedimenter evrimi. – Maden Tetkik ve Arama Dergisi, **97**: 1–20.
- WENZ, W. (1938): Gastropoda, Teil 1. – In: O. H. SCHINDEWOLF (ed.): Handbuch der Paläozoologie, **6/1**: 1639 pp. – Berlin (Borntraeger).

Plate 1

Fig. 1–3: *Alvania erentoezae* nov. sp.
Scale bar: 0.5 mm

Fig. 1: height: 1.43 mm, width: 1.03 mm, protoconch height: 0.3 mm, protoconch width: 0.33 mm (Holotype: NHMW 2005z0034/0001)

Fig. 2: height: 2.0 mm, width: 1.23 mm, protoconch height: 0.24 mm, protoconch width: 0.37 mm (Paratype: NHMW 2005z0034/0002)

Fig. 3: shell height: 1.58 mm, shell width: 0.79 mm, protoconch height: 0.2 mm, protoconch width: 0.25 mm (Paratype: NHMW 2005z0034/0003)

