

Ann. Naturhist. Mus. Wien	109 A	19–27	Wien, September 2007
---------------------------	-------	-------	----------------------

Taxonomic identity of *Eostega lebedinskyi* LAMBRECHT, 1929 (Aves) from the middle Eocene of Romania

by Jiří MLÍKOVSKÝ¹

Manuscript submitted on February 27th 2006,
the revised manuscript on December 4th 2006

(With 4 text-figures)

Summary

Eostega lebedinskyi LAMBRECHT, 1929 from the middle Eocene of Cluj-Manastur in Romania was identified as a member of the modern family Sulidae. It is the second oldest representative of the Sulidae known so far.

Zusammenfassung

Eostega lebedinskyi LAMBRECHT, 1929 aus dem Mitteleozän von Cluj-Manastur in Rumänien wurde als Mitglied der rezenten Familie Sulidae erkannt. Es ist der zweitälteste Vertreter der Familie, der bis jetzt beschrieben wurde.

Introduction

Eostega lebedinskyi LAMBRECHT, 1929 is one of the very few avian taxa described from the Paleogene of Eastern Europe (MLÍKOVSKÝ 1996, 2002; MAYR 2005). The bird was described by LAMBRECHT (1929) on the basis of an incomplete mandible found in the limestone quarry at Koloszmanostor near Koloszvár in southeastern Austria-Hungary (now Manastur in Cluj in northern Romania) prior to 1891. In 1890, the specimen was presented by János NÁGY to the Society of Transylvanian Museum (Erdélyi Múzeum Egyesület) according to an old label written in Hungarian and attached to the specimen. At a later date, the specimen was forwarded by Anton KOCH to the Naturhistorisches Hofmuseum (now Naturhistorisches Museum) in Wien, Austria (LAMBRECHT 1929: 1269), where it is still deposited, bearing inventory number 1930/0001/0013.

The specimen was originally thought to be the bill of a turtle according to the Hungarian label, but LAMBRECHT (1929) correctly recognized its avian affinities and described on its basis *Eostega lebedinskyi*. He stated that *Eostega* is a "steganopode" (= pelecaniiform sensu WETMORE 1930, 1960) bird, but did not include it in a family, concluding only that it is intermediate between *Phalacrocorax* (i.e. Phalacrocoracidae) and *Sula* (i.e. Sulidae). Later, LAMBRECHT (1933) included *Eostega* in his new family Elopterygidae, based on *Elopteryx* ANDREWS, 1913 from the late Cretaceous of Romania. However, the

¹ Department of Zoology, National Museum, Václavské náměstí 68, 115 79 Praha 1, Czech Republic. – E-mail: jiri.mlikovsky@nm.cz

latter genus is probably non-avian (OLSON 1985: 93). BRODKORB (1963) continued to list *Eostega* in the Eopterygidae. OLSON (1985: 193) correctly stated that *Eostega* "is quite probably peleciform, but needs comparison with other Eocene Peleciformes such as *Prophaethon*, *Limnofregata*, and *Protplotus*, before its familial allocation can be determined."

I reexamined the holotypical mandible of *Eostega lebedinskyi* LAMBRECHT, 1929 with results presented below.

Systematic paleontology

Order Ardeiformes WAGLER, 1831

Family Sulidae REICHENBACH, 1849

Genus *Eostega* LAMBRECHT, 1929

Eostega LAMBRECHT, 1929: 1272 [Type (by monotypy and n.g. n.sp. convention): *Eostega lebedinskyi* LAMBRECHT, 1929.]

Masillastega MAYR, 2002: 508 [Type (by original designation): *Masillastega rectirostris* MAYR, 2002.]

Included species: *Eostega lebedinskyi* LAMBRECHT, 1929, and *Eostega rectirostris* (MAYR, 2002), new combination.

Diagnosis (LAMBRECHT 1929): Extensive description and comparisons, but no diagnosis given.

Diagnosis (LAMBRECHT 1933: 289): "Mandible with double proc. coronoideus as in *Phalacrocorax* and *Sula*, but less pneumatized" (my translation).

Diagnosis (here emended): Typical sulid with a strong, straight mandible, having a single fenestra, which differs from mandibles of modern sulid genera in having: (1) processus coronoideus located above internal fenestra mandibularis, (2) medial fossa aditus canalis neurovascularis sharply bordered and narrow, (3) cotyla medialis of the fossa articularis quadratica long and slender, (4) cotyla caudalis of the same fossa narrow and clearly bordered posteriorly, and (5) dorsal margin of the intermediate part of ramus mandibulae rather narrow, with medial ridge lower than lateral ridge.

Distribution: Middle Eocene (Lutetian) of Cluj-Manastur, Romania, and Messel, Germany.

Eostega lebedinskyi LAMBRECHT, 1929

Eostega lebedinskyi LAMBRECHT, 1929: 1272 (Figured by LAMBRECHT 1929, fig. 12-13, and LAMBRECHT 1933, fig. 103).

Holotype: Partial mandible in slab, lacking symphyseal part of mandibular ramus; Naturhistorisches Museum Wien, 1930/0001/0013.

Age and locality: Cluj-Manastur, Cluj Province, Romania; 46.47 N, 23.36 E. Middle Eocene, Lutetian, MP 13 (see LAMBRECHT 1929, BOMBIȚA 1963, VLAICU-TĂȚĂRIM 1963, KESSLER 1996).




Fig. 1: Original label attached to the specimen.

F a m i l i a l a f f i l i a t i o n : Robust mandible with a broad and posteriorly flattened caudal part of rami clearly indicates that *Eostega lebedinskyi* belongs to the Pelecaniformes (sensu WETMORE 1930, 1960). Within this order, the holotypical mandible of *Eostega lebedinskyi* differs from the same element of the Pelecanidae in having: (1) caudal joint laterally not narrowed, (2) posterior facet of the caudal joint not deeply excavated, and (3) fossa aditus canalis mandibulae not restricted, and (4) in being not swollen. It differs from the same element of the Fregatidae in having: (1) Rami not flattened latero-medially, (2) posterior facet of the caudal joint not declined laterally, (3) caudal joint more broad in dorsal view, and (4) processus coronoideus more broad. It differs from the same element of the Phaethontidae in having: (1) processus coronoideus present, (2) caudal joint broad in dorsal view, and (3) caudal joint oval, not rounded. It differs from the same element of the Phalacrocoracidae and Anhingidae in having: (1) rami dorso-ventrally broad, (2) fossa auditus canalis neurovascularis mandibulae more restricted, (3) caudal joint broad, and (4) caudal joint not separated from the intermedial part of rami by a neck. It differs from the same element of the fossil Pelagornithidae (sensu lato) in having no bony teeth (cf. HARRISON & WALKER 1976b). It differs from the same element of the ancestral phaethon *Prophaethon shrubsolei* ANDREWS, 1899 from the early Eocene of England, currently separated at the familial level as Prophaethontidae (HARRISON & WALKER 1976a, 1977), in having: (1) caudal joint dorso-ventrally broad, and (2) rami heavily build. It differs from the same element of the ancestral frigatebird *Limnofregata azygosternon* OLSON, 1977 in having: (1) rami more robust, and (2) caudal joint dorso-ventrally broad. Mandibles of the fossil Plotopteridae are unknown, but this highly specialized family is known only from the Oligocene and early Miocene of the northern part of the Atlantic Ocean, incl. California, Washinton and Japan (HOWARD 1969; OLSON 1979, 1980; OLSON & HASEGAWA 1979, 1985, 1996; GOEDERT 1988; GOEDERT & CORNISH 2002). Its occurrence in the middle Eocene Paratethys Sea is thus less probable.




On the other hand, the holotypical mandible of *Eostega lebedinskyi* agrees with the same element of the Sulidae in having: (1) rami robust, but not swollen, (2) posterior facet of the caudal joint almost flat, (3) caudal joint not narrowed laterally, (4) processus coronoideus well developed, and (5) caudal joint broad, but not rounded. Although it is difficult to identify isolated bones of early Tertiary birds (cf. OLSON 1985), it is possible to conclude, that *Eostega lebedinskyi* belongs in the family Sulidae.

Generic affiliation (Tab. 1): The family Sulidae consists of three modern and several Neogene genera. The modern genera include *Sula* BRISSON, 1760, *Morus* VIEILLOT, 1816, and *Papasula* OLSON & WARHEIT, 1988 (see OLSON & WARHEIT 1988, VAN TETS et al. 1988). Below, I present a more detailed description of the holotypical mandible of *Eostega lebedinskyi* and add in parentheses, whether the given condition is similar (+) to that found in the modern genera, or differs (x) from it: (1) processus coronoideus is narrow and distinctly bordered (x *Sula*, *Papasula*, + *Morus*, where it is somewhat longer, and posteriorly less well bordered), (2) processus coronoideus is located above the anterior end of medial fossa aditus canalis (x *Sula* and *Morus*, where it is located more posteriorly, x *Papasula*, where it is located more anteriorly), (3) fossa aditus canalis is well defined (x *Sula*, + *Morus* and *Papasula*), (4) fossa aditus canalis narrow (+ *Sula*, x *Morus*, where it is large and broad, x *Papasula*, where it is very narrow and very short), (5) cotyla medialis long and slender (x *Sula*, *Morus* and *Papasula*, where it is short and rounded), (6) cotyla caudalis narrow (x *Sula*, *Morus* and *Papasula*, where it is broader), (7) cotyla caudalis posteriorly clearly bordered (x *Sula*, *Morus* and *Papasula*, where the posterior border is less distinct), (8) posterior facet of the caudal joint positioned in the axis of ramus (+ *Papasula*, x *Sula* and *Morus*, where it is bend medially), (9) intermediate part of ramus not swollen (+ *Sula* and *Morus*, x *Papasula*, where this part of ramus is swollen), (10) dorsal border of rami narrow (x *Sula*, *Morus* and *Papasula*, where it is broader), (11) medial ridge of the dorsal border of rami lower than lateral ridge (x *Sula*, *Morus* and *Papasula*, where this condition applies only to the posterior end of the rami, while medial and lateral ridges are equally high elsewhere). These characters clearly indicate, that *Eostega* is different from all modern genera of sulids.

Tab. 1. Measurements of the bills of *Eostega lebedinskyi* and Recent sulids [in mm]. A – length from the posterior end of the bill to the anterior end of coronoid facet. B – length from the posterior end of the bill to the anterior end of os angulare (measured on the medial side of the bone). C – maximum transversal width of the joint.

	n	A	B	C
<i>Eostega lebedinskyi</i>	1	ca. 33.5	74	15+
<i>Papasula abboti</i>	1	29.5	50	15.0
<i>Sula leucogaster</i>	3	21.5–23.0	29–30	11.2–12.0
<i>Sula neboxii</i>	3	28.0–30.0	48–50	13.5–13.9
<i>Sula variegata</i>	1	28.0	43	13.8
<i>Sula dactylatra</i>	3	29.5–31.0	53–54	14.6–14.8
<i>Sula sula</i>	3	22.0–23.5	39–40	11.3–11.8
<i>Morus bassanus</i>	3	34.0–35.0	64–65	16.9–17.1
<i>Morus capensis</i>	3	30.5–31.5	55–56	14.8–15.0
<i>Morus serrator</i>	3	30.0–32.0	52–56	14.6–14.8

The Sulidae have a rich Neogene record (BRODKORB 1963, HARRISON 1978, OLSON 1985, MLÍKOVSKÝ 2002), but only few Paleogene fossils have been assigned to the Sulidae so far.



Figs. 2–4: Holotype of *Eostega lebedinskyi* LAMBRECHT. (2) General view, (3) detail of left ramus and (4) detail of right ramus (NHMW 1930/0001/0013).

Masillastega rectirostris MAYR, 2002 was described on the basis of a skull from the middle Eocene (MP 11) of Messel, Germany (MAYR 2002). The mandible of this species is very similar to that of *Eostega*. MAYR (2002) claimed that mandibular rami are convex in *Eostega*, while they are straight in *Masillastega*. In fact, mandibular rami are

straight in both these forms. Indeed, the only difference MAYR (2002) did find was in the size of the two mandibles, maximum length being about 152 mm in *Eostega* (estimated, because the tip of the mandible is absent – LAMBRECHT 1929) and 135 mm in *Masillastega*. Such a slight difference in size alone is certainly no reason for separating genera. Hence, I synonymize here *Masillastega* MAYR, 2002 with *Eostega* LAMBRECHT, 1929. Until more data on Paleogene sulids of Europe are available I retain *rectirostris* as a separate species, transferring it to the genus *Eostega* LAMBRECHT, 1929 as *Eostega rectirostris* (MAYR, 2002), new combination.

Mergus ronzonei GERVAIS, 1848-1852 from the early Oligocene (MP 21) of Ronzon, France, was transferred to the genus *Sula* BRISSON, 1760 by MILNE-EDWARDS (1867: 271). HARRISON (1975a) suggested that the species belongs in the Phalacrocoracidae. I restudied the holotype of this species – an imperfect pelvis in a slab –, concluding that it is indeterminate at the genus and perhaps even at the family level and relegating it to the Aves incertae sedis (MLÍKOVSKÝ 2002: 260; see also CHENEVAL 1984, OLSON 1985).

Sula arvernensis MILNE-EDWARDS, 1867 was described on the basis of an incomplete pelvis in a slab and an incomplete sternum in a slab, both of which originated from the late Oligocene (MP 30) of Gannat, France. HARRISON (1975a,b) separated *arvernensis* at the genus level as *Empheresula* HARRISON, 1975b. The taxonomic position of *Empheresula arvernensis* within the Sulidae is uncertain. A comparison with *Eostega lebedinskyi* is impossible because of the lack of similar elements. However, even if these two forms would be found congeneric in future, the genus name *Eostega* LAMBRECHT, 1929 would have priority of *Empheresula* HARRISON, 1975b.

Neogene genera of sulids include *Microsula* WETMORE, 1938 (Maryland), *Palaeosula* HOWARD, 1958 (California), *Miosula* MILLER, 1925 (California), *Enkurosula* KAŠIN, 1977 (France, incl. *Pseudosula* HARRISON, 1975a, which is preoccupied), and *Sarmatosula* GRIGORESCU & KESSLER, 1977 (Romania). Of them, *Microsula* WETMORE, 1938 was synonymized with the modern *Morus* VIEILLOT, 1816 (OLSON & RASMUSSEN 2001). Mandibles were not described for any of these genera, so that direct comparison with *Eostega* is impossible. Moreover, these genera do not reflect proper phylogeny of the Sulidae and should be restudied (OLSON 1985 and pers. communication in 1997, MLÍKOVSKÝ 2002). Until the Oligo-Miocene history of sulids is clarified, *Eostega* should be regarded as a valid genus.

Paleogene sulids of Europe

The record of Paleogene sulids of Europe is limited to *Eostega rectirostris* (MAYR) from the middle Eocene (Lutetian, MP 11) of Messel, Germany (MAYR 2002), *Eostega lebedinskyi* LAMBRECHT from the middle Eocene (Lutetian, MP 13) of Cluj, Romania (LAMBRECHT 1929, 1933, MLÍKOVSKÝ 2002, this paper), and to an unidentified sulid from the late Oligocene of Thalberg, Germany (GÖHLICH in DARGA et al. 1999). Although the record is meager, it shows that sulids were members of the Paleogene Paratethys avifauna at least in the middle Eocene and in the late Oligocene.

Acknowledgments

I was allowed to study the holotype of *Eostega lebedinskyi* owing to courtesy of Ortwin SCHULTZ and Gudrun DAXNER-HÖCK (Wien). Storrs L. OLSON (Washington, D.C.) allowed me to use the comparative collection of avian skeletons in the National Museum of Natural History of the Smithsonian Institution in Washington, D.C., when I was a short-term fellow of the Institution in January/February 1997. Daniel GOUJET (Paris) allowed me to study fossil birds in the collections of the Muséum National d'Histoire Naturelle in Paris, France. Erika GÁL (Budapest) kindly translated for me the old Hungarian label attached to the holotype. Two anonymous referees provided helpful comments on an earlier draft the manuscript. I am obliged to all of these persons.

References

- ANDREWS, C.W. (1899): On the remains of a new bird from the London Clay of Sheppey. – Proc. zool. Soc. London **1899**: 776-785.
- (1913): On some bird remains from the Upper Cretaceous of Transsylvania. – Geol. Mag. (5) **10**: 193-196.
- BOMBIȚA, G. (1963): Contribuții la corelarea eocenului epicontinental din R. P. Română [Contributions to the correlation of the epicontinental Eocene in Romania]. – 115 pp. – București (Editura Academiei Republicii Populare Română). [In Romanian.]
- BRODKORB, P. (1963): Catalogue of fossil birds: Part 1 (Archaeopterygiformes through Ardeiformes). – Bull. Florida State Mus., Biol. Sci. **7**: 179-293.
- CHENEVAL, J. (1984): Les oiseaux aquatiques (Gaviiformes à Ansériiformes) du gisement aquitain de Saint-Gérard-le-Puy (Allier, France): révision systématique. – Palaeovertebrata, **14**: 33-115.
- DARGA, R., M. BÖHME, U.B. GÖHLICH & G.E. RÖSSNER (1999): Reste höherer Wirbelthiere aus dem Alttertiär des Alpenvorlandes bei Siegsdorf/Oberbayern. – Mitt. Bayer. Staatssamml. Paläontol. hist. Geol., **39**: 91-114.
- GERVAIS, P. (1848-1852): Zoologie et Paléontologie françaises (Animaux vertébrés), ou Nouvelles recherches sur les animaux vivants et fossiles de la France. Vol. 1. – iv + 271 pp. – Paris (Arthus).
- GOEDERT, J.L. (1988): A new late Eocene species of Plotopteridae (Aves: Pelecaniformes) from northwestern Oregon. – Proc. Calif. Acad. Sci., **45**: 97-102.
- GOEDERT, J.L. & J. CORNISH (2002): A preliminary report on the diversity and stratigraphic distribution of the Plotopteridae (Pelecaniformes) in Paleogene rocks of Washington State, USA. – In: Z. ZHOU & F. ZHANG (Eds.): Proceedings of the 5th Symposium of the Society of Avian Paleontology and Evolution: 63-76. – Beijing (Science Press).
- GRIGORESCU D. & E. KESSLER (1977): The middle Sarmatian avian fauna of South Dobrogea. – Revue roum. Géol., Géophys. Géogr. (Géologie) **21**: 93-108.
- HARRISON, C.J.O. (1975a): The taxonomic status of MILNE-EDWARD'S [sic] fossil sulids. – Bull. Brit. Ornithol. Club, **95**: 51-54.
- (1975b): *Empheresula*: a new name for *Parasula* HARRISON 1975. – Bull. British Ornithol. Club, **95**: 175.
- (1978): Fossil boobies and gannets. – In: J.B. NELSON: The Sulidae: gannets and boobies: 974-977. – London (Oxford University Press)

- HARRISON, C.J.O. & C.A. WALKER (1976a): A reappraisal of *Prophathon shrubsolei* ANDREWS (Aves). – Bull. Brit. Mus. (Nat. Hist.), Geology **27**: 1-30.
- (1976b): A review of the bony-toothed birds (Odontopterygiformes): with descriptions of some new species. – Tertiary Res., Spec. Paper **2**: 1-62.
- (1977): Birds of the British Lower Eocene. – Tertiary Res., Spec. Paper **3**: 1-52.
- HOWARD, H. (1958): Miocene sulids of southern California. – Los Angeles Co. Mus., Contrib. Sci., **25**: 1-15.
- (1969): A new avian fossil from Kern County, California. – Condor, **71**: 68-69.
- KAŠIN, G.N. (1977): Novoe rodovoe nazvanie ptic vzamen preokkupirovanogo [A new generic name of birds replacing a preoccupied one]. – Paleontol. Ž., 1977/3: 139.
- KESSLER E. (1996): Tertiary avian localities of Romania. – Acta Univ. Carol., Geologica **39**: 703-709. – Praha.
- LAMBRECHT, K. (1929): Mesozoische und tertiäre Vogelreste aus Siebenbürgen. – In: E. SCIKI (Ed.): X^e congrès international de zoologie: 1262-1275. – Budapest (Stephaneum).
- (1933): Handbuch der Palaeornithologie. – 1024 pp. – Berlin (Gebr. Borntraeger).
- MAYR, G. (2002): A skull of a new peleciform bird from the Middle Eocene of Messel, Germany. – Acta palaeontol. pol., **47**: 507-512.
- (2005): The Paleogene fossil record of birds in Europe. – Biol. Rev., **80**: 515-542.
- MILLER, L. (1925): Avian remains from the Miocene of Lompoc, California. – Carnegie Inst. Washington, Publ. **349**: 107-117.
- MILNE-EDWARDS, A. (1867-1868): Recherches anatomiques et paléontologiques pour servir à l'histoire des oiseaux fossiles de la France. Vol. 1. – 472 pp. – Paris (Victor Masson et Fils).
- MLÍKOVSKÝ, J. (1996): Tertiary avian faunas of Europe. – Acta Univ. Carol., Geologica, **39**: 777-818. – Praha.
- MLÍKOVSKÝ, J. (2002): Cenozoic birds of the world. Part 1. Europe. – 406 pp. – Praha (Ninox Press).
- OLSON, S.L. (1977): A Lower Eocene frigatebird from the Green River Formation of Wyoming (Pelecaniformes: Fregatidae). – Smithson. Contrib. Paleobiol., **35**: i-iii, 1-33.
- (1979): Fossil counterparts of giant penguins from the North Pacific. – Science, **206**: 688.
- (1980): A new genus of penguin-like peleciform bird from the Oligocene of Washington (Pelecaniformes: Plotopteridae). – Nat. Hist. Mus. Los Angeles, Contrib. Sci., **330**: 51-57.
- (1985): The fossil record of birds. – In: D.S. FARNER & J.R. KING (Eds.): Avian biology **8**: 79-252. – New York (Academic Press).
- OLSON, S.L. & Y. HASEGAWA (1979): Fossil counterparts of giant penguins from the North Pacific. – Science, **206**: 688-689.
- (1985): A femur of *Plotopterus* from the early middle Miocene of Japan (Pelecaniformes: Plotopteridae). – Bull. natn. Sci. Mus. Tokyo, (Ser. C) **11**: 137-140.

- (1996): A new genus and two new species of gigantic Plotopteridae from Japan (Aves: Plotopteridae). – *J. Vert. Paleontol.*, **16**: 742-751.
- OLSON, S.L. & RASMUSSEN, P. (2001): Miocene and Pliocene birds from the Lee Creek Mine, North Carolina. – In: RAY, C.E. & BOHASKA, D.J. (eds.): *Geology and paleontology of the Lee Creek Mine, North Carolina, III*. – *Smithsonian Contributions to Paleobiology*, **90**: 233-365.
- , K.I. WARHEIT (1988): A new genus for *Sula abbotti*. – *Bull. Brit. Orn. Club*, **108**: 9-12.
- VAN TETS, G.F., C.W. MEREDITH, P.J. FULLAGAR & P.M. DAVIDSON (1988): Osteological differences between *Sula* and *Morus*, and a description of an extinct new species of *Sula* from Lord Howe and Norfolk Inlands, Tasman Sea. – *Notornis*, **35**: 35-57.
- VLAICU-TĂTĂRIM, N. (1963): Stratigrafia eocenului din regiunea de la sud-vest de Cluj [Stratigraphy of the Eocene in the region south-west of Cluj]. – 204 pp. – București (Editura Academiei Republicii Populare Română). [In Romanian.]
- WETMORE, A. (1930) A systematic classification for the birds of the world. – *Proc. U. S. natl. Mus.*, **76/24**: 1-8.
- (1938): A Miocene booby and other records from the Calvert Formation of Maryland. – *Proc. U. S. natl. Mus.*, **85**: 21-25.
- (1960): A classification for the birds of the world. – *Smithsonian misc. Coll.*, **139/11**: 1-37.