

# A new locality of *Groenlandia densa* (L.) FOURR. (*Potamogetonaceae*) in the surroundings of the southern Harz Mountains (Central Germany)

St. Meyer\* & A. Hoch\*\*

*Groenlandia densa* (L.) FOURR. (= *Potamogeton densus* L.) constitutes one of the exceedingly rare plant species ascribed to the southern Harz Mountain area and its surroundings (Saxony-Anhalt, Thuringia). The first reliable record of *Groenlandia densa* in this region can be found in WALLROTH 1822: "Potamogeton densus L., In fossis uliginosis ad Wiehe" (51°16'N, 11°25'E).

Seen herbarium material from this region:

*Potamogeton densus* L.: Flora thuring[iaca], In Gewässern bei Grossfurra, 08.1892 Hermann Sterzing (51°23'N, 10°48'E) [GOET, PRC]; – Tümpel b[ei] Nordhausen, 20.07.1892 Louis Osswald (51°29'N, 10°48'E) [JE], see also OSSWALD 1893; *Potamogeton densus* L. var. *lancifolius* Koch: Flora hercynica, in fossis Nordhaus[en], 23.07.1892 Adolf Vocke (51°29'N, 10°48'E) [W 2007-0005336, PRC]; Flora hercynica, in fossis aqua repletis Nordhausen, 23.07.1892 Adolf Vocke (51°29'N, 10°48'E) [PRC]; – Flora hercynica, in aquis stagnantibus Nordhaus[en], 23.07.1892 Adolf Vocke (51°29'N, 10°48'E) [JE]; *Potamogeton densus* L. var. *lanceolata*: In fossis Ziegelei pr[ope] Petersdorf, 07/08.1892 Adolf Vocke (51°31'N, 10°49'E) [GOET].

Of these documented localities in the surroundings of Großfurra, Wiehe, Petersdorf, and Nordhausen today exist no actual records since more than 100 years.

In the vegetation period of 2009, a new locality has been found: Saxony-Anhalt, district of Mansfeld-Südharz: Einzingen, alt. c. 170 m, 25.06.2009 Armin Hoch (51°26.717'N, 11°21.642'E) [GOET]. This locality is c. 20–40 km distant from the earlier known places. There are more than 5.000 individuals growing in a small pond in the middle of the village. The pond has an inflow from several springs as well as a permanent outflow.

## Acknowledgements

We thank the curators Ernst Vitek (W), Jochen Heinrichs (GOET), Jan Stepánek (PRC) and Jochen Müller (JE) for granting access to the material and for assistance during personal visits of the authors.

## References

- OSSWALD L., 1893: Beiträge zur Flora von Nord-Thüringen. – Mitt. Thüring. Bot. Ver. 3/4: 57-59.  
WALLROTH F.W., 1822: Schedulae criticae de plantis floriae Halensis selectis. Corollarium novum ad C. Sprengelii Floram Halensem. Accedunt generum quorumdam specierumque omnium definitiones novae, excursus in stirpes difficiliores et icones V Tomus I, Phanerogamia. – Halae: Car. Aug. Kümmelius.

Stefan Meyer, Georg-August-University of Göttingen, Albrecht-von-Haller-Institut for Plant Sciences, Dept. of Ecology and Ecosystem Research, Untere Karspüle 2, D-37073 Göttingen, Germany – smeyer1@gwdg.de

\*\* Armin Hoch, Karstlandschaft Südharz Biosphere Reserve, Hallesche Straße 68a, D-06536 Roßla, Germany – armin.hoch@lvwa.sachsen-anhalt.de

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Annalen des Naturhistorischen Museums in Wien](#)

Jahr/Year: 2009

Band/Volume: [111B](#)

Autor(en)/Author(s): Meyer Stefan, Hoch Armin

Artikel/Article: [A new locality of \*Groenlandia densa\* \(L.\) Fourr. \(Potamogetonaceae\) in the surroundings of the southern Harz Mountains \(Central Germany\). 285](#)