113	619–653	Wien, Mai 2011
	113	113 619–653

Early Barremian dasycladalean algae from Serre de Bleyton (Drôme, SE France)

By Ioan I. BUCUR¹

(With 7 plates and 3 figures)

Manuscript submitted on September 22th 2010, the revised manuscript on December 17th 2011

Abstract

Dasycladalean calcareous algae are described from the Barremian deposits from Serre de Bleyton (Drôme, France). The association consists of 15 taxa and is characteristic for the Lower Barremian series of the northern margin of the Tethys. The algae have been transported by gravitational flows (turbidites) from a shallower area, *i.e.* a carbonate platform margin, to a deeper, basinal setting.

Keywords: calcareous algae, dasycladales, Early Barremian, SE France

Zusammenfassung

Dasycladale Kalkalgen aus Ablagerungen des Barremium von Serre de Bleyton (Drôme, Frankreich) werden beschrieben. Die Vergesellschaftung besteht aus 15 Arten und ist typisch für das Untere Barremium des Tethys Nordrandes. Die Algen stammen aus Flachwasserbereichen (z.B. dem Rand einer Karbonatplatform) und wurden durch Turbidite in tiefere Beckensedimente verfrachtet.

Schlüsselworte: Kalkalgen, Dasycladales, Unteres Barremium, Südost-Frankreich

Introduction

This paper is part of a larger project deeling with the description of the micro- and macropaleontologic assemblages from Serre de Bleyton (SE France), and refers to a rich assemblage of isolated, nicely-preserved specimens of dasycladalean algae. The samples have been collected by Gero MOOSLEITNER (Salzburg) from Lower Cretaceous turbiditic sediments cropping out in a small area on the southeastern flank of the Serre de Bleyton

¹ Babeş-Bolyai University, Department of Geology, str. M. Kogălniceanu nr. 1, 400084 Cluj-Napoca, Romania; e-mail: ioan.bucur@ubbcluj.ro



Fig. 1. Location of the study area (asterisk indicates sampling site).

Range (Fig. 1). Besides algae, foraminifers and numerous other fossils assigned to various groups of invertebrates have been also recovered. Some of these have been already studied, the results having been published in 2010 by JÄGER (crinoids), JANSSEN (belemnites), LÖSER (corals), LUKENDER (ammonoids), RIEGRAF & MOOSLEITNER (rhyncolites), TAYLOR (bryozoans), and VILLIER (asteroids). For the the general characteristics of the sediments providing the Serre de Bleyton fossils, see KROH et al. (2010). Calcareous algae were collected from the sample profiles 1 and 2 described by KROH et al. (2010).

Material and Methods

The selected paleontological material has been washed and sorted by Gero MOOSLEITNER; it was recovered from the alteration area of outcrops 1 and 2 (KROH et al. 2010). The algae have been studied under a stereomicroscope. Images of the studied material were obtained using both the stereomicroscope, and scanning electron microscope (SEM). In order to achieve a better assignment at species level, a large number of specimens were embedded in synthetic resin and thin sectioned. These samples were studied under a petrographic microscope and images made by using a digital camera. The identified algal association consists of 15 taxa (11 determined at species level, while 4 at the genus level, with an uncertain assignment in the case of one of them). No quantitative evaluation has been preformed, but one can estimate that the most frequent species within the association are represented by *Actinoporella podolica*, *Salpingoporella muehlbergii* and *Pseudoactinoporella fragilis*.

R e pository: The studied specimens and thin sections are housed at the Paleontological Museum, Department of Geology, Babeş-Bolyai University Cluj-Napoca, under the inventory numbers 23813 to 23837.

Systematic Palaeontology

Order Dasycladales PASCHER, 1931

Family Polyphysaceae Kützing, 1843

Genus Actinoporella GÜMBEL in ALTH, 1882

Actinoporella podolica (Alth, 1878), emend. Conrad, Praturlon & Radoičić, 1974

(Pl. 1, Figs 1–5; Pl. 3, Figs 1–3; Pl. 4, Figs 1–6; Pl. 6, Figs 6, 7, 11; Pl. 7, Figs 1–6)

Selected synonymy

1878 Gyroporella podolica ALTH – ALTH: 83, Pl. VI, Figs 1-8

- 1970 Verticiloporella dalmatica RAVIV & LORCH CONRAD: 69, Pl. 5, Figs 1–3.
- 1974 Actinoporella podolica Gümbel in Alth, 1882 Conrad et al.: 4, Figs 1(1-8), 2 (1-7), 4-6, 8-12.
- 1974 Actinoporella sp. CANÉROT, Pl. X, Figs 4-6.
- 1978 Actinoporella podolica (ALTH, 1878) emend. CONRAD, PRATURLON & RADOIČIĆ, 1974 –
- BASSOULLET et al.: 30, Pl.2, Figs 4–7 [with extended synonymy].
- 1980 Ainoporella podolica Gümbel Arnaud-Vanneau, Pl. 112, Fig. 4.
- 1989 Actinoporella podolica (Alth, 1878) Dragastan: 28, Pl. 14, Figs 4–5.
- 1992 Actinoporella podolica GÜMBEL, emend. CONRAD et al., 1974 -
 - RAVIV & LORCH, Pl. 2, Figs a-o, Text-fig. 7.
- 1994 Actinoporella podolica (ALTH) GRANIER, Pl. 1, Figs 1, 4, 8–10.
- 1996 Actinoporella podolica (ALTH) CLAPS et al., Pl. 3, Fig. 14.

1996 Actinoporella podolica Conrad, Praturlon & Radoičić, 1974 – Sokač: 27, Pl. XX, Figs 1–3, 5–13.

- 2001 Falsolikanella sp. BUCUR, Pl. 12, Figs 3-7
- 2001 Actinoporella? sp. BUCUR, Pl. 12, Fig. 8
- 2006 Actinoporella podolica HUSINEC & SOKAČ, Figs 4H(c), 4I.
- 2006 Actinoporella podolica (ALTH) SCHLAGINTWEIT et al., Fig. 3 (3).
- 2008 Actinoporella sp. gr. podolica GRANIER, Pl. 2, Figs F, G.
- 2008 Actinoporella podolica (ALTH) HOSSEINI & CONRAD, Pl. 3, Figs B-E.
- 2010 Actinoporella podolica (ALTH) TAHERPOUR et al.: 4, Fig. 6f.

Description and remarks: Isolated verticilated discs resulted from breaking of the thallus at interverticillar space level (Pl. 1, Figs 1–5; Pl.3, Figs 1–3). The verticils consist of long, well-interconnected phloiophorous laterals within the verticil and slightly inclined towards the top. The axial cavity is relatively narrow compared to the external diameter of verticils, and is surrounded by circular bulges on both upper and lower sides resulting from calcification of the small secondary laterals located on one side and the other of the coronae. This feature allows a relatively straightforward identification of this alga by examining the external morphology of verticils only.

Nevertheless, the thin sections cross-cutting through numerous isolated verticils (Pl. 4, Figs. 1–6; Pl. 6, Figs. 6, 7, 11; Pl. 7, Figs. 1–6) do not always reveal the diagnostic features

Annalen des Naturhistorischen Museums in Wien, Serie A 113



Fig. 2. Two different types of transverse (horizontal) sections through verticillar discs of *Actinoporella podolica*. A-A' – the section cut the corona-like structure; B-B' – the section don't cut the corona-like structure.

of the species. In some longitudinal or diagonal sections to the thallus axis (Pl. 7, Figs 3, 6) the corona-type structures can be recognized, which allow a relatively simple identification. Transversal sections to the thallus axis (*i.e.* horizontal sections at verticil level) do not always expose corona-type structures. Fig. 2 illustrates two types of transversal sections that differ by the level at which the verticil is segmented. In the case of an A-A' section, the segmentation plane cross-cuts the corona-type structures, thus they are clearly visible (*e.g.*, Pl. 4, Figs. 1–3). In the case of a B-B' section, the corona-type structures are not intercepted, while the general aspect of the alga may be mistaken as similar sections through the verticils of other species (*e.g. Clypeina* or *Falsolikanella*) (*e.g.* Pl. 4, Figs 4–6). It is also worth mentioning that in some large specimens with numerous laterals, the latter show a certain degree of imbrication that imprints a peculiar aspect to the sections shown (*e.g.*, Pl. 6, Figs 6, 7, 11; Pl. 7, Figs 1, 2, 4, 5).

CONRAD et al. (1974) presented a detailed description of *Actinoporella podolica* as well as a thallus reconstruction. When considering that calcification is more intense in the verticil (the laterals being closely connected), one can easily relate this to the frequent breaking of the thallus at interverticilar level, leading to the frequent presence of this alga only as isolated verticils. The general thallus morphology with verticils in vertical succession is fossilised only in internal platform, low energy depositional environments (*e.g.* RADOIČIĆ 1968: Pl. XVI, Fig. 1, as "*Dasycladacea*" dalmatica).

622

Dimensions:

D = 0.72-2.45 mmd = 0.30-0.82 mm d/D = 0.20-0.47 l = 0.5-1.02 p(distal) = 0.15-0.40 w = 16-26

Genus Clypeina MICHELIN, 1845

Clypeina sp.

(Pl. 6, Figs 12–13)

Remarks: Two of the sectioned disc-like verticils proved to belong to a *Clypeina* species. Even if not very frequent, *Clypeina* specimens are present in the assemblage together with *Actinoporella*, as isolated discs.

Dimensions:

D = 1.0 mm d = 0.52 mm l = 0.30 mmp (diatal) = 0.12 mm

Genus Pseudoclypeina RADOIČIĆ, 1975, non 1970

Pseudoclypeina sp.

(Pl. 1, Figs 1- pars, 8; Pl. 4, Figs 15-17; Pl. 5, Figs 1-3)

Remarks: A relatively large amount of fragments of disc-like verticils, characterised by a distal "bulbous" enlargement was found in the *Actinoporella*-type assemblage (Pl. 1, Fig. 1). The thin sections (Pl. 4, Figs 15–17; Pl. 5, Figs 1–3) point to the existence of some long primary laterals that carry a bush of shorter secondary laterals. This type of verticilated discs can be assigned to two genera: *Rajkaella* (DRAGASTAN & BUCUR 1988) and *Pseudoclypeina* (RADOIČIĆ 1975). As a rule, *Rajkaella* is characterized by verticilated discs with free primary laterals along most of the lateral length (DRAGASTAN & BUCUR 1993), while in the case of *Pseudoclypeina* species the laterals of one verticil coalesce over the whole length (RADOIČIĆ 1970). The differences between *Rajkaella* (as *Radoiciciella*) and *Pseudoclypeina* were emphasized by DRAGASTAN & BUCUR (1993) and GRANIER (1990), even if some of the specimens originally assigned by DRAGASTAN & BUCUR (1993) to *Radoiciciella* (= *Rajkaella*) could be more likely attributed to *Pseudoclypeina* (*e.g.*, MASSE 1976: Pl. 5, Fig. 16; FOURCADE et al. 1972: Pl. 2, Figs 5–8). We have assigned the specimens identified at Serre de Bleyton to the genus *Pseudoclypeina* exactly based on the compact aspect of the verticilated discs. However, it is clear that a review of all algae assigned to the two genera would be needed for a final clarification of their relationships.

Dimensions

 $l_1 = 0.70 - 0.85 \text{ mm}$ $l_2 = 0.15 - 0.30 \text{ mm}$ $p_1 = 0.10 - 0.20 \text{ mm}$ $p_2 = 0.05 - 0.10 \text{ mm}$

Family Triploporellaceae Berger & KAEVER, 1992

Genus Angioporella Conrad, Masse & Radoičić, 1973

Angioporella fouryae Conrad, Masse & Radoičić, 1973

(Pl. 2, Figs 1, 2; Pl. 3, Fig. 4; Pl. 4, Figs 7-11)

Selected synonymy

1973 Angioporella fouryae n. gen., n. sp. - MASSE et al.: 384, Text-fig. 1, Pl. 1, Figs 1-8.

1976 Angioporella fouryae MASSE et al., 1973 – CONRAD & PEYBERNÈS: 180, Fig. 5(a)

1980 Angioporella fouryae Masse, Conrad & Radoičić – Arnaud-Vanneau, Pl. 112, Fig. 9.

1994 Angioporella aff. fouryae Masse, Conrad & Radoičić – Bucur: 151, Pl. II, Fig. 28.

2007 Angioporella fouryae MASSE, CONRAD & RADOIČIĆ – BUCUR et al., Fig. 4 (11).

Description and remarks: Thallus cylindrical, externally slightly undulated, made up of a succession of verticils of fertile laterals (large and less numerous) and sterile laterals (tubular, and more numerous). The supposed fertile laterals cannot be observed at the exterior because they are surrounded by the distal parts of the sterile laterals, vertically oriented. Specimens submitted to advanced dissolution with acetic acid (Pl. 2, Fig. 1) show the morphology of the fertile laterals, the distal part of which is quadrangular, higher than wider. The thin sections (Pl. 4, Figs 7–10) provide supplementary data on the internal structure. The central cavity is also slightly undulated, but if the external constrictions are set at the level of sterile laterals, the axial cavity is larger at this level and thinner at the level of the fertile laterals. Longitudinal and longitudinal-oblique sections from Pl. 4, Figs 7–10 illustrate also the morphology of the fertile laterals, higher than wide, and almost trapezoidal in outline. The sterile laterals, tubular, slightly enlarged to the exterior, are visible in the transverse section of Pl. 4, Fig. 11.

I did not observe any elements that can be ascribed to reproductive structures within the cavities corresponding to the fertile laterals. Also, I can not observe with certitude the peduncular pore linking the fertile laterals to the axial cavity. The possible connection visible at the upper-left two verticils of the specimen in Pl. 4, Fig. 10 could represent the effect of secondary dissolution. A possible explanation for the absence of the link pore between fertile laterals and the axial cavity is given by MASSE et al. (1973) by the relatively late calcification of the fertile ampulae.

624

625

Dimensions

L (maximum observed) = 3.6 mmD = 0.75-1.55 mmd = 0.18-0.90 mmh = 0.78-0.90 mml (sterile) = 0.35 mmp (sterile) = 0.07-0.10 mmp (fertile) = 0.35-0.70 mmw (sterile) = 29-30w (fertile) = approx. 8

Genus Deloffrella GRANIER & MICHAUD, 1987

Deloffrella quercifoliifpora GRANIER & MICHAUD, 1987 (Pl. 6, Fig. 1)

Selected synonymy

1987 Deloffrella quercifoliipora n. gen., n. sp. – GRANIER & MICHAUD: 1093, Text-fig. 3, Pl. I, Figs 1–10. 2001 Deloffrella quercifoliipora GRANIER & MICHAUD – BUCUR, Pl. 3, Figs 4–6. 2002 Deloffrella quercifoliipora GRANIER & MICHAUD – BUCUR: 42, Pl. 3, Figs 1–5 [with synonymy]. 2003 Deloffrella quercifoliipora GRANIER & MICHAUD – BUCUR et al.: 218, Pl. I, Figs 1–3. 2005 Deloffrella quercifoliipora GRANIER & MICHAUD – SCHLAGINTWEIT, Pl. II, Figs 8, 10, 12.

Description: The only specimen in longitudinal-oblique section was found accidentally, when sectioning *Salpingoporella* samples. Specimens of *Deloffrella* could not be identified by external morphological examination only, given their scarcity in the association, on the one hand, and the possibility of taking them as less well-preserved specimens of *Salpingoporella*, on the other. The specimen illustrated in Pl. 6, Fig. 1 provides insight into the structure of the laterals. At least three orders of laterals are visible, the relatively larger primary ones being clearly noticeable both in the longitudinal and oblique parts of the section.

R e m a r k s: Based on the available published information, this represents the first mention of the species *Deloffrella quercifoliipora* in Lower Cretaceous deposits in southeastern France. Following CONRAD (pers. comm.) although rare, *Deloffrella quercifoliipora* is present in Berriasian and upper Barremian deposits of Switzerland (CLAVEL et al., unpublished). Annalen des Naturhistorischen Museums in Wien, Serie A 113

Dimensions

L (maximum observed) = 1.65 mmD = 0.46 mmD = 0.12 mmH = 0.10 mm

Genus Falsolikanella GRANIER, 1987

Falsolikanella danilovae (RADOIČIĆ ex BARATTOLO, 1978) GRANIER et al., 2000 (Pl. 2, Figs 9–12; Pl. 3, Fig. 7; Pl. 4, Figs 12–14)

Selected synonymy

- 1968 Likanella? danilovae n. sp. RADOIČIĆ: 238, Pl. I, Figs 1–3; Pl. II, Figs 1–6; Pl. III, Figs 1–6; Pl. IV, Figs 1–4; Pl. V, Figs 1–4; Pl. VI, Figs 1–5; Pl. VII, Figs 1–4.
- 1973 Likanella(?) danilovae RADOIČIĆ, 1968 JAFFREZO: 81, Pl. 2, Figs 12, 13.
- 1978 Likanella? danilovae RADOIČIĆ, 1968 BASSOULLET et al.: 142, Pl. 16, Fig. 9–11 [with synonymy].
- 1979 Likanella(?) danilovae ARNAUD-VANNEAU, Pl. 110, Figs 1-2
- 1989 Likanella? aff. danilovae Radoičić, 1968 Conrad & Masse: 282, Pl. II, Fig. 7
- 1993 Likanella? danilovae Luperto Sinni & Masse, Pl. 1, Figs 14, 15.
- 1994 Praturlonella danilovae (RADOIČIĆ) 1968 BUCUR: 153, Pl. VII, Figs 1-8.
- 2000 Falsolikanella danilovae (RADOIČIĆ, 1975) BUCUR: 58, Pl. V, Figs 1-11, 13, 14.
- 2000 Falsolikanella danilovae Radoičić ex Barattolo, 1978, n. comb. Granier et al.: 177, Pl. I,

Figs 1–4, 6.

2007 Falsolikanella danilovae (RADOIČIĆ) – BUCUR et al.: 133, Fig. 4(13).

Description and remarks: *Falsolikanella* is present in the dasycladalean association as subspherical or thick-discoidal corpuscles with rounded borders ("segments" or "rings" of the moniliform-type thallus; see RADOIČIĆ 1968: Fig. 1). Externally, (Pl. 2, Figs 9–12; Pl. 3, Fig. 7) the laterals distal termination can be noticed. The thin sections illustrated in Pl. 4, Figs 12–14 represent horizontal-subhorizontal sections through the subspheroidal segments (*i.e.* transversal sections at the level of the segments of laterals as related to the axis of the thallus).

A short comment on the generic assignment of this alga (*Likanella*, *Selliporella*, *Praturlonella*, *Falsolikanella*) is given by BUCUR (2000). Further discussion on this assignment is beyond of the scope of this paper, even if maybe necessary, taking into account that some authors still attribute this species to other genera.

Dimensions

D = 0.8-1.9 mmD = 0.25-0.70 mm L = 0.35-0.60 mm P (distal) = 0.12-0.25 mm

626

Genus Montiella Morellet & Morellet, 1922

Montiella? elitzae (BAKALOVA, 1971)

Selected synonymy

- 1970 Cylindroporella elitzae sp. BAKALOVA: 126, Pl. III, Figs 1–8.
- 1973 Cylindroporella maslovi n. sp. SRIVASTAVA: 699, Figs 16–18.
- 1974 Cylindroporella barnesii (cf. SAINT-MARC, 1970) PRESTAT, Pl. 8, Fig. 3.
- 1978 Cylindroporella elitzae BAKALOVA, 1971 BASSOULLET et al.: 71, Pl. 7, Figs 6, 7.
- 1979 Cylindroporella sp. cf. C. sugdeni ELLIOTT AZEMA et al., Pl. XXXI, Figs. 1-4.
- 1980 *Montiella? elitzae* (Вакаlovа) nov. comb. Radoičić: 114, Pl. 1, Figs 1, 2; Pl. 2, Figs 1–4; Pl. 3, Figs 1–4.
- 1986 Cylindroporella cf. elitzae BAKALOVA MU: 86, Pl. V, Figs 3, 4, 6–8.
- 1989 Montiella? elitzae (BAKALOVA, 1971) RADOIČIĆ, 1980 CONRAD & MASSE: 282, Pl. II, Figs 1–3.
- 1993 *Bakalovaella elitzae* (BAKALOVA), nov. comb. BUCUR: 100, Pl. &, Figs 1–18 [with extended synonymy]
- 1994 Bakalovaella elitzae (BAKALOVA, 1971) BUCUR: 156, Pl. XIII, Figs. 3–12.
- 2000 Bakalovaella elitzae (BAKALOVA) BUCUR et al.: 442, Pl. 3, Figs 1-5.
- 2000 Bakalovaella elitzae (BAKALOVA, 1971) DRAGASTAN: 129, Pl. 2, Figs 16-17.
- 2001 Bakalovaella elitzae (BAKALOVA) BUCUR, Pl. 13, Figs 1–3.
- 2003 Bakalovaella elitzae (BAKALOVA, 1971) BUCUR et al.: 218, Pl. I, Fig. 6.
- 2007 Montiella elitzae (BAKALOVA) BUCUR et al.: 133, Fig. 4(8).
- 2010 Montiella? elitzae (BAKALOVA, 1971) RADOIČIĆ, 1980 TAHERPOUR et al., Fig. 8 (a-d).

Description and remarks: As in the case of *Deloffrella*, *Montiella*? *elitzae* has been identified only in thin sections, in the attempt to obtain sections through specimens of *Piriferella paucicalcarea* – a species that is essentially similar based only on external morphology. The longitudinal-tangential sections in Pl. 6, Figs 2–3 show the arrangement of the large ovoidal, fertile ampoules in relationship to the subjacent tubular, sterile laterals showing a significantly-smaller diameter. The section illustrated in Pl. 6, Fig. 2 crosscuts, in its central part, the two types of laterals, close to the point where these diverge from the primary lateral.

Montiella? elitzae has been described in detail by BUCUR (1993, under *Bakalovaella elitzae*). Additional data on the general morphology of its thallus have been published by TAHERPOUR et al. (2010) by the description and illustration of a complete, exceptionally well-preserved specimen in longitudinal section. This presented both the lower, pedunculate side lacking fertile ampoules, and the upper, terminal side lacking sterile secondaries.

Concerning the generic assignment, the genus *Bakalovaella* and the combination *Bakalovaella elitae* proposed by BUCUR (1993) has not been well-received by most subsequent authors. It is however worth to include here the comment in Taherpour et al. (2010): "In the Paleocene type-species *M. munieri*, the presence of two types of secondary laterals is inferred, yet not physically demonstrated. In view of this equivocal situation, the genus *Bakalovaella* BUCUR, 1993 (type-species *B. elitzae*) will possibly prove to be a well founded taxon rather than a junior synonym of *Montiella*".

Annalen des Naturhistorischen Museums in Wien, Serie A 113

Dimensions

L (maximum observed) = 3.7 mm (up to 7 mm in Taherpour et al., 2010) D = 0.85-1.1 mmP (sterile) = 0.09 mmP (fertile) = 0.25-0.30 mm

Genus Piriferella Sokač, 1996

Piriferella paucicalcarea (Conrad, 1970) Conrad, Schlagintweit & Bucur, 2009

Selected synonymy
1970 Heteroporella? paucicalcarea n. sp. – CONRAD: 68, Text-fig. 5, Pls III, IV.
1970 Heteroporella? paucicalcarea CONRAD, 1970 – JAFFREZO: 80, Pl. 3, Figs 9, 11, 16.
1978 Heteroporella? paucicalcarea CONRAD, 1970 – BASSOULLET et al.: 133, Pl. 15, Figs 5–8.
1978 Heteroporella? paucicalcarea CONRAD – ARNAUD-VANNEAU, Pl. 111, Figs 3–5.
1989 Heteroporella? paucicalcarea CONRAD, 1970 – CONRAD & MASSE: 281, Pl. II, Fig. 11.
1994 Heteroporella? paucicalcarea CONRAD, 1970 – BUCUR: 152, Pl. VI, Fig. 13.
2000 Similiclypeina paucicalcarea (CONRAD, 1970) nov. comb. – BUCUR: 60, Pl. IV, Fig. 7. [with extended synonymy]
2007 Similiclypeina paucicalcarea (CONRAD) – BUCUR et al.: 133, Fig. 4(6, 9).
2009 Piriferella paucicalcarea (CONRAD, 1970) n. comb. – CONRAD et al.: 28(c)

Description: The detached specimens show large and somehow irregular pores at the periphery (Pl. 1, Figs 6, 7), representing the distal termination of the laterals. They are better noticed in the broken specimen illustrated in Pl. 2, Fig. 13. The thin sections (Pl. 6, Figs 5, 8) do not provide additional information. For a detailed description see CONRAD (1970) and BASSOULLET et al. (1978).

Genus Pseudoactinoporella Conrad, 1970 emend. Conrad & Peybernes, 1976

Pseudoactinoporella fragilis CONRAD, 1970, emend. CONRAD & PEYBERNES, 1976 (Pl. 2, Figs 3–6; Pl. 3, Figs 9, 10; Pl. 5, Figs 4–6)

Selected synonymy

- 1970 Pseudoactinoporella fragilis n. gen, n. sp. CONRAD: 66, Text-fig. 4, Pl. I, Figs 1–3; Pl. II; Pl. VIII, Fig. 4.
- 1972 Pseudoactinoporella fragilis Conrad Fourcade et al.: 240, Pl. 3, Figs 7–9.
- 1973 Triploporella sp. SRIVASTAVA: 702, Fig. 21.
- 1974 Pseudoactinoporella fragilis CANÉROT, Pl. XVII, Fig. 2.
- 1976 Pseudoactinoporella fragilis CONRAD CONRAD & PEYBERNES: 188, Fig. 12 (a, b, d-f),
- 1978 Pseudoactinoporella fragilis CONRAD, 1970 BASSOULLET et al.: 212, Pl. 25, Figs 6-8.

1980 Pseudoactinoporella fragilis Conrad – Arnaud-Vanneau, Pl. 111, Figs 1, 2.

- 1989 Pseudoactinoporella fragilis Conrad, 1970 Conrad & Masse: 283, Pl. II, Fig. 8.
- 1994 *Pseudoactinoporella fragilis* CONRAD, 1970 emend CONRAD & PEYBERNES, 1976 BUCUR: 153, Pl. VIII, Figs 9–17.

628

1994 Actinoporella fragilis CONRAD, 1970, n. comb. – GRANIER: 115. [with extended synonymy] 2000 Pseudoactinoporella fragilis CONRAD, 1970 – BUCUR: 60, Pl. IV, Figs 1–6. [with extended synonymy] 2001 Pseudoactinoporella fragilis CONRAD – BUCUR, Pl. 13, Figs 5, 6, 8.

Description and remarks: *Pseudoactinoporella fragilis* is one of the most frequent species in the dasycladalean association from Serre de Bleyton. The detached specimens (Pl. 2, Figs 2–5; Pl. 3, Figs 9, 10) show the general funnel-, or trumpet-like morphology, with a dominantly cylindrical lower side and a splayed upper side. The same morphology is noticeable also in thin sections (Pl. 5, Figs. 4–6). Moreover, the shape of the phloiophorous laterals, shorter in the lower part of the thallus and longer in the upper one is visible. The diameter of the axial cavity is relatively constant along the thallus. Thin sections also give evidence that the small lateral "appendices" connected to the proximal side of the laterals, were located on either their upper, or their lower side. These "appendices" have been noticed by Fourcade et al. (1972), and they were used as emendation criteria by CONRAD & PEYBERNÈS (1976).

GRANIER (1994) has transferred the species *fragilis* to the genus *Actinoporella*, considering that the small secondary laterals used for the genus emendation practically represent elements of a "corona"-type structure. On the other hand, BUCUR (2000) brought forward arguments for maintaining the species in the genus *Pseudoactinoporella*, which are considered by the author to be still valid.

Dimensions

L (maximum observed) = 4.56 mmD (upper part) = 1.4 mmD (lower part) = 0.72-0.80 mmd = 0.28-0.35 mml = 0.25 mmp (distal) = 0.12 mmh = 0.12-0.15 mm

Genus Salpingoporella PIA in TRAUTH, 1918 emend. CARRAS et al., 2006

Recently, CARRAS et al. (2006) presented the species in detail belonging to the genus *Salpingoporella*, including extended synonymy lists and complete descriptions. Accordingly, I will present only brief synonymy-related selections mainly including additional citations, and I will add some general remarks on the three identified species of *Salpingoporella*.

Salpingoporella genevensis (CONRAD, 1969) CONRAD et al., 1973 (Pl. 2, Fig. 7(sg), 8; Pl. 3, Fig. 8; Pl. 5, Figs 10, 11)

Selected synonymy

- 1969 Pianella genevensis n. sp. CONRAD: 571, Figs 1-9
- 1970 Salpingoporella genevensis CONRAD, 1969 JAFFREZO: 79, Pl. 1, Figs 6, 8, 10.
- 1978 Salpingoporella genevensis (CONRAD) 1969 BASSOULLET et al.: 239, Pl. 29, Figs 5-8.
- 1980 Salpingoporella genevensis (CONRAD) ARNAUD-VANNEAU, Pl. 113, Figs 6, 7.
- 1989 Salpingoporella genevensis (CONRAD) 1969 CONRAD & MASSE: 283, Pl. II, Figs 9, 10.
- 1989 Salpingoporella genevensis (CONRAD, 1969) DRAGASTAN: 15, Pl. 6, Figs 1-6.
- 1993 Salpingoporella genevensis LUPERTO SINNI, Pl. 2, Fig. 11.
- 1996 Salpingoporella genevensis CONRAD CLAPS et al., Pl. 2, Figs 4, 7.
- 2000 Salpingoporella genevensis (CONRAD) BUCUR, Pl. 1, Figs 12, 13.
- 2006 Salpingoporella genevensis (CONRAD, 1969) CARRAS et al.: 473, Pl. II, Figs 5–9 [with extended synonymy]
- 2006 Salpingoporella genevensis HUSINEC & SOKAČ, Fig. 5A.
- 2007 Salpingoporella genevensis (CONRAD) BUCUR et al.: 133, Fig. 4(10).

Salpingoporella melitae RADOIČIĆ, 1975

(Pl. 2, Fig. 7(A),?Pl. 5, Fig. 8)

Selected synonymy

- 1967 Salpingoporella melitae n. sp. RADOIČIĆ, Pl. 1, Figs 1–4; Pl. 2, Figs 1, 2; Pl. 3, Figs 1–3; Pl. 4, Figs 1, 2 (nomen nudum: syntypes)
- 1975 Salpingoporella melitae RADOIČIĆ: 277. [designation of a lectotype]
- 1978 *Salpingoporella melitae* RADOIČIĆ, 1967 BASSOULLET et al.: 244, Pl. 29, Figs 16, 17. [with synonymy]
- 1980 Salpingoporella melitae RADOIČIĆ ARNAUD-VANNEAU, Pl. 113, Figs 3-5.
- 1989 Salpingoporella melitae RADOIČIĆ, 1965 CONRAD & MASSE: 283, Pl. II, Fig. 12.
- 1989 Salpingoporella melitae RADOIČIĆ, 1967 DRAGASTAN: 13, Pl. 5, Figs 3, 4.
- 1993 Salpingoporella melitae LUPERTO SINNI & MASSE, Pl. 2, Figs 8-10.
- 1994 Salpingoporella melitae RADOIČIĆ, 1967 BUCUR: 154, Pl. X, Figs 1-7, 10, 11.
- 1996 Salpingoporella melitae CLAPS et al., Pl. 3, Figs 1-6.
- 1996 Salpingoporella melitae RADOIČIĆ, 1967 SOKAČ: 3, Pl. II, Figs 6–13; Pl. III, Figs 1, 2; Pl. XIX, Fig. 7; Pl. XX, Figs 2(B), 3(B). [with selected synonymy]
- 2000 Salpingoporella melitae RADOIČIĆ BUCUR, Pl. I, Figs 7, 8.
- 2001 Salpingoporella melitae RADOIČIĆ BUCUR & COCIUBA, Pl. 3, Figs 3, 4.
- 2001 Salpingoporella melitae RADOIČIĆ BUCUR, Pl. 5, Figs 2, 4.
- 2006 Salpingoporella melitae RADOIČIĆ Ć, 1975 CARRAS et al.: 478, Pl. VI, Figs 6-8.
 - [with extended synonymy]
- 2007 Salpingoporella melitae RADOIČIĆ BUCUR et al.: 133, Fig. 4(7).

Salpingoporella muehlbergii (LORENZ, 1902) PIA in TRAUTH, 1918 (Pl. 2, Fig. 7; Pl. 5, Fig. 7)

Selected synonymy 1902 Diplopora Muehlbergii n. sp. – LORENZ: 52–54, Figs 3–6,?7. 1918 Salpingoporella muehlbergii – PIA in TRAUTH: 211–213, Fig. 4(a).

- 1978 Salpingoporella muehlbergii (LOREZ), 1902 BASSOULLET et al.: 245, Pl. 30, Figs 1-4.
- 1980 Salpingoporella muehlbergii (LORENZ) ARNAUD-VANNEAU: 113, Figs 1, 2.
- 1989 Salpingoporella muehlbergii (LORENZ) 1902 CONRAD & MASSE: 283, Pl. II, Fig. 6.
- 1989 Salpingoporella muehlbergii (LORENZ, 1902) CONRAD, 1970 DRAGASTAN: 13, Pl. 5, Figs. 5-6.
- 1991b Salpingoporella muehlbergii (LORENZ) SCHLAGINTWEIT, Pl. 18, Figs 3–5.
- 1993 Salpingoporella muehlbergii LUPERTO SINNI & MASSE, Pl. 2, Figs 1, 2.
- 1994 Salpingoporella muehlbergii (LORENZ, 1902) emend. CONRAD, 1970 BUCUR: 154, Pl. IV, Fig. 19; Pl. IX, Figs 7–12, 16, 18.
- 1996 Salpingoporella muehlbergii CLAPS et al., Pl. 4, Figs 4, 5.
- 1996 Salpingoporella muehlbergii (LORENZ, 1902) РІА in TRAUTH, 1918 SOKAČ: 2, Pl. I, Figs 1–13; Pl. XX, Figs 4(B0, 5(B).
- 2000 Salpingoporella muehlbergii (LORENZ) BUCUR, P. I, Figs 9–11.
- 2001 Salpingoporella muehlbergii (LORENZ) BUCUR, Pl. 5, Fig. 1.
- 2001 Salpingoporella muehlbergii (LORENZ) BUCUR & COCIUBA, Pl. 3, Figs 1, 2.
- 2006 *Salpingoporella muehlbergii* (LORENZ, 1902) PIA *in* TRAUTH, 1918 CARRAS et al.: 480, Pl. VII, Figs 1–4 [with extended synonymy].
- 2006 Salpingoporella muehlbergii HUSINEC & SOKAČ, Fig. 5(C, D).
- 2008 Salpingoporella muehlbergii (Lorenz, 1902) Pia, 1918 Sokač & Grgasović: 263, Pl. VI, Figs 9–11, Pl. VII.
- 2010 Salpingoporella muehlbergii TAHERPOUR et al., Fig. 7 (f-g)

Remarks: Among the three species, the most frequent one is without doubt *Salpingoporella muehlbergii*. Moreover, it is also one of the most frequent Hauterivian-Aptian dasycladalean algae from the Tethys area. *Salpingoporella melitae* is much scarcer in the association and attempts to section specimens of this species have failed. Nevertheless one can mention the presence of some specimens (Pl. 5, Fig. 8) that according to their sizes can be assigned at the limit between *S. muehlbergii* and *S. melitae*. *Salpingoporella genevensis* is also rarely found in the association; however it is easily recognizable given the quadrangular appearance of its laterals (Pl. 2, Fig. 8, Pl. 3, Fig. 8; Pl. 5, Fig. 10). The close-up view in Pl. 5, Fig. 11 illustrates the aspect of the laterals in tangential section and shows the fingerprints of the reproductive cysts in the calcitic wall around the laterals.

? Genus *Salpingoporella* PIA *in* TRAUTH, 1018, or ? Genus *Neogyroporella* YABE & TOYAMA 1949

?Salpingoporella/?Neogyroporella sp.

(Pl. 1, Figs 9, 10; Pl. 3, Figs 11, 12; Pl. 5, Fig. 9; Pl. 6, Fig. 4; Pl. 7, Fig. 10)

Remarks: This relatively large alga (external diameter between 1.2 and 1.65 mm) shows individualized verticillar discs with laterals that do not seem to alternate from one verticil to the other (a typical feature for *Salpingoporella* species). Moreover, one can notice (Pl. 1, Fig. 9; Pl. 6, Fig. 4) a kind of spacing of verticils along the algal axis. The laterals seem to be phloiophlorous but their exact shape is difficult to observe given a general marginal abrasion affecting the identified specimens. However, the specimen illustrated in Pl. 5, Fig. 9 seems to show more vesiculiform laterals, resembling the typical

Annalen des Naturhistorischen Museums in Wien, Serie A 113

	Kimm.	Tithon.	Berr.	Valan.	Hauterivian	Barremian	Aptian
Deloffrella quercifoliipora							
Actinoporella podolica		_					
Angioporella fouryae							
Falsolikanella danilovae							
Montiella? elitzae							
Piriferella paucicalcarea							
Pseudoactinoporella fragilis							
Salpingoporella genevensis							
Salpingoporella melitae							-
Salpingoporella muehlbergii							

Fig. 3. Generalized stratigraphic rage of ten dasycladalean algae from Serre de Bleyton identified at specific level (based on CONRAD & MASSE 1989; GRANIER & DELOFFRE 1993; BUCUR 1999; CLAVEL et al. 2010).

feature of the species of genus *Neogyroporella* (algae with vesiculiform-type laterals with euspondyl arrangement, as compared to species of the genus *Gyroporella* showing an aspondyl pattern of the vesiculiform laterals (cf. BASSOULLET et al. 1978, p. 172). It is also worth mentioning the relatively large diameter of the axial cavity compared to the external diameter (d/D = 0.60-0.75, for values of the external diameter (D) between 1.2–1.65 mm and of the axial cavity diameter (d) between 0.85–1.05 mm.

Genus Triploporella Steinmann, 1880

Triploporella sp.

(Pl. 3, Figs 5, 6; Pl. 7, Figs 7–9)

Remarks: Some specimens of a large alga (external diameter between 2.1–2.3 mm) show features characteristic of the genus *Triploporella*: verticils consisting of large, tubular laterals of quadrangular-oval section. Within the laterals (Pl. 7, Figs 8, 9) lens-shaped, cyst-containers –type reproductive structures are noticeable, with sizes of 0.2/0.1 mm, the diameter of the cysts being 0.04–0.05 mm. Most probably, the advanced degree of erosion of the external surface is the cause for the lack of secondary laterals located at the distal end of the primary ones. Based on general morphology, sizes, and internal structure, the specimens from Serre de Bleyton can most probably be assigned to *Triploporella praturlonii* BARATTOLO, 1982.

Forma-genus Russoella BARATTOLO, 1984

Russoella radoicicae BARATTOLO, 1984 (Pl. 6, Figs. 9–10)

R e m a r k s: Rare, small-sized (equatorial diameter = 0.17-0.19; axial diameter = 0.12-0.14 mm; diameter of the cysts = 0.04-0.05 mm) lens-shaped corpuscles have been identified in the sediment filling the axial cavity or the inner space of some *Actinoporella* specimen laterals (*e.g.*, Pl. 7, Figs 2, 4). These correspond, morphologically, and in terms of size, to the species *Russoella radoicicae*. Most probably, such corpuscles originate from specimens of *Triploporella* where, inside the laterals, similarly shaped and sized cyst-containers have been identified (*e.g.* Pl. 7. Figs 8, 9).

Discussion

Stratigraphic significance of the dasycladalean assemblage from Serre de Bleyton

Figure 3 illustrates the generalized stratigraphic distribution of the ten dasycladalean species identified at Serre de Bleyton. It is worth mentioning that most of these species have been identified in Barremian deposits from various regions in southern France (CONRAD 1970; CONRAD & PEYBERNES 1976; ARNAUD-VANNEAU 1980; CONRAD & MASSE 1989; MASSE 1993; BUCUR et al. 2007; CLAVEL et al. 2010). The association of species *S. genevensis*, *P. paucicalcarea*, *A. fouryae*, *P. fragilis* and *F. danilovae* is typical for the Late Hauterivian–Early Barremian (cf. CONRAD & MASSE 1989; GRANIER & DELOFFRE 1993; BUCUR 1999; CLAVEL et al. 2010). Most probably, an Early Barremian age can be assigned to the dasycladalean-bearing deposits from Serre de Bleyton.

Paleoenvironmental interpretation

The dasycladalean association from Serre de Bleyton is contained in turbiditic sediments within a basinal succession. The dasycladalean algae have been transported from a carbonate platform margin towards a deeper, basinal setting. Also in talus (slope) deposits they are calibrated by ammonites and echinoids (CLAVEL et al. 2007). CONRAD (1970; 1979) considered that the corresponding species inhabited the median and external area of a carbonate platform. The same type of environment was assumed also for such relatively large dasycladalean by BUCUR & SĂSĂRAN (2005). It is noteworthy that the same species in the association from Serre de Bleyton have been identified in Lower Barremian shallow waters deposits (carbonate platforms) in the entire area around the Vocontian Basin (CONRAD 1970; MASSE 1976, 1993; ARNAUD-VANNEAU 1980; BUCUR et al. 2007; CLAVEL et al. 2007; CLAVEL et al. 2007; CLAVEL et al. 2010).

Annalen des Naturhistorischen Museums in Wien, Serie A 113

Conclusions

The dasycladalean association from Serre de Bleyton consists of 15 taxa, of which 11 have been identified at the species level. The association is typical for peri-Vocontian carbonate platform deposits of early Barremian age, having been transported from shallower waters at the platform margin to deeper, basinal areas.

Acknowledgements

Above all, I would like to thank Gero MOOSLEITNER (Salzburg) who sent me the algal samples and who invited me to contribute to the study of the fossils from Serre de Bleyton. I am also grate-ful to Marc Conrad (Genève), Bruno Granier (Brest) and Felix Schlagintweit (München) for their preliminary remarks on the algae described in this paper. Marc Conrad and Patrick GENOT (Nantes) also supported my research with some litterature – I acknowledge their support also in this way. My colleagues Dana POP (BBU) and Mike KAMINSKI (KFUPM) are thanked for correcting the English version of the text. Part of this study was carried out in Erlangen (Geozentrum Nordbayern, Institut für Paläontologie) with financial support from Alexander von HUMBOLDT Foundation. I thank the A.v. HUMBOLD Foundation as well as the collegues from Erlangen (Roman KOCH – logistic support, Birgit LEIPNER-MATA – thin sections, and Christian SCHULBERT – SEM photos). Thank are also due to my PhD students Camelia CATINCUT and Mihai MICHETIUC for some of the thin sections through the dasycladalean algae.

The reviewers, Marc CONRAD and Bruno GRANIER are grattefully acknowledge for their final remarks on the paper, as well as the editor, Andreas KROH for his remarks and help with figure 1.

References

- ALTH, A. (1878): O Galicyjskich gatunkach skamieniałych otwornic rodzaju *Gyroporella* GÜMB. Rozprawy i Sprawozdania z Posiedzeń Wydziału Matematyczno-Przyrodniczego Akademii Umiejętności, **5**: 71–112.
- (1882): Versteinerungen des Nizniower Kalksteines. Beiträge zur Paläontologie Österreich-Ungarns und des Orients, 1: 183–332.
- ARBNAUD-VANNEAU, A. (1980): Micropaléontologie, paléoécologie et sédimentologie d'une plateforme carbonatée de la marge passive de la Tethys: L'Urgonien du Vercors septentrional et de la Chartreuse (Alpes Occidentales). – Géologie Alpine, Mémoire **11**, 874 p., Grenoble
- AZEMA, J., FOUCAULT, A., FOURCADE, E., GARCIA-HERNANDEZ, M., GONSALES-DONOSO, J. M., LINARES, A., LINARES, D., LOPEZ-GARPIDO, A. C., RIVAS, P. & VERA, J.A. (1979): Las Microfacies del Jurassico y Cretacico de las zonas externas de las Cordilleras Beticas. 83 p. Universidad de Granada.
- BAKALOVA, D. (1971): Nouvelles espèces de Dasycladaceae (Algae) dans les sédiments urgoniens du Prébalkan central. – Bulletin of the Geological Institute, Series Paleontology, **20**: 123–128.
- BARATTOLO, F. (1978): Su di una nuova dasicladacea (alghe verdi) nel Paleocene dell'Appennino meridionale. Bolletino della Società dei Naturalisti in Napoli, **87**: 1–76.

- -----(1982): Oservazioni su *Triploporella praturlonii* n. sp. (Alghe verdi, dasicladali) del Cretacico infeirore dell'Appennino centrale. Geologica Romana, **21**: 825–847.
- (1984): Su alcuni organi riproduttori di Dasicladacee nel Cretacico dell'Appennino. Bollettino della Società dei Naturalisti in Napoli, 92 (1983): 1–33.
- BASSOULLET, J.P., BERNIER, P., CONRAD, M.A., DELOFFRE, R. & JAFFREZO, M. (1978): Les Algues Dasycladales du Jurassique et du Crétacé. Geobios, Mémoire spécial **2**, 330 p., Lyon
- BUCUR, I.I. (1993): Some new or poorly known calcareous algae (Dasycladales, Gymnocodiaceae) in the Lower Cretaceous deposits from the Reşiţa-Moldova Nouă zone (Southern Carpathians, Romania). – Revista Española de Micropaleontologia, **25**/1: 93–126
- (1994): Algues calcaires de la zone de Reşiţa-Moldova Nouă (Carpathes Méridionales, Roumanie). – Revue de Palobiologie, 13/1: 147–209.
- (1999): Stratigraphic significance of some skeletal algae (Dasycladales, Caulerpales) of the Phanerozoic. – In: FARINACCI, A & LORD, A.R. (eds): Depositional episodes and Bioevents. Palaeopelagos Special Publication nº 2: 53–104
- (2000): Lower Cretcaeous dasyclad algae from Pădurea Craiului massif (Northern Apuseni Mountains, Romania) – Acta Palaeontologica Romaniae, 2 (1999): 53–72.
- (2001): Lower Cretaceous algae of the Reşiţa-Moldova Nouă zone. In: BUCUR, I.I., FILIPESCU, S. & SĂSĂRAN, E. (eds): Algae and carbonate platforms in the western part of Romania. 4th Regional Meeting of IFAA, Cluj-Napoca, Romania, August 29–September 5, 2001. Field trip guidebook. – pp. 137–166, Cluj-Napoca (Cluj-University Press).
- (2002): Nouvelles découvertes de dasycladales fossiles dans les dépôts du Crétacé inférieur de la zone de Reşiţa-Moldova Nouă (Carpathes Méridionales, Roumanie). – Acta Palaeontologica Romaniae, 3: 41–52.
- —, ARNAUD-VANNEAU, A. & ARNAUD, H. (2007): Remarks on *Triploporella praturlonii* Barattolo 1982 from the Lower Barremian of Southern Vercors (SE France). Neues Jahrbuch für Geologie unfd Paläontologie, Abhandlungen, 244/2: 133–141.
- & COCIUBA, I. (2001): Lower Cretaceous algae of Pădurea Craiului. In: BUCUR, I.I., FILIPESCU, S. & SĂSĂRAN, E. (eds): Algae and carbonate platforms in the western part of Romania. 4th Regional Meeting of IFAA, Cluj-Napoca, Romania, August 29–September 5, 2001. Field trip guidebook. – pp. 119–127, Cluj-Napoca (Cluj-University Press).
- --- & Săsăran, E. (2005): Relationship between algae and environment: an Early Cretaceous case study, Trascău Mountains, Romania. Facies, **51**: 274–286.
- ----, SENOWBARI-DARYAN, B & MAJIDIFARD, M.R. (2003): Neocomian microfossil association from the Taft area near Yazd (Central Iran). Facies, **49**: 217–222.
- CANÉROT, J. (1974): Recherches géologiques aux confins des chaînes ibériques et catalanes (Espagne). Thèse, 517 p., Toulouse.
- CARRAS, N., CONRAD, M.A. & RADOIČIĆ, R. (2006): *Salpingoporella*, a common genus of Mesozoic Dasycladales (calcareous green algae). Revue de Paléobiologie, **25**/2: 457–617.
- CLAPS, M.P., PARENTE, M., NERI, C. &, BOSELLINI, A. (1996): Facies and Cycles of the S. Giovanni Rotondo Limestone (Lower Cretaceous, Gargano Promontory, Southern Italy): The Borgo Celano Section. – Annali dell'Università di Ferrara, 7 (suppl.): 5–35.

- CLAVEL, B., BUSNARDO, R., CHAROLLAIS, J., CONRAD, M.A. & GRANIER, B. (2010): Répartition biostratigraphique des orbitolinidés dans la biozonation à ammonites (plate-forme urgonienne du Sud-Est de la France). Partie 1: Hauterivien supérieur – Barrémien basal. – Carnets de Géologie/Notebooks on Geology, Article 2010/06 (CG2010_A06), 53 p.
- —, CHAROLLAIS, J., CONRAD, M., JAN DU CHÊNE, R., BUSNARDO, R., GARDIN, S., ERBA, E., SCHROEDER, R., CHERCHI, A., DECROUEZ, D., GRANIER, B., SAUVAGNAT, J. & WEIDMANN, M. (2007): Dating and progradation of the Urgonian limestone from the Swiss Jura to South-East France. [Datierung und Progradation des Urgons vom Schweizer Jura bis Südost-Frankreich.] – Zeitschrift der deutschen Gesellschaft für Geowissenschaften, 158/4: 1025–1062.
- CONRAD, M.A. (1969): *Pianella genevensis* n. sp., une dasycladacée cladospore des calcaires urgoniens. Eclogae Geologicae Helvetiae, **62**/2: 571–582.
- (1970): Barremian and Lower Aptian Dasycladaceae in the area surrounding Geneva (Switzerland). – Geologica Romana, 9: 63–100.
- (1979): The Lower Cretaceous algae in the area surrounding Geneva (Switzerland): biostratigraphy and depositional environments. In: FLÜGEL, E. (ed.): Fossil Algae. Recent results and developments. pp. 295–300, Berlin (Springer Verlag).
- & MASSE, J.-P. (1989): Les Algues calcaires des formations carbonatées de l'Hautérivien-Barrémien pro parte du Jura vadois et neuchâtelois (Suisse). – Mémoires de la Société neuchâteloise des Sciences naturelles, 11: 277–290.
- & PEYBERNÈS, B. (1976): Hauterivian-Albian Dasycladaceae from the Urgonian limestones in the French and Spanish Pyrénées. – Geologica Romana, 15: 175–197.
- —, PRATURION, A. & RADOIČIĆ, R. (1973): Reinstatement of the genus Salpingoporella PIA (Dasycladaceae), followed by Salpingoporella steinhauseri, n. sp. – Comptes Rendus des Séances, SPHN Genève, 7/2–3:103–111.
- ----, PRATURLON, A. & RADOIČIĆ, R. (1974): The genus Actinoporella GÜMBEL in ALTH 1882, Dasycladales, green algae. A revision. Geologica Romana, 13: 1–15.
- —, SCHLAGINTWEIT, F. & BUCUR, I.I. (2009): *Piriferella somalica* (Dasycladalean calcareous green algae) from Tithonian deposits in the Northern Calcareous Alps synonymy followed by an emendation of *Piriferella* SOKAČ 1998 and *Similiclypeina* BUCUR 1993. In: BASSO, D., CARAGNANO, A., BROCCHI, V. & BENZANI F. (eds) Abstract book of the 6th Regional Symposium of IFAA, Milano, Italy, 1–5 sept. 2009. Annali dell'Università di Ferrara, Sez. Museologia Scientifica e Naturalistica, Volume speciale 2009.
- DRAGASTAN, O. (2000): Early Cretaceous algae of Aliman (Southern Dobrogea): a revision and description of two new species from East Carpathians. – Acta Palaeontologica Romaniae, 2 (1999): 125–137.
- & BUCUR, I.I. (1988): Comments on the genus *Radoiciciella*. Analele Universității București, Geologie, 37: 97–99.
- ---- & BUCUR, I.I. (1993): The dasyclad genus *Radoiciciella* and its representatives from Romania. Revista Española de Micropaleontologia, **25**/2: 5–23
- FOURCADE, E., JEREZ, L., RODRIGUEZ, T. & JAFFREZO, M. (1972): El Jurasico terminal y el Cretacico inferior de la Sierra de la Muela (Provincia de Murcia). Consideraciones sobre las biozonas con foraminiferas del Albense-Aptense del Sureste de España. – Revista Española de Micropaleontologia, Numero. Extraordinario XXX Aniversario E.N. Adaro: 215–248.

- GRANIER, B. (1987): Révision de *Likanella campanensis* AZEMA, et JAFFREZO, 1972, algue dasycladacée du Crétacé inférieur du sud-est de l'Espagne. Revue de Paléobiologie, **6**/2: 207–212.
- (1990): The case of the genus *Radoiciciella*, Dasycladalean Algae from the Tethyan Upper Jurassic and Lower Cretaceous. – Comunicações dos Serviços geológicos de Portugal, 75: 29–37.
- (1994): The genus *Actinoporella* (GÜMBEL in ALTH, 1881) and its representatives. A review.– Proceedings of the International Symposium and Field-Meeting "Alpine Algae '93". – Beiträge zur Paläontologie, **19**: 13–127.
- (2008): Holostratigraphy of the Kahmah regional series in Oman, Quatar and the United Arab Emirates. – Carnets de Géologie/Notebooks in Geology, Article 2008/07 (CG 2008_A07), 33 p.
- & DELOFFRE, R. (1993): Inventaire critique des algues dasycladales fossiles. II^o partie Les algues dasycladales du Jurassique et du Crétacé. Revue de Paléobiologie, **12**/1: 19–65.
- —, BUCUR, I.I. & TRABOLD, G. (2000): Falsolikanella danilovae RADOIČIĆ ex BARATTOLO 1978, n. comb., a diploporacean alga from the Urgonian facies. – Acta Palaeontologica Romaniae, 2 (1999): 177–181.
- & MICHAUD, F. (1987): Deloffrella quercifoliipora n. gen., n. sp., une algue dasycladacée nouvelle du Kimméridgien et du Portlandien du Sud-Est du Mexique. – Bulletin de la Societé géologique de France, (8), 3/6: 1089–1096.
- HOSSEINI, S.A. & CONRAD, M.A. (2008): Calcareous algae, foraminifera and sequence stratigraphy of the Fahliyan Formation at Kuh-e-Surmeh (Zagros basin, SW of Iran). – Geologia Croatica, 61/2–3:215–237.
- HUSINEC, A. & SOKAČ, B. (2006): Early Cretaceous benthic associations (foraminifera and calcareous algae) of a shallow tropical-water platform environment (Mljet Island, southern Croatia). Cretaceous Research, 27: 418–441.
- JAFFREZO M. (1973): Les algues calcaires du Jurassique supérieur et du Crétacé inférieur des Corbières. Première partie. Revue de Micropaléontologie, **16**/2: 75–88.
- JANSSEN, N.M.M. (2010): Barremian invertebrates from Serre de Bleyton (Drôme, SE France): Belemnites. Annalen des Naturhistorichen Museums Wien, Serie A, **112**: 659–672.
- JÄGER, M. (2010): Crinoids from the Barremian (Lower Cretaceous) of the Serre de Bleyton (Drôme, SE France). Annalen des Naturhistorichen Museums Wien, Serie A, **112**: 733–774.
- KROH, A., LUKENDER, A & MOOSLEITNER, G. (2010): The invertebrate fauna from the Barremian of Serre de Bleyton (Drôme, SE France). – Annalen des Naturhistorichen Museums Wien, Serie A, 112: 569–574.
- LORENZ, T. (1902): Geologische Studien im Grenzbebiete zwischen helvetischer und ostalpiner Facies. – Berichte der naturforschenden Gesellschaft zu Freiburg im Breisgau, **12**: 34–62.
- Löser, H. (2010): The Barremian coral fauna of the Serre de Bleyton mountain range (Drôme, SE France). Annalen des Naturhistorichen Museums Wien, Serie A, **112**: 575–612.
- LUKENDER, A. (2010): Barremian ammonoids from Serre de Bleyton (Drôme, SE France). Annalen des Naturhistorichen Museums Wien, Serie A, **112**: 613–626.

- LUPERTO-SINNI, E. & MASSE, J.-P. (1993): The Early Creataceous dasycladales from the Apilia region (Southern Italy): biostratigraphic distribution and paleobiogeographic significance. – In: BARATTOLO, F. et al. (eds): Studies on fossil benthic algae – Bolletino della Società Paleontologica Italiana, Special volume 1: 295–309.
- MANCINELLI, A. (1992): Distribuzione delle dasycladali (Alghe verdi) nei sedimenti del Cretacico inferiore in facies di Piattaforma carbonatica dell'Appenino centro-meridionale (Lazio e Abruzzo). Studi Geologici Camerti, **12**: 7–29.
- MASSE, J.-P. (1976): Les calcaires urgoniens de la Provence (Valanginien-Aptien). Stratigraphie, Paléontologie, les paléoenvironnments et leur évolution. – Thése Doctorat d'Etat, Université Aix-Marseille II: 445 p.
- (1993): Early Cretaceous Dasycladales biostratigraphy from Provence and adjacent regions (South of France, Switzerland, Spain). A reference for Mesogean correlations. – In: BARATTOLO F. et al. (eds): Studies of fossil benthic algae – Bolletino della Società Paleontologica Italiana, Special Volume 1: 311–324.
- —, CONRAD, M.A. & RADOIČIĆ, R. (1973): Angioporella fouryae n. gen., n. sp., une algue calcaire (dasycladaceae) du Barrémien du sud de la France. – Eclogae Geologicae Helvetiae, 66/2: 383–387.
- MICHELIN, H. (1840–1847): Iconographie zoophytologique. Description par localités et terrains des Polypiers sossiles de France et pays environnants. 384 p., Paris (Bertrand ed.).
- MORELLET, L. & MORELLET, J. (1922): Nouvelle contribution ă l'étude des dasycladacées tertiaires. Mémoires de la Société géologique de France, Paléontologie, **58**: 5–33.
- PIA, J. von (1918): Dasycladaceae. In: TRAUTH, F. (ed.). Das Eozänvorkommen bei Radstadt im Pongau und seine Beziehungen zu den Gleichalterigen Ablagerungen bei Kirchberg am Wechsel und Wimpassing am Leithagebirge. Denkschriften der Kaiserliche Akademie der Wissenschaften in Wien, Matematisch-naturwissenschaftliche Klasse, 95: 209–213.
- RADOIČIĆ, R. (1967): *Salpingoporella melitae* sp. nov. des sédiments crétacés inférieurs des Dinarides externes. Annales Géologiques de la Peninsule Balkanique, **33**: 121–126.
- (1968): Likanella? danilovae spec. nov. et quelques autres Dasycladacées crétacées inférieures des Dinarides externes. – Vesnik (Geologija), Ser. A, 26: 177–193.
- (1970): The new dasycladacean genus Pseudoclypeina (a preliminary report). Bulletin Scientifique, Section A: 15/1–2.
- (1975): Linoporella buseri sp. nov. from the Liassic of the Julian Alps (a preliminary report). Bulletin Scientifique, Conseil des Académies des Sciences et des Arts de la RSF de Yougoslavie, Section A: Sciences Naturelles, Techniques et Médicales, 20/9–10: 277–278.
- (1980): Contribution to the reconsideration of s group of cylindroporelliform dasyclads. Bulletin de l'Académie serbe des sciences et des arts, classe des sciences naturelles et mathématiques, **20**: 109–115.
- RAVIV, V. &, LORCH, J. (1992): Mesozoic Dasycladaceae (Chlorophyceae) from Israel. Israel Journal of Botany, **41**/1: 1–54.
- RIEGRAF, W. & MOOSLEITNER, G. (2010): Barremian rhyncholites (Lower Cretaceous Ammonoidea: calcified upper jaws) from the Serre de Bleyton (Départment Drôme, SE France). – Annalen des Naturhistorichen Museums Wien, Serie A, 112: 627–658.

- SCHLAGINTWEIT, F. (1991a): Neritische Oberjura- und Unterkreide Kalkgerölle aus den Losensteiner Schichten (Alb – Cenoman) der Typlokalität Stiedelsbachgraben (Oberosterreich; Nordliche Kalkalpen). – Mitteilungen der Gesellschaft der Geologie- und Bergbaustudenten in Österreich, 37: 83–95.
- (1991b): Allochthone Urgonkalke im Mittleren Abschnitt der Nordlichen Kalkalpen: Fazies, Paläontologie und Paläogeographie. – Munchner Geowissenschaftliche Abhandlungen (A), 20: 1–119.
- (2005): Neogyroporella? gawlicki n. sp., a new dasycladale from the Upepr Jurassic-Lower Cretaceous "Lärchberg Formation" of the Northern Calcareous Alps, Austria. – Geologia Croatica, 58/2: 103–117.
- —, GAWLICK, H.-J., MISSONI, S., LEIN, R. & HOXHA, L. (2006): Late Jurassic to Early Cretaceous Dasycladales and benthonic foraminifera from the Munella Carbonate Platform s.l. of the Mirdita zone (Albania). – In: SUDAR, M., ERCEGOVAC, M. & GRUBIĆ, A. (eds) – XVIIIth Congress of the Carpathian-Balkan Geological Association, Belgrad 2006, Proceedings: 527–530.
- SOKAČ, B. (1996): Taxonomic review of some Barremian and Aptian Calcareous Algae (Dasycladales) from the Dinaric and Adriatic Karst Regions of Croatia. Geologia Croatica, **49**/1: 1–79.
- & GRGASOVIČ, T. (2008): on some Salpingoporella species from the Lower Cretaceous of Dinaric Karst. – Geologia Croatica, 61/2–3: 251–271.
- SRIVASTAVA, N.K. (1973): Neocomian calcareous algae from Bolshoi Balkan, USSR. Neues Jahrbuch für Geologie und Paläontologie, Monatshefte, 11: 641–708.
- STEINMANN, G. (1880): Zur kenntnis fossiler Kalkalgen (Siphoneen). Neues Jahbuch für Mineralogie, Geologie und Paläontologie, 2: 130–140.
- TAHEPOUR KHALIL ABAD, M., CONRAD, M.A., ARYAEI, A.A. & ASHOURI, A.R. (2010): Barremian-Aptian Dasycladalean algae new and revisited, from the Tirgan Formation in thr Kopet Dagh, NE Iran. – Carnets de Géologie/Notebooks on Geology, Article 2010/05 (CG 2010_A05), 13 p.
- TAYLOR, P. D. (2010): Barremian bryozoans from Serre de Bleyton (Drôme, SE France) Annalen des Naturhistorichen Museums Wien, Serie A, **112**: 673–700.
- YABE, H. & TOYAMA, S. (1949): New Dasycladaceae from the Jurassic Torinosu Limestone of the Sakawa Basin. II. Proceedings of Japan Academy, **25**/7: 160–164.

Isolated specimens of dasycladalean algae from Serre de Bleyton. Stereomicroscope photomicrographs.

- Fig. 1. Dasycladalean assemblage dominated by *Actinoporella* (A) specimens. Disc fragments of *Pseudoclypeina* sp. (Pc) are also present.
- Figs 2–5. *Actinoporella podolica* (ALTH, 1878). Isolated verticils of laterals showing corona-like protuberances made up by the small secondary laterals around the central stem, and the phloiophorous large secondary laterals. Fig. 3 represents an enlargement of the specimen in the lower-central part of Fig. 2, and Fig. 5 an enlargement of the upper part of Fig. 2.
- Fig. 6. Piriferella paucicalcarea (CONRAD, 1970). View of the external surface.
- Fig. 7. Assemblage dominated by *Piriferella paucicalcarea* (CONRAD, 1970). Most diagnostic is the broken specimen in the center-right part of the photo.
- Fig. 8. Fragments of verticillar discs of *Pseudoclypeina* sp. The fragments originating from different specimens are arranged to suggest the shape of a verticil.
- Figs 9, 10. Unidentified dasycladalean alga (*?Salpingoporella-?Neogyroporella*). See the thin section through the specimen from Fig. 9 in Pl. 6, Fig. 4.



Isolated specimens of dasycladalean algae from Serre de Bleyton. Stereomicroscope photomicrographs

- Figs 1–2. Angioporella fouryae MASSE, CONRAD & RADOIČIĆ, 1973. Assemblage with isolated specimens artificially corroded (with acetic acid) showing the successive alternance of sterile and fertile verticils (Fig. 1); a small specimen in external view showing two verticils of sterile laterals and the thin vertical "bulges" between the vertical parts of the sterile laterals covering the fertile ampoulae.
- Figs 3–6. *Pseudoactinoporella fragilis* CONRAD, 1970. Assemblages with isolated specimens in figs. 3 and 4; Fig. 5 shows the external morphology of the specimen in the lower-central part of Fig. 4, and Fig. 6 shows the upper part of the same specimen.
- Fig. 7. Assemblage with isolated specimens of Salpingoporella. Most specimens belong to Salpingoporella muehlbergii (LORENZ, 1902). Rare Salpingoporella genevensis (CONRAD, 1969) (Sg) and Salpingoporella melitae RADOIČIĆ, 1975 (A) are also present.
- Fig. 8. Salpingoporella genevensis (CONRAD, 1969). External view of an isolated specimen.
- Figs 9–12. *Falsolikanella danilovae* (RADOIČIĆ). Assemblage dominated by relatively small specimens (see enlargements in figs 11 and 12). Fig. 10 presents an external view of a larger annular segment of laterals.
- Fig. 13. *Piriferella paucicalcarea* (CONRAD). Close-up view of the specimen in Pl. 1, Fig. 7 (center-right).



SEM photomicrographs of dasycladalean algae from Serre de Bleyton.

- Figs 1–3. *Actinoporella podolica* (ALTH, 1878). Corona-like structure developed by the small secondary laterals is clearly visible on both sides of the verticillar discs.
- Fig. 4. Angioporella fouryae MASSE, CONRAD & RADOIČIĆ, 1973. Artificially slightly corroded specimen showing the verticils of thin sterile laterals surrounding the larger fertile laterals.
- Figs. 5–6. *Triploporella* sp. Specimen in two different views showing only the first-order laterals.
- Fig. 7. *Falsolikanella danilovae* (RADOIČIĆ, 1978). Annular segment made up of divergent arranged laterals.
- Fig. 8. Salpingoporella gr. genevensis (CONRAD, 1969). Broken specimen showing the arrangement of laterals.
- Figs 9–10. *Pseudoactinoporella fragilis* CONRAD, 1970. Lateral (Fig. 9, upper specimen and Fig. 10), and upper side (Fig. 9, lower specimen) view showing the characteristic enlargement of the thallus in the upper part, and the arrangement of the laterals in a verticil, respectively.
- Figs 11, 12. Unidentified dasycladalean algae (*?Salpingoporella-?Neogyroporella*). See the thin section through the specimen from Fig. 11 in Pl. 7, Fig. 10.



Thin sections through isolated specimens of dasycladalean algae from Serre de Bleyton.

- Figs 1–6. *Actinoporella podolica* (ALTH, 1878). Transverse sections through isolated verticils. The corona-like structure is very well visible in Figs 1-3, but in less visible, even if still present, in the other specimens (Figs 4-6).
- Figs 7–11. Angioporella fouryae MASSE, CONRAD & RADOIČIĆ, 1973. Figs 7–9, longitudinal (7) and longitudinal-oblique (8, 9) sections through corroded specimens showing the alternate arrangement of sterile and fertile laterals. Fig. 10 illustrates a longitudinal-oblique section through a non-corroded specimen. The possible connection between the fertile laterals and the main stem is visible in the case of the upper two verticils (left side). Fig. 11 is a transverse section through a verticil of sterile laterals.
- Figs 12–14. *Falsolikanella danilovae* (RADOIČIĆ, 1978). Thin sections through isolated annular segments.
- Figs 15–17. *Pseudoclypeina* sp. Thin sections through fragments of disc-like verticils. Long primary laterals, and shorter secondaries are visible.



Thin sections through isolated specimens of dasycladalean algae from Serre de Bleyton.

- Figs 1–3. *Pseudoclypeina* sp. Thin sections through fragments of verticillar discs showing the primary and secondary laterals.
- Figs 4–6. *Pseudoactinoporella fragilis* CONRAD, 1970. Longitudinal-oblique sections showing the enlargement of the thallus in the upper part, as well as the shape and arrangement of the laterals.
- Fig. 7. Salpingoporella muehlbergii (LORENZ, 1902). Longitudinal section.
- Fig. 8. *Salpingoporella* gr. *muehlbergii* (LORENZ, 1902) melitae Radoičić. Transverse section.
- Fig. 9. Unidentified dasycladalean alga (?Salpingoporella-?Neogyroporella). Transverse section.
- Figs 10, 11. Salpingoporella genevensis (CONRAD, 1969). Tangential section. Fig. 11 is a close-up view of the left side of Fig. 10 showing the characteristic morphology of the laterals in tangential section, as well as the imprint of reproductive cysts.



Thin sections through isolated specimens of dasycladalean algae from Serre de Bleyton.

- Fig. 1. *Deloffrella quercifoliipora* GRANIER & MICHAUD, 1987. Longitudinal-oblique section showing the typical morphology of the laterals.
- Figs 2, 3. *Montiella? elitzae* (BAKALOVA, 1971). Longitudinal-tangential sections showing the position of the fertile and sterile secondary laterals.
- Fig. 4. Unidentified dasycladalean alga (?Salpingoporella-?Neogyroporella). Longitudinal-tangential section through the specimen in Pl. 1, Fig. 9.
- Figs 5, 8. *Piriferella paucicalcarea* (CONRAD, 1970); oblique (5) and tangential (8) sections.
- Figs 6, 7, 11. *Actinoporella podolica* (ALTH, 1878). Transverse (horizontal) sections through isolated verticillar discs with a large number of laterals partly overlapping each other.
- Figs 9, 10. *Russoella radoicicae* BARATTOLO, 1984. Small reproductive structures deriving most probably from *Triploporella* specimens.
- Figs 12, 13. *Clypeina* sp. Transverse, slightly oblique (Fig. 12) section, and oblique-tangential (Fig. 13) section.



Thin sections through isolated specimens of dasycladalean algae from Serre de Bleyton.

- Figs 1–6. Actinoporella podolica (ALTH, 1878). Transverse (horizontal) sections (1, 2, 4, 5), longitudinal (vertical) section (3), and longitudinal-obique (vertical-oblique) section (6) through isolated verticils of laterals. The corona-like structures are visible in figs 3 and 6.
- Figs 7–9. *Triploporella* sp. Transverse (7) and longitudinal-tangential (8) sections (Fig. 8 is a thin section of the specimen illustrated in Pl. 3, Fig. 6). Fig. 9 is a close-up view of the lower-right part of Fig. 8, showing the presence of *Russoella*-like reproductive structures inside the laterals.
- Fig. 10. Unidentified dasycladalean alga (?*Salpingoporella*-?*Neogyroporella*). Thin sections of the specimen illustrated in Pl. 3, Fig. 11.



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Annalen des Naturhistorischen Museums in Wien

Jahr/Year: 2011

Band/Volume: 113A

Autor(en)/Author(s): Bucur Ioan I.

Artikel/Article: <u>Early Barremian dasycladalean algae from Serre de Bleyton (Drôme, SE France). 619-653</u>