Ann. Naturhistor. Mus. Wien

Othonna rechingeri B. NORD. spec. nova, a hexaploid succulent from South Africa

By BERTIL NORDENSTAM ¹)

Mit 1 Textabbildung

Manuskript eingelangt am 2. Februar 1971

Abstract

Othonna rechingeri B. NORD. (Compositae-Senecioneae-Othonninae) is described from Calvinia Division, Cape Province, South Africa. The new species is hexaploid with the chromosome number 2n = 60. The closely related species O. patula SCHLTR. is diploid with 2n = 20.

Introduction

In connection with the Kirstenbosch Golden Jubilee celebrations in 1963, my highly esteemed friend, Professor KARL-HEINZ RECHINGER, paid a visit to South Africa. In addition to the official excursion program we made a private collecting tour together, i. a. to some of the most interesting parts of the Vanrhynsdorp and Calvinia Divisions. Among the many rare plants found, was a remarkable, undescribed species of *Othonna*, which I have great pleasure in naming after my fellow traveller on that memorable excursion.

Othonna rechingeri B. Nord., sp. nov.

Orig. coll.: NORDENSTAM 765, Cape Province, Calvinia Division, Klip Koppies, c. 2 miles E. of Nieuwoudtville, 27. VII. 1962 (S holotype; isotypes distributed to BOL, K, PRE and W).

Illustr.: Fig. 1.

O. patulae SCHLTR. affinis, sed foliis angustioribus longioribus, capitulis discoideis differt.

Herba perennis glabra; caulis foliatus erectus brevis e caudice tuberoso basali exiens, apice stolones elongatos et pedunculum scapiformen emittens. Tuber parvum partim subterraneum crassum viride vel brunneolum. Stolones pauci horizontales sensim arcuati graciles elongati foliati. Folia alterna sparsa, in apice caulis valde conferta, erecta sessilia linearia teretia carnosa laevia viridia apice rotundata mucronulata. Pedunculus erectus (repetite) bifurcatus scapiformis nudus sed bracteis reductis et interdum 1-2 foliis basalibus instructus, ramis capituliferis arcuatim nutantibus. Capitula hemi-

¹) Museum of Natural History, Stockholm 50, Sweden.

B. NORDENSTAM

sphaerica lutiflora discoidea heterogama. Involucri bracteae 8 basi connatae oblongae – anguste ovatae. Flores marginales 15-20 hermaphroditi; corolla basi tubulosa, superne campanulata; stylus bilobatus; pappi setae copiosae caducae albae; achenium oblongo-obovatum – anguste oblongum apice albo-annulatum, indumento corticate appresso argenteo-cinereo vel brunneolo, humido mucoso. Flores centrales 30-50 feminei-steriles; corolla basi tubulosa, superne campanulata; stylus simplex sterilis apice appendice



Fig. 1. Othonna rechingeri B. NORD. (NORDENSTAM 765). — A: Habit, $\times \frac{1}{2}$. — B: Marginal floret, $\times 5$. — C: Style apex, from marginal floret, $\times 20$. — D: Achenes, $\times 5$. — E: Central floret, $\times 5$. — F: Style apex, from central floret, $\times 20$. — G: Stamens, from central floret, $\times 10$. — Del. auct.

conica instructus; ovarium anguste oblongum glabrum abortivum; pappi setae paucae caducae albae.

A glabrous *perennial herb* with erect flowering stems from a basal tuber, emitting long, leafy stolons. *Tuber* rounded, partly underground and with fibrous roots from the base, 1-2 cm in diam., firm, green or brownish. *Stem* with an erect or decumbent, short (up to a few cm long), leafy portion and an elongated erect peduncle, terete, fleshy. *Stolons* 1-5 (seldom more) from the apex of the leafy stem, spreading horizontally, gently arching, slender, up

Othonna rechingeri B. NORD. spec. nova, a hexaploid succulent from South Africa 141

to 20 cm long (rarely longer), ending in rooting tubers developing new stems and leaves. Leaves alternate, laxly set or somewhat crowded at the stem apex, erect, sessile, linear, terete, fleshy, smooth, light green, 1-10 cm long, 1,5-4 mm thick (shortest on the stolons, longest and thickest near the stem apex), apically rounded and mucronulate. Peduncle 10-30 cm long, forking once or 2-3 times, with leaf-like bracts at the ramifications, otherwise leafless or with 1-2 distant leaves, with arching-nodding capituliferous branches. Capitula hemispherical, 1-1,5 cm in diam., yellow-flowered, discoid, heterogamous with tubular perfect marginal florets and tubular male (Q-sterile) central florets. Receptacle slightly convex, shallowly alveolate. Involucral bracts uniseriate, 8, basally connate, oblong – narrowly ovate, 6-8 mm long, 2-4 mm wide, \pm herbaceous, many with broad, membranous margins, acute or obtuse, indistinctly many veined or with 3-5 purplish lines, sometimes purplish tipped. Marginal florets c. 15-20. Corolla c. 4 mm long; tube 1,5 mm long, basally slightly inflated; limb narrowly companulate, 2,5 mm long; lobes ovate-deltoid, 0,6-0,7 mm long, the tips somewhat cucullate, acute, with minutely papillate exteriors. Style slightly flattened, with swollen base; style branches c. 1 mm long, flattened, apically tipped with an acute, triangular appendage surrounded by short sweeping-hairs. Anthers 1,6-1,8 mm long incl. the ovate, obtuse appendage. Pappus bristles copious (c. 100-150), 3-4 mm long, caducous, erect, straight, white. Achene oblong-obovate narrowly oblong, 2,5-3 mm long, 1 mm wide, faintly ribbed or subterete, coated with a densely appressed, silvery grey - brownish hair-coat, becoming mucilaginous when soaked, apically with a distinct white annulus. Central florets c. 30-50. Corolla 4,5 mm long; tube 1,5 mm long, cylindrical; limb narrowly campanulate, 3 mm long; lobes deltoid, 0,7-0,8 mm long. Anthers 1,8-2 mm long incl. the ovate, obtuse appendage; filaments distinctly swollen apically. Style simple, sterile, tipped with a blunt conical appendage; cone 0,3 mm long, with short sweeping-hairs at the base. Ovary narrowly oblong, 1,5 mm long, glabrous, c. 5-veined, abortive. Pappus bristles few (c. 5-15), erect, straight, white, caducous.

Flowering period: July-Sept.

Chromosome number: 2n = 60 (in sections of root-tips from NORDEN-STAM 765 and 3017, modified Navashin's fixative, paraffin method, stain crystal violet).

Collections: Cape Province, Calvinia Division: Papelfontein, 2200 ft., VIII. 1897, SCHLECHTER 10908 (COI, G in herb. Barbey-Boissier, S) — Flats below Klip Koppies, c. 2 miles E. of Nieuwoudtville, VII. 1962, NORDENSTAM 765 (BOL, K, PRE, S, W) — Same locality, IX. 1963, NORDENSTAM 3017 (S).

Othonna rechingeri B. NORD. belongs to sect. Carnosae HARV. and is no doubt closely related to O. patula SCHLTR. These two species deviate from all others by their peculiar, rooting stolons. The runners evidently serve as an effective means of vegetative propagation. I have had both species in cultivation for several years, and I found that they easily spread to other pots. When the stolon tip comes into contact with soil, a small tuber is formed, which rapidly roots and develops to a new individual.

O. rechingeri and O. patula are very similar in their vegetative parts, the former, however, having slightly longer and narrower leaves. The chief difference is found in the flowerheads, which are radiate in O. patula, but discoid in O. rechingeri. In other words, the two species have quite different types of marginal florets. These are normal, ligulate, female ray-florets in O. patula, but in O. rechingeri their place is taken by perfect tubular florets. The latter type is rare in the genus, occurring only in a few taxa of sect. Carnosae, viz., O. rechingeri, O. carnosa var. discoidea, and O. sparsiflora (cf. NORDENSTAM 1961, p. 125). Most discoid taxa of the genus (formerly recognized as the genus Doria LESS.) have reduced, more or less tubular, female marginal florets, which are clearly derived from normal ray-florets.

The chromosome number obtained, 2 n = 60, indicates that the species is a hexaploid. Recent studies have shown that the basic number in the genus is invariably x = 10, and that polyploidy has played a considerable role in the intrageneric evolution (NORDENSTAM 1967 and 1969). The allied species *O. patula* SCHLTR was found to be diploid (2n = 20, counted on NORDENSTAM 2976 from Cape Province, Vanrhynsdorp Division, 15 miles E. of Vanrhynsdorp on road to Van Rhyns Pass, 1963). Another obvious ally, viz. *O. protecta* DTR., is also a diploid (NORDENSTAM 1967 and 1969). The latter is a widespread karroo species, lacking runners, but otherwise very similar to *O. patula*.

O. rechingeri is only known from the vicinity of Nieuwoudtville in Calvinia Division. SCHLECHTER's collection was distributed under the MS. name O. "gracilis", which if published would have become a later homonym of O. gracilis HIERN. SCHLECHTER's locality "Papelfontein" is according to JESSOP (1964, p. 140) probably the same as Papekuilsfontein, a place situated only about 13 miles south of my locality.

Literature cited

JESSOP, J. P. (1964): Itinerary of Rudolf Schlechter's collecting trips in Southern Africa. – Journ. S. Afr. Bot. 30: 129–146.

NORDENSTAM, B. (1961): Die Gattungszugehörigkeit von Euryops sparsiflorus S. Moore. – Mitt. Bot. München 4: 125–126.

- (1967): Chromosome numbers in Othonna (Compositae). Bot. Notiser 120: 297-304.
- (1969): Chromosome studies on South African vascular plants. Ibid. 122: 398-408.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Annalen des Naturhistorischen Museums in Wien

Jahr/Year: 1971

Band/Volume: 75

Autor(en)/Author(s): Nordenstam Bertil

Artikel/Article: Othonna rechingeri B. Nord. Spec.nova, a hexaploid succulent from South 139-142