

A Short Note on the Occurrence of the Upper Triassic Oyster *Umbrostrea? montiscaprilis* (KLIPSTEIN, 1843) (Mollusca: Bivalvia) in the Northern Alpine Raibl Beds of the Schafberg, Salzburg, Austria

ISTVÁN SZENTE¹, HARALD LOBITZER² & FELIX SCHLAGINTWEIT³

3 Text-Figures, 1 Plate

Österreichische Karte 1:50.000
Blatt 65 Mondsee

Northern Calcareous Alps
Northalpine Raibl Beds
Salzkammergut
Schafberg
Bivalvia

Contents

Zusammenfassung	27
Abstract	27
Introduction	27
Geological Setting and Localities	28
Bivalves	28
Acknowledgements	30
References	31
Plate	32

Eine kurze Mitteilung über das Vorkommen der obertriassischen Auster *Umbrostrea? montiscaprilis* (KLIPSTEIN, 1843) (Mollusca: Bivalvia) in den Nordalpinen Raibler Schichten des Schafbergs, Salzburg, Österreich

Zusammenfassung

Aus den Nordalpinen Raibler Schichten des Schafberg-Nordfußes wird eine für das Karnium charakteristische Bivalven-Faunula, die drei Arten umfasst, beschrieben. Trotz Rekristallisation weisen die Schalenstrukturen einiger Exemplare von "*Ostrea*" *montiscaprilis* KLIPSTEIN, 1843 darauf hin, dass diese Art der Gattung *Umbrostrea* HAUTMANN, 2001 angehört.

Abstract

Northern Alpine Raibl Beds exposed at the northern foot of the Schafberg yielded a bivalve faunula consisting of three species characteristic of the Carnian stage. Although the shells are generally recrystallized, preserved structures of some valves of "*Ostrea*" *montiscaprilis* KLIPSTEIN, 1843 indicate that this species belongs to the genus *Umbrostrea* HAUTMANN, 2001.

Introduction

Oysters are a distinct and successful group of marine bivalves. Due to the predominantly calcitic composition of their shells, they have a good fossil record especially from the early Jurassic onwards. By contrast, key characters of classification such as inner features as well as mineralogy and structure of the valves are much less known in Triassic oysters, as a consequence of the frequent loss or recrystallization of their inner, presumably aragonitic shell layers (HAUTMANN, 2001a, 2006a, b). In fact, generic

assignment of several Triassic species is still debated. The last decade saw the publication of a remarkable series of papers dealing with the origin and early evolution of oysters (CHECA & JIMÉNEZ-JIMÉNEZ, 2003; CHECA et al., 2006; HAUTMANN, 2001a, b, 2004, 2006a, b; MALCHUS, 2008; MÁRQUEZ-ALIAGA et al., 2005). In the light of this renewed interest, the occurrence of "*Ostrea*" *montiscaprilis* – originally described by KLIPSTEIN (1843) from the Carnian of the Monte Caprile (Goat Hill, if translated) of the Southern Alps – in the Northern Alpine Raibl Group of the Schafberg (Sheep Mountain, if translated) seems worth describing.

1 ISTVÁN SZENTE: Eötvös Museum of Natural History, H 1117 Budapest, Pázmány P. s. 1/c, Hungary. szente@ludens.elte.hu

2 HARALD LOBITZER: Lindaustraße 3, A 4820 Bad Ischl, Austria. harald.lobitzer@aon.at

3 FELIX SCHLAGINTWEIT: Lerchenauerstraße 167, D 80935 München, Germany. EF.Schlagintweit@t-online.de

Geological Setting and Localities

Since the paper by MOJSISOVICS (1866) fossiliferous exposures of the Northern Alpine Raibl Formation are known from the surroundings of the Eisenaueralm on the northern slope of Mt. Schafberg, south of the eastern end of the lake Mondsee. The outcrops form a largely WNW-ESE striking, approx. 2 km long, wedge-shaped patch on the northern slope of the hill Weinkogel. A good and easily accessible succession of Raibl Beds usually strikingly poor in fossils is exposed along a creek (Sch 1 in Text-Fig. 1, see also SIBLÍK, 2008). It comprises an alternation of limestones with grey marl and marly sandstones beds. At some places, however, bivalves occur in large quantities in limestone beds. Two localities were studied (Text-Fig. 1). Already MOJSISOVICS (1866) mentions fossil findings from there: in the grey sandstones of the Lunz Formation “*Equisetites*” fragments and from the overlying “Cardita Beds” *Avicula aspera* PICHLER, “*Cidaris*” spines, *Pecten*, *Plicatula* and a bed with *Ostrea Montis Caprillis*.

Locality Sch 1 was already reported by SIBLÍK (2008). At the point where the marked tourist path leading from Schafing to the Eisenaueralm crosses the creek, dark grey limestone slabs with abundant remains of *Umbroscaprilis montiscaprilis* can be found in the drift of the creek. Although obviously transported, these slabs were found to provide the only opportunity to collect *U. montiscaprilis* by hand tools.

Two types of fossiliferous Raibl limestones could be distinguished macroscopically, which differ from each other in microfacies as well. Microfacies of the greenish-gray limestone containing abundant shells of *U. montiscaprilis* is characterized by shell fragments of the fore-mentioned species (Text-Fig. 2a). The bluish-grey limestone slabs contain abundant black onkoids of up to 15 mm. Identified fossils include solenoporacean(?) algae, small agglutinated foraminifers, gastropods and echinoderms (Text-Fig. 2b–d). A similar onkolithic bindstone microfacies was reported by BELOCKY et al. (1999) from the Raibl Beds of the Gaisberg/Kirchberg in Tirol.

Slightly upstream, above the small waterfall exposing unfossiliferous marl beds the bed of the creek is formed by limestone bedding planes displaying abundant bivalves among which *Schafhaeutlia? sp. cf. mellingi* (HAUER, 1857) and *U. montiscaprilis* could be identified (Text-Fig. 3a, b).

Some tens of meters upstream, megalodontid? bivalves can be seen in large quantities in a limestone bank forming the right flank of the creek (Text-Fig. 3c, d). The specimens are preserved as internal moulds with conjoined, closed valves and can not be extracted from the compact rock with hand tools.

Raibl Beds are also exposed higher on the right flank of the creek valley, along the path leading from Kreuzstein to the Eisenaueralm (Locality Sch 2). Two bivalve specimens, representing *S. mellingi* and *Rossiodus cf. columbella* (HOERNES, 1855) were found there by Dr. Miloš Siblík, Prague.

Bivalves

Umbroscaprilis montiscaprilis (KLIPSTEIN, 1843)

(Pl. 1, Figs. 1–9, 11?)


Material: about a dozen specimens, presumably all left valves, embedded in compact limestone.

Description: Inaequilateral, backward-curved, higher than long shells ornamented with up to 15 squamose, antimarginal ribs/plicae whose number increases with intercalation of new ones at the postero-ventral region. Attachment area is subordinate if compared with the height of the valve. Umbonal cavity is well defined. Internal features can not be studied in the available material.

The shell structure has been completely obscured by recrystallization in most cases (e.g. Pl. 1, Figs. 5, 6). Some sections, however, display two shell layers of different structure (Pl. 1, Figs. 7, 8). The outer one seems to be of prismatic nature while the inner one is formed by



Text-Fig. 1.
Map showing the bivalve localities studied.



Text-Fig. 2.

Characteristic microfacies of Raibl limestones.

a: fragment of *U. montiscaprilis*, ostracods and echinoderms; b: gastropod with coated and recrystallized shell; c: dark-coloured onkoids; d: partly recrystallized solenoporacean alga?

The bar represents 3 mm.

sparry calcite. In another section layering of the middle part of the shell was found to be still preserved (Pl. 1, Fig. 9). No structural chambers ("Hohlräume") were observed. Some sections show elongated bodies composed of sparry calcite, associated with finely foliated calcite structures (Pl. 1, Fig. 11). Their relationship to *U. montiscaprilis* is, however, not justified.

Remarks: "*Ostrea*" *montiscaprilis*, usually assigned to the genera/subgenera *Lopha* or *Alectryonia* was frequently recorded from various Carnian formations of the Northern Calcareous Alps (see TOLLMANN, 1976) but it was only rarely described and figured (e.g. WÖHRMANN, 1889, p. 200, Pl. 6, Figs. 1–3). The Schafberg specimens agree well in shape with the type (KLIPSTEIN, 1843), as well as with those more recently described and figured in the literature, e.g. by JELEN (1989) and LIEBERMAN (1979) (see also DIENER, 1923 for older references). The species was also recorded from North America (e.g. STANLEY, 1979), however, the specimens referred to have never been described or figured thus it is doubtful whether they are conspecific (McROBERTS, 1997).

U. montiscaprilis differs from *Actinostreon haidingerianum* (EMM-RICH, 1853), a common species in the Rhaetian of the Northern and Southern Alps as well as of the NW Carpathians, by having more ribs/plicae (see e.g. ZAPFE, 1967, p. 438, Pl. 3, Figs. 7a, b; GOETEL, 1917, p. 169, Pl. 9, Figs. 4a, b.) and – probably – by its aragonitic inner shell layers (see below).

Uncertainty concerning the generic assignment of "*O. montiscaprilis*" roots in the lack of appropriate knowledge of its shell mineralogy and structure. Differences between Triassic lophate oysters and *Lopha* RÖDING, 1789 were already recognized by MALCHUS (1990) who erected the new genus *Palaeolopha* based on *Ostrea haidingeriana* as type species, and including – although doubtfully – *Palaeolopha montiscaprilis* as well. According to HAUTMANN (2001a), however, the shell of "*O. haidingeriana*" is entirely calcitic and similar in microstructure to that of *Actinostreon* BAYLE, 1878 as documented by SIEWERT (1972). Thus, *Palaeolopha* should be considered as a junior synonym of *Actinostreon*. On the other hand, evidence presented by HAUTMANN (2001a, b) and MÁRQUEZ-ALIAGA et al. (2005) suggest that shells of



a



b



c



d

Text-Fig. 3.

Field occurrence of Northern Alpine Raibl Beds.

a: unfossiliferous marl and limestone beds; b: bedding plane with bivalves (*Schafhaeutlia?* sp. cf. *mellingi*, *U.?* *montiscaprilis*); c, d: megalodontid? bivalves. The hammer is 33 cm long, the hand lens is 32 mm wide.

Middle and early Late Triassic oysters contained aragonite layers as well. On the basis of shell mineralogy, these forms should be assigned to *Umbrostrea* HAUTMANN, 2001, whose valves consist of an outer layer of calcitic prisms, a thin middle layer of foliated calcite and a thick, originally aragonitic inner layer. Although shell preservation of the Schafberg specimens is far from suitable, the data available make their assignment to *Umbrostrea* the most plausible. The almost exclusive appearance of recrystallized, presumably originally aragonitic shell parts may be due to the peeling off of the thin outer calcitic layers, in a way found by SANDERS et al. (2007) in diceratid rudists. Spalling of the outer, prismatic layer of *U. iranica* HAUTMANN, 2001 was also figured by HAUTMANN (2001a).

***Schafhaeutlia mellingi* (HAUER, 1857)**

Pl. 1, Figs. 10, 12–14

An internal mould of a left valve and another one formed by conjoined, closed valves were found. The umbo placed near the mid-length of the valves as well as the strong conical cardinal tooth of the left valve preserved as external mould are characteristic features of *S. mellingi*, a species

well represented in the Carnian of the Northern Calcareous Alps (see e.g. WÖHRMANN, 1889; TOULA, 1910). In the older literature it is referred to as *Gonodon*, *Gonodus* or *Corbis mellingi*.

***Rossiodus* cf. *columbella* (M. HOERNES, 1855)**

Pl. 1, Fig. 15

A single internal mould of a left valve may represent *R. columbella*, a characteristic Upper Carnian – Norian megalodontid species as described and figured by VÉGH-NEUBRANDT (1982).

Acknowledgments

Dr. Miloš Siblík (Prague) provided us with bivalve specimens he collected. His help and kindness is gratefully acknowledged here. The field-work of I. Szenté and H. Lobitzer was supported on the basis of the bilateral agreement between the Geologische Bundesanstalt (Vienna) and the Geological Institute of Hungary (Budapest). This is a contribution of the OTKA Project K 81298.

References

- BELOCKY, R., SLAPANSKY, P., EBLI, O., OGORELEC, B. & LOBITZER, H. (1999): Die Uran-Anomalie in der Trias-Deckscholle des Gaisberg/Kirchberg in Tirol (Österreich) – Geophysikalische, geochemische und mikrofazielle Untersuchungen. – Abh. Geol. B.-A., **56/2**, 13–33.
- CHECA, A.G., JIMÉNEZ-JIMÉNEZ, A.P. (2003): Rib fabrication in Ostreoidea and Plicatuloidea (Bivalvia, Pteriomorphia) and its evolutionary significance. – Zoomorphology, **122**, 145–159.
- CHECA, A.G., JIMÉNEZ-JIMÉNEZ, A.P., MÁRQUEZ-ALIAGA, A. & HAGEDORN, H. (2006): Further comments on the origin of oysters. – Palaeogeography, Palaeoclimatology, Palaeoecology, **240**, 672–674.
- DIENER, C. (1923): Fossilium Catalogus, Pars 19, Lamellibranchiata triadica. – 257 pp, Junk, Berlin.
- GOETEL, W. (1917): Die Rhätische Stufe und der unterste Lias der subtriatrischen Zone in der Tatra. – Bull. Acad. Sci. Cracovie, Cl. Sci. Math. Nat. Série A, Sci. Math. 1–222, 12 Pls.
- HAUTMANN, M. (2001A): Taxonomy and phylogeny of cementing Triassic Bivalves (Families Prospondylidae, Plicatulidae, Dimyidae and Ostreidae). – Palaeontology, **44**, 339–373.
- HAUTMANN, M. (2001B): Die Muschelfauna der Nayband-Formation (Obertrias, Nor-Rhät) des östlichen Zentraliran. – Beringeria, **29**, 3–181, Würzburg.
- HAUTMANN, M. (2004): Early Mesozoic evolution of alivincular bivalve ligaments and its implications for the timing of the “Mesozoic marine revolution”. – Lethaia, **37**, 165–172.
- HAUTMANN, M. (2006A): Shell mineralogical trends in epifaunal Mesozoic bivalves and their relationship to seawater chemistry and atmospheric carbon dioxide concentration. – Facies, **52**, 417–433.
- HAUTMANN, M. (2006B): Shell morphology and phylogenetic origin of oysters. – Palaeogeography, Palaeoclimatology, Palaeoecology, **240**, 668–671.
- JELEN, B. (1989): The Karnian bivalves from Lesno brdo, Slovenia, NW Yugoslavia and their paleobiological significance. – Geologija, **31/32**, 11–127.
- KLIPSTEIN, A. v. (1843): Beiträge zur geologischen Kenntniss der östlichen Alpen. – 312 pp., 20 Taf., Giessen.
- LIEBERMAN, H.M. (1979): Die Bivalven- und Ostracodenfauna von Raibl und ihr stratigraphischer Wert. – Verh. Geol. B.-A., Wien **1979/2**, 85–131, 5 Pls.
- MALCHUS, N. (1990): Revision der Kreide-Austern (Bivalvia: Pteriomorphia) Ägyptens (Biostratigraphie, Systematik). – Berliner Geowiss. Abh. Reihe A, **125**, 231 p., Berlin.
- MALCHUS, N. (2008): Problems concerning early oyster evolution: a reply to Márquez-Aliaga and Hautmann. – Palaeogeography, Palaeoclimatology, Palaeoecology, **258**, 130–134.
- MÁRQUEZ-ALIAGA, A., JIMÉNEZ-JIMÉNEZ, A.P., CHECA, A.G. & HAGEDORN, H. (2005): Early oysters and their supposed Permian ancestors. – Palaeogeography, Palaeoclimatology, Palaeoecology, **229**, 127–136.
- MCRROBERTS, C.A. (1997): Late Triassic (Norian–Rhaetian) bivalves from the Antimonio Formation, Northwestern Sonora, Mexico. – Revista Mexicana de Ciencias Geológicas, **14**(2), 167–177.
- MOJSISOVICS, E. v. (1866): Ueber die Gliederung der Trias zwischen dem Hallstätter und Wolfgang-See. – Jb. Geol. R.-A., **16**, Verh., 160–164, Wien.
- SANDERS, D., LUKESCH, M., RASSER, M. & SKELTON, P. (2007): Shell beds of diceratid rudists ahead a low-energy gravelly beach (Tithonian, Northern Calcareous Alps, Austria): Paleocology and Taphonomy. – Austrian J. Earth Sci., **100**, 186–199, Wien.
- SIBLIK, M. (2008): Bericht 2008 über Untersuchungen von Brachiopoden in den Raibler Schichten auf Blatt 65 Mondsee. – Jb. Geol. B.-A., **148/2**, 276–277, Wien.
- SIEWERT, W. (1972): Schalenbau und Stammesgeschichte von Austern. – Stuttgarter Beitr. Naturk. B (Geol. Paläont), **1**, 1–57.
- STANLEY, G.D. JR. (1979): Paleocology, structure, and distribution of Triassic coral buildups in Western North America. – The University of Kansas Paleontological Contributions, **65**, 1–58.
- TOLLMANN, A. (1976): Analyse des klassischen nordalpinen Mesozoikums. Stratigraphie, Fauna und Fazies der Nördlichen Kalkalpen. – 580 pp, Deuticke, Wien.
- TOULA, F. (1910): Schichten mit *Gervillella* (“*Perna*”) *Bouéi* v. HAUER, am Gaumannmüllerkogel an der Weissenbacher Strasse. – Jb. Geol. R.-A., **59/2**, 383–406, Taf. 12, Wien.
- VÉGH-NEUBRANDT, E. (1982): Triassische Megalodontaceae. Entwicklung, Stratigraphie und Paläontologie. – 526 p. Akadémiai Kiadó, Budapest.
- WÖHRMANN, S. v. (1889): Die Fauna der sogenannten *Cardita*- und Raibler-Schichten in den Nordtiroler und bayerischen Alpen. – Jb. Geol. R.-A., **39/1**, 181–258, Taf. 5–10, Wien.
- ZAPFE, H. (1967): Beiträge zur Paläontologie der nordalpinen Riffe. Die Fauna der Zlambach-Mergel der Fischerwiese bei Aussee, Steiermark. – Ann. Naturhist. Mus. Wien, **71**, 413–480.

Received: 10. September 2010, Accepted: 13. October 2010

Plate 1

The specimens in Figs. 2–4, 10 and 12–15 are coated with ammonium-chloride.

The scale bar is 3 mm in Figs. 5–7 and 11, and 1 mm in Figs. 8, 9.

Figs. 1–9, 11?: *Umbroscapria? montiscaprilis* (KLIPSTEIN, 1843).

Fig. 1: characteristic occurrence and preservation of valves, 0.6x.

Figs. 2–4: left valves.

Figs. 5, 6: longitudinal section of specimen in Fig. 2, umbo is to the left, acetate peels.

Figs. 7, 8: cross section of ribs displaying traces of an outer calcitic prismatic? and an inner, originally aragonitic? shell layer, acetate peel.

Fig. 9: preserved structure of the inner, recrystallized shell layer, acetate peel.

Fig. 11: fine foliated calcite structures associated to recrystallized shell fragments, questionably interpreted as peeled off shell layers of *U.? montiscaprilis*, acetate peel.

Figs. 10, 12–14: *Schafhaeutlia mellingi* (HAUER, 1857).


Fig. 10: internal mould of a left valve bearing fine commarginal and radial ornamentation.

Fig. 12: lateral view.

Fig. 13: frontal view.

Fig. 14: dorsal view.

Fig. 15: *Rossiodus cf. columbella* (M. HOERNES, 1855).



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Abhandlungen der Geologischen Bundesanstalt in Wien](#)

Jahr/Year: 2010

Band/Volume: [65](#)

Autor(en)/Author(s): Szent-Istvanyi Szentistvanyi Gyula, Lobitzer Harald,
Schlagintweit Felix

Artikel/Article: [A Short Note on the Occurrence of the Upper Triassic Oyster
Umbrostrea? montiscaprilis \(KLIPSTEIN, 1843\) \(Mollusca: Bivalvia\) in the Northern
Alpine Raibl Beds of the Schafberg, Salzburg, Austria 27-33](#)