

Zollner Formation

CLAUDIA SPALLETTA, HANS-PETER SCHÖNLAUB, MARIA CRISTINA PERRI, CORRADO VENTURINI & MONICA PONDRELLI

Österreichische Karte 1:50.000

Blatt BMN 197 Kötschach

Blatt BMN 198 Weißbriach

Blatt BMN 199 Hermagor

Blatt UTM 3109 Oberdrauburg

Blatt UTM 3110 Kötschach-Mauthen

Blatt UTM 3116 Sonnenalpe Naßfeld

Blatt UTM 3117 Nötsch im Gailtal

Carta Topografica d'Italia 1:50.000

Foglio 018 Passo di Monte Croce Carnico

Foglio 031 Ampezzo

Foglio 032 Tolmezzo

Foglio 033 Tarvisio

Definition

Well bedded light gray, greenish to black chert cm to dm thick, interlayered with rare limestone lenses, and black siliceous shale.

Description


The Zollner Formation consists mainly of radiolarian chert with interbedded rare limestone and black siliceous shale. The limestones are represented by centimetric/decimetric lenses and rare layers of wackestone/mudstone; present along the whole unit. The black shales are more abundant in the lower part of the formation.

Fossil content


Conodonts, graptolites (in the lowermost part), radiolarians.

Depositional environment

Pelagic.



Areas of outcrop of the Zollner Formation with indication of the stratotype of the lower and upper boundary (asterisks), A: Wasserfall Section; B: Rio Chianaletta Section. Reference section (square), Oberbuchach 3 section.



The Zollner Formation type section. a) sketch of the Wasserfall Section (after SCHÖNLAUB, 1985); b) log of the Rio Chianaletta Section (modified after PERRI & SPALLETTA, 1998). Legend: 1. biomicrite; 2. radiolarian-enriched mudstone; 3. radiolarian chert with interbedded limestone levels and lenses; 4. radiolarian chert; 5. breccia of the Hochwipfel Formation; 6. sandstone and pelite of the Hochwipfel Formation; c) view of the section in the field (photo M.C. PERRI).

Stratotype

Wasserfall Section near Lake Zollner (SCHÖNLAUB, 1985), at coordinates N 46°36'18", E 13°04'11" for the lower boundary.

Rio Chianaletta Section (PERRI & SPALLETTA, 1998), at coordinates N 46°36'13.7", E 12°50'02.8" for the upper boundary.

Reference sections

Oberbuchach 3 section (JAEGER & SCHÖNLAUB, 1980) below Gundersheimer Alm at coordinates N 46°37'29.1", E 13°05'52.6", where the transition with the Bischofalm Formation is well exposed.

Type area

Carnic Alps.

Main outcrop areas

The Zollner Formation crops out mainly in the Cima Ombladet-Rio Chianaletta area to the west, in a wide area of the Central Carnic Alps, and in the area between Mt. Poludnig and the Gail River to the east.

Thickness

The maximum estimated thickness is more than 100 m. Continuous sections exposing the whole unit are not known.

Boundaries

Underlying units – Bischofalm Formation (conformable, gradual contact).

Overlying units – Hochwipfel Formation (conformable, gradual contact) in most cases affected by tectonic strain.

Lateral units – Findenig Formation, Valentin Formation and Pal Grande Formation.

Derivation of name

After Lake Zollner.



Views of the Zollner Formation in the field. a) radiolarian chert with limestone lenses at Rio Chianaletta (photo C. SPALLETTA); b) radiolarian chert in the Pramasio Pass area (photo C. VENTURINI).

Synonymy

Radiolariti (liditi): SPALLETTA et al. (1982).
 Zollner Formation: SCHÖNLAUB (1985); SCHÖNLAUB et al. (2004).
 Radiolarites and shales: SPALLETTA & VENTURINI (1990).
 Rio Chianaletta Radiolarites: BRIME et al. (2008).
 Radiolariti del Rio Chianaletta: SPALLETTA (2009).
 Zollner-Formation/Zollner Formation: SUTTNER & KIDO (2014).

Chronostratigraphic age

Devonian – Carboniferous: Lochkovian to lower Viséan.

Biostratigraphy

Conodonts. – The youngest level of the formation has been assigned to the *texanus-homopunctatus* Zone (PERRI & SPALLETTA, 1998).

Graptolites. – *Monograptus hercynicus* Zone for the oldest level of the formation (HERZOG, 1983, 1988).

Complementary references -

Remarks -

References

- BRIME, C., PERRI, M.C., PONDRELLI, M., SPALLETTA, C. & VENTURINI, C. (2008): Polyphase metamorphism in the eastern Carnic Alps (N Italy-S Austria): Clay minerals and conodont Colour Alteration Index evidence. – *International Journal of Earth Sciences*, **97**/6, 1213–1229, Berlin-Heidelberg.
- HERZOG, U. (1983): Zur Gewinnung von Conodonten aus Lyditen und zur zeitlichen Basis des Hochwipfel-Karbon im Gebiet des Poludnig (Östliche Karnische Alpen). – *Carinthia II*, **173**/93, 363–369, Klagenfurt.
- HERZOG, U. (1988): Das Paläozoikum zwischen Poludnig und Oisternig in den Östlichen Karnischen Alpen. – *Carinthia II*, **47**, 1–123, Klagenfurt.
- JAEGER, H. & SCHÖNLAUB, H.P. (1980): Silur und Devon nördlich der Gundersheimer Alm in den Karnischen Alpen (Österreich). – *Carinthia II*, **170**/90, 403–444, Klagenfurt.
- PERRI, M.C. & SPALLETTA, C. (1998): Conodont distribution at the Tournaisian/Viséan boundary in the Carnic Alps (Southern Alps, Italy). – In: SZANIAWSKI, H. (ed.): *Proceedings of the Sixth European Conodont Symposium (ECOS VI)*. – *Palaeontologia Polonica*, **58**, 225–245, Warszawa.
- SCHÖNLAUB, H.P. (1985): Das Paläozoikum der Karnischen Alpen. – In: SCHÖNLAUB, H.P. (ed.): *Arbeitstagung der Geologischen Bundesanstalt 1985 Kötschach-Mauthen, Gailtal* – Geologische Bundesanstalt, 34–52, Wien.
- SCHÖNLAUB, H.P., HISTON, K. & POHLER, S. (2004): The Palaeozoic of the Carnic Alps. – In: SCHÖNLAUB, H.P. (ed.): *Field Trip Carnic Alps Guidebook*. June 23–24, 2004, Carinthia, Austria. – Geologische Bundesanstalt, 2–32, Wien.

SPALLETTA, C. (2009): Radiolariti del Rio Chianaletta. – In: VENTURINI, C. (ed.): Note Illustrative del Foglio 031 Ampezzo. – Carta Geologica d'Italia alla scala 1:50000, Istituto Superiore per la Protezione e la Ricerca Ambientale (ex-Agenzia per la Protezione dell'Ambiente e per i Servizi Tecnici, Servizio Geologico d'Italia), 46-47, Stampa A.T.I. – S.EL.CA. srl. – L.A.C. srl. – System Cart srl., Firenze.

SPALLETTA, C. & VENTURINI, C. (1990): Stratigraphic correlation form of the Palaeozoic sequence in the Carnic Alps. – Rendiconti della Società Geologica Italiana, **12**, 417–421, Roma.

SPALLETTA, C., VAI, G.B. & VENTURINI, C. (1982): La Catena Paleocarnica. – In: CASTELLARIN, A. & VAI, G.B. (eds.): Guida alla geologia del Sudalpino centro-orientale. – Guide Geologiche Regionali, Società Geologica Italiana, 281–292, Bologna.

SUTTNER, T.J. & KIDO, E. (2014): Zollner-Formation/Zollner Formation. – In: PILLER, W.E. (ed.): The lithostratigraphic units of the Austrian Stratigraphic Chart 2004 (sedimentary successions), Vol.I - The Paleozoic Era(them). – Abhandlungen der Geologischen Bundesanstalt, **66**, 73, Wien.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Abhandlungen der Geologischen Bundesanstalt in Wien](#)

Jahr/Year: 2015

Band/Volume: [69](#)

Autor(en)/Author(s): Spalletta Claudia, Schönlaub Hans-Peter, Perri Maria Cristina, Venturini Corrado, Pondrelli Monica

Artikel/Article: [Zollner Formation 141-144](#)