

**FIRST RECORD OF *AGATHIS RUBENS* TOBIAS FROM TURKEY
(HYMENOPTERA: BRACONIDAE: AGATHIDINAE)**

Özlem Çetin ERDOĞAN & Ahmet BEYARSLAN

Trakya University, Faculty of Arts and Science, Department of Biology. 22030
Edirne, Turkey, e-mail: cetinozlem@hotmail.com

**Izvešček - PRVA NAJDBA VRSTE *AGATHIS RUBENS* TOBIAS V TURČIJI
(HYMENOPTERA: BRACONIDAE: AGATHIDINAE)**

Vrsta najezdnikov *Agathis rubens* Tobias, 1963, je bila opisana iz Kazahstana in do zdaj ni bila znana od drugod. Navajamo prvo najdbo te vrste v Turčiji.

Forty-six species of *Agathis* are known from the West Palaearctic region. Of these, 29 have been recorded from Europe (Simbolotti and van Achterberg, 1999). Generally, agathidines are more diverse in humid tropical regions but the genus *Agathis* is more speciose in temperate regions. According to Nixon (1986) *Agathis-Microdus*-group genera are restricted mainly to the Northern Hemisphere. Hitherto, 18 species of *Agathis* have been reported from Turkey (Nixon, 1986; Zettel and Beyarslan, 1992; Cetin and Beyarslan, 2001; Simbolotti and van Achterberg, 1999; Beyarslan et.al. 2002a, 2002b).

We present here the first record of *Agathis rubens* Tobias, 1963 from Turkey. *Agathis rubens* was described on the specimens from Kazakhstan (Transcaucasia) (Tobias, 1963) and has never been recorded in any other region up to now. We found it in Western Turkey (Afyon Prov. Sincanli. Güney, 1250 m, 39° 08' 16N, 30° 58' 37E, 30. 6. 1998, 1♀). The material is deposited in the Zoological Museum of the Department of Biology, Trakya University (TU), Turkey. The species differs from other species of the genus by very weak or undeveloped notauli, combined with almost uniform reddish-yellow body and dark wings. The female is 4.7 mm long.

References

- Beyarslan, A., F. Inane, Ö. Cetin & M. Aydogdu**, 2002a: Braconiden von den türkischen Inseln Imbros und Tenedos (Hymenoptera, Braconidae, Agathidinae, Braconinae, Cheloninae, Microgastrinae). *Entomofauna*, 23(15): 173-188.
- Beyarslan, A., F. Inane, Ö. Cetin, M. Aydogdu**, 2002b: Braconidae Species of Turkish Aegean Region, International Symposium: Parasitic Hymenoptera; Taxonomy and Biological Control, 14-17 May, 2001, Kőszeg-Hungary, 285-290.

- Cetin, Ö. & A. Beyarslan**, 2001: The Agathidinae (Hymenoptera: Braconidae) Fauna of the Marmara Region. *Doga Turkish Journal of Zoology*, 25(3): 257-268.
- Nixon, G.E.J.**, 1986: A revision of the European Agathidinae (Hymenoptera: Braconidae). *Bulletin of the British Museum Natural History (Entomology)* Ser. 52 (3): 183- 242.
- Simbolotti, G. & C. van Achterberg**, 1999: Revision of the West Palearctic species of the genus *Agathis* Latreille (Hymenoptera: Braconidae). *Zoologische Verhandelingen, Leiden* 325: 1-167.
- Tobias, V.I.**, 1963: Parazititseskie perepontosatokrylye roda *Agathis* Latr. (Hymenoptera: Braconidae) Kazachstana i srednej Azii. *Entomologicheskoe Obozrenie*, 42: 864-883.
- Zettel, H. & A. Beyarslan**, 1992: Über Agathidinae aus der Türkei (Hymenoptera: Braconidae). *Entomofauna*, 13(5): 121-132.

Received / Prejeto: 7. 6. 2004