

LJUBLJANA, DECEMBER 2005

Vol. 13, št. 2: 77–80

A NEW SPECIES OF THE GENUS *MYCTERODUS* SPINOLA (HEMIPTERA: FULGOROMORPHA: ISSIDAE) FROM CROATIA

Vladimir M. GNEZDILOV¹ & Michael R. WILSON²

¹Zoological Institute, Russian Academy of Sciences, Universitetskaya nab.1,
199034, St. Petersburg, Russia. E-mail: hemipt@zin.ru

²Department of Biodiversity & Systematic Biology, National Museum of Wales,
Cathays Park, Cardiff CF10 3NP,
UK. E-mail: mike.wilson@nmgw.ac.uk

Abstract A new species of the genus *Mycterodus* (subgenus *Semiroodus*) – *Mycterodus croaticus* sp. n. from Croatia (Dubrovnik, Ombla, Loznica) is described. It is closely related to *Mycterodus pallens* Stål, 1861, but differs in more acutely angulate anterior margin of coryphe and straight ventral margin of dorso-lateral phallobase lobe under semicircular toothed process.

KEY WORDS: Hemiptera, Issidae, *Mycterodus*, *Semiroodus*, new species

Izvleček – NOVA VRSTA RODU *MYCTERODUS* SPINOLA (HEMIPTERA: FULGOROMORPHA: ISSIDAE) IZ HRVAŠKE

Opisana je nova vrsta rodu *Mycterodus* (podrod *Semiroodus*) – *Mycterodus croaticus* sp. n. iz Hrvaške (Dubrovnik, Ombla, Loznica). Je v bližnjem sorodstvu z vrsto *Mycterodus pallens* Stål, 1861, vendar se od nje razlikuje po bolj ostrih vogalih prednjega roba korife in ravnem trebušnem robu dorzolateralnega režnja falobaze pod polkrožnim zebastim izrastkom.

KLJUČNE BESEDE: Hemiptera, Issidae, *Mycterodus*, *Semiroodus*, nova vrsta

Introduction

According to recently published review of the subgenus *Semiroodus* of the genus *Mycterodus* (Gnezdilov & Drosopoulos, 2005), *Mycterodus pallens* Stål, 1861 recorded by Melichar (1906) from Croatia is in fact distributed only in Greece. Examined material from Croatia belongs to a new species closely related to *M.*

pallens, but differs in the structure of male genitalia illustrated first by Dlabola (1985, figs 18-22) under the name *M. pallens*.

Material and methods

The material examined is deposited in the following collections: BMNH – the Natural History Museum (London, UK), NMWC – National Museum of Wales (Cardiff, UK), MMBC – Moravian Museum, Collection of L. Melichar (Brno, Czech Republic), SMNH – Swedish Museum of Natural History (Stockholm, Sweden), ZIN – Zoological Institute, Russian Academy of Sciences (St. Petersburg, Russia).

Taxonomy


Mycterodus croaticus sp. n.

Holotype: ♂, Croatia, Dubrovnik, 23. V 1952, W.J. Le Quesne (BMNH).


Paratypes: Croatia: 1 ♀, as holotype (BMNH); 1 ♂, Dubrovnik, "89-51", "Dr. Eger." (ZIN); 1 ♂, Ombla, 27. V. 1934 (BMNH); 1 ♂, 2 ♀, Loznica, 7. VI. 1934 (BMNH and ZIN); 1 ♀, same locality, 24. 27. VI. 1934 (BMNH), all leg. D. Aubertin; 1 ♀, "Dalmatien", P. Novak leg. (NMWC); 1 ♂, 1 ♀, Ragusa (Dubrovnik) (MMBC).

Description: Externally very similar to *M. pallens* Stål, 1861 (syntype examined, SMNH) (Fig. 1), but distinguished by more acutely angulate anterior margin of coryphe (Fig. 2) and the structure of male genitalia (Figs 3-5).

Coloration: General coloration from light yellowish brown to dark brown. Apical cells of fore wings with dark brown spots. Metope with sublateral keels bordered



Figs 1-2: *Mycterodus* spp., head, pronotum, and scutellum, dorsal view. 1 – *Mycterodus pallens* Stål, ♀, syntype; 2 – *Mycterodus croaticus* sp. n., ♀, paratype.



Figs 3-5: *Mycteroodus croaticus* sp. n., male genitalia. 3 – stylus, lateral view; 4 – anal tube, dorsal view; 5 – penis, lateral view.

apically by black stripe and with V-shaped dark brown spot under the keels. Traces of the sensory pits of metope between its upper margin and sublateral keels black. Pronotum with a couple of dark brown dots medially. Each fore wing usually with three dark brown dots: one dot between M and CuA basally and two dots between second branche of R and first branche of M (one dot near Median bifurcation and another dot distally). Hind wings yellowish light brown. Socle setae and teeth of legs dark brown.

Male genitalia: Pygofer with weakly convex hind margin. Anal tube elongate, narrows apically (in dorsal view) (Fig. 4). Anal column short (0.15 times as long as anal tube) and wide. Dorso-lateral lobes of phallobase widening before apex (in lateral view). Ventral margin of each lobe with large semicircular toothed process, the margin under the process straight (in lateral view) (Fig. 5). Apical processes of aedeagus long and narrow apically, each with long and broad apical lobe and excavated margin of phallotrema (in lateral view). Aedeagal hooks gradually narrow

to apex, about 0.5 times as long as aedeagus. Stylus with straight hind margin; caudo-dorsal angle rounded (Fig. 3).

Body length: Males 6.2-7.0 mm. Females 6.4-6.9 mm.

Etymology: The species name is derived from the name of the country – Croatia.

Comparison: The species clearly differs from *M. pallens* in straight ventral margin of dorso-lateral phallobase lobe under semicircular toothed process (Fig. 5).

Acknowledgements

We are pleased to thank The Royal Society (London, UK) for financial support to Vladimir M. Gnezdilov for collaborative study. We are sincerely grateful to Mr. Mick Webb (London, UK), Dr. Pavel Lauterer and Dr. Igor Malenovsky (Brno, Czech Republic), and Dr. Bert Viklund (Stockholm, Sweden) for the opportunity to examine the material.

References

- Dlabola J.**, 1985: Neue mediterrane, eremische und ostafrikanische Issiden-Taxone (Hom., Auchenorrhyncha). *Acta Musei Nationalis Pragae*, **40** B(3-4): 217-243.
- Gnezdilov V.M. & Drosopoulos S.**, 2005: Review of the subgenus *Semiroodus* Dlabola of the genus *Mycteroodus* Spinola (Homoptera, Fulgoroidea, Issidae). *Annales de la Société Entomologique de France*, **40** (3-4), 2004: 235-241.
- Melichar L.**, 1906: Monographie der Issiden (Homoptera). *Abhandlungen der K.K. Zool.-Botan. Gesellschaft in Wien*, **3** (4): 1-327.

Received / Prejeto: 13. 10. 2005

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Acta Entomologica Slovenica](#)

Jahr/Year: 2005

Band/Volume: [13](#)

Autor(en)/Author(s): Gnezdilov Vladimir M., Wilson Michael R.

Artikel/Article: [A new species of the genus Mycterodus Spinola \(Hemiptera: Fulgoromorpha: Issidae\) from Croatia Nova vrsta rodu Mycterodus Spinola \(Hemiptera: Fulgoromorpha: Issidae\) iz Hrvatske 77-80](#)