

Identification of Central European ground beetles of the genus *Bembidion* (Coleoptera: Carabidae) using DNA barcodes: A case study of selected species¹

Michael J. RAUPACH, Karsten HANNIG & Johann-Wolfgang WÄGELE

¹ Dedicated to Prof. Gerd Müller-Motzfeld (†)

Abstract: Identification of Central European ground beetles of the genus *Bembidion* (Coleoptera: Carabidae) using DNA barcodes: A case study of selected species - Species identification based on morphological characters is challenging when closely related species are found at a given locality. In this context, the identification of immature stages is difficult, especially due to lack of distinct morphological differences. Sequence diversity within a fragment of the mitochondrial cytochrome *c* oxidase subunit 1 gene (COI), the so-called DNA barcode for animals, has been shown to represent an effective molecular tool for valid species identification. In this study we examined the performance of DNA barcodes in the discrimination of 21 ground beetle species of the genus *Bembidion* from Central Europe, analysing 101 specimens. Our results revealed low intraspecific variation (mean = 0.19%) and a clear barcoding gap. Although further work is needed to examine patterns of sequence diversity on a broader geographical scale as well as more specimens and additional species, our case study encourages the use and effectiveness of DNA barcodes for a reliable identification of Central European *Bembidion* species.

1 Introduction

The Carabidae count as one of the largest and most diverse insect families, with no less than an estimated 40,000 described species inhabiting all terrestrial habitat types from the sub-arctic to wet tropical regions (ARNETT & THOMAS 2000, ARNDT et al. 2005). Ground beetles, which constitute more than 750 species, are also an important element of the Coleopteran fauna of Central Europe (MÜLLER-MOTZFELD 2004). Carabidae are known to be a species-rich group with a high level of morphological and ecological diversity (e.g. LÖVEI & SUNDERLAND 1996), exhibiting different levels of habitat selectivity that range from generalist to specialist. As a consequence, carabids are often numerically dominant in collections of ground-dwelling arthropods and can be used as indicator organisms for assessments of environmental pollution, habitat classification, conservation, or the characterization of soil nutrients (e.g. LÖVEI & SUNDERLAND 1996, RAINIO & NIEMELÄ 2003, AVGIN & LUFF 2010). However, many species

are difficult to identify morphologically and usually require the help of highly trained taxonomists due to different degrees of morphological variability and polymorphisms within and between species. It is also very difficult or impossible to identify the larvae or females of some species, e.g. various *Amara* species (HURKA 1996).

In this context, DNA barcodes may represent a new effective tool for valid species identification for large-scale biodiversity studies (e.g. VALENTINI et al. 2008, SMITH & FISHER 2009, HEBERT et al. 2010). During recent years DNA barcoding has emerged as a rapid method for species discovery and biodiversity assessment (e.g. HAJIBABAEI et al. 2006, STOECKLE & HEBERT 2008, RADULOVICI et al. 2009, ZHOU et al. 2009). A DNA barcode itself consists of a 648 bp region of the mitochondrial cytochrome *c* oxidase 1 (COI) gene. The concept of DNA barcoding is based on the assumption that each species will most likely have a unique DNA barcode and that intraspecific COI variation is lower than interspecific COI variation, generating the so-called barcoding gap (HEBERT

et al. 2003a, 2003b, 2004). In contrast to DNA taxonomy, which focuses on the classification of both known and undescribed species (TAUTZ et al. 2003, VÖGLER & MONAGHAM 2007), the main goal of DNA barcoding is to identify unknown specimens in terms of a known and existing classification (HEBERT et al. 2003a). The effectiveness of DNA barcoding for the identification of species has been proven in many studies, analysing both vertebrate and invertebrate taxa (e.g. CLARE et al. 2007, COSTA et al. 2007, HUBERT et al. 2008, MACHIDA et al. 2009, RIVERA & CURRIE 2009, ROBINSON et al. 2009, UTHICKE et al. 2010). However, there are only few studies that analyse the usefulness of COI barcodes for the species identification of ground beetles (GREENSTONE et al. 2005, MADDISON 2008, RAUPACH et al. 2010a, 2010b).

In this study we tested the ability of DNA barcodes to discriminate a variety of Central European species of the genus *Bembidion*. The genus *Bembidion* is the largest genus within the Carabidae with more than 1,200 species worldwide (LINDROTH 1985, MADDISON 2008). All species are small (less than 9 mm), move very fast and most species live close to water (LINDROTH 1985). We analysed a subset of 21 species, including a variety of morphologically similar and obviously closely related or sibling species, such as the species pairs e.g. *Bembidion lampros*/*Bembidion properans* (DESENDER & CRAPPE 1983, MÜLLER 1971), *Bembidion millerianum*/*Bembidion ruficorne* (REITTER 1908, MÜLLER-MOTZFELD 1982) or *Bembidion deletum*/*Bembidion incognitum* (MÜLLER-MOTZFELD 2004).


2 Material and Methods

In total, 101 adult ground beetle specimens of the genus *Bembidion* were collected by the authors in various localities in Austria and Germany (Tab. 1). All beetles were identified by one of the authors of this article (KH) using the keys in MÜLLER-MOTZFELD (2004). Numbers of analysed specimens per species ranged between two and ten beetles. Specimens are deposited in the collection of the Zoologisches Forschungsmuseum Alexander Koenig (ZFMK) in Bonn, Germany. Total genomic DNA was extracted from dissected legs of specimens or complete specimens using the QIAmp® Tissue Kit (Qiagen GmbH, Hilden), following the manufacturer's extraction protocol. Polymerase chain reaction (PCR) has been used

for amplifying the COI barcode fragment using the primer pair LCO1480 and HCO2198 (FOLMER et al. 1994). The PCR mix contained 4 µl Q-Solution, 2 µl 10x Qiagen PCR buffer, 2 µl dinucleotide triphosphates (dNTPs, 2 mmol/µl), 0.1 µl of each primer (both 50 pmol/µl), 1 µl of DNA template with of between 2 and 150 ng/µl, 0.2 µl Qiagen Taq polymerase (5 U/µl), and filled up to 20 µl with sterile H₂O. PCR amplification reactions were conducted in Thermal Cycler GeneAmp® PCR System 2700/2720 (Applied Biosystems, Darmstadt) or the Eppendorf Mastercycler® Pro system. The PCR thermal conditions included an initial denaturation at 94 °C (5 min), followed by 38 cycles at 94 °C (denaturation, 45 s), 48 °C (annealing, 45 s), 72 °C (extension, 80 s), and a final extension step at 72 °C (7 min). Negative and positive controls were included with each round of reactions. All PCR products were purified with the QIAquick® PCR Purification Kit (Qiagen GmbH, Hilden). Purified PCR products were cycle-sequenced and sequenced in both directions at a contract sequencing facility (Macrogen, Seoul, Korea) on an ABI3730 XL automatic DNA sequencer, using the same primers as used in PCR. Double-stranded sequences were assembled using the SeqMan™ II programme (DNASTAR, Inc., Konstanz, Germany) or the Geneious® 5.3 programme package (DRUMMOND et al. 2010). We also performed BLAST searches to confirm the identity of all new COI sequences (ALTSCHUL et al. 1990). Aligned sequences were translated to amino acid sequences to check for nuclear mitochondrial pseudogenes (numts) using Geneious® 5.3 (DRUMMOND et al. 2010). Various *Bembidion* COI sequences were already published as part of a comprehensive study analysing the utility of DNA barcodes and nuclear rDNA expansion segments for carabid species identification (see RAUPACH et al. 2010a). All analysed sequences are available in GenBank (see Tab. 1).

Sequences were aligned using MUSCLE version 3.6 (EDGAR 2004) with default settings. The analysis of intra- and interspecific genetic variability of the analysed *Bembidion* species based on Kimura 2-parameter distances (K2P; KIMURA 1980) were calculated using MEGA 5.03 (TAMURA et al. 2011). Intra- as well as interspecific K2P distances were plotted as histogram (Figure 1) using PAST version 1.94b (HAMMER et al. 2001). Base frequencies were obtained using MEGA 5.03 (TAMURA et al. 2011). We also performed a neighbour-joining cluster analysis (SAITOU & NEI 1987) for a graphical repre-

Fig. 1: Histogram of the calculated intra- and interspecific Kimura 2-parameter distances of the COI barcode fragment for the analysed *Bembidion* specimens.



sensation of patterns of nucleotide divergences using MEGA 5.03, based on K2P distances (Figure 2). Four COI sequences of *Omophron limbatum* were used as outgroup (GenBank accession numbers GU347265-GU347268). Finally, bootstrap support values were calculated by re-sampling and analysing 1,000 replicates (FELSENSTEIN 1985).

3 Results

Our analysis revealed low GC-contents (32.6 %) for the analysed sequences, as known in most insects (MIN & HICKEY 2007, CLARE et al. 2008, WEI et al. 2010). Individual mean nucleotide content was A = 0.299, G = 0.166, C = 0.16 and T = 0.375. The analysis of pairwise COI nucleotide divergences for all *Bembidion* species showed higher interspecific than intraspecific divergences (Figure 2). All specimens of the same species grouped together, even when samples were obtained from geographically disparate areas. Consequently, all analysed species were identified in the sampled material and a clear barcoding gap was revealed. Mean intraspecific genetic divergence for the analysed 21 *Bembidion* species was 0.19 %. Maximum intraspecific divergence was observed in *Bembidion properans* (1.92 %), followed by *Bembidion testaceum* (1.27 %) and *Bembidion litorale* (0.95 %). Interspecific distances ranged between 0.0458 and

0.1998 (mean = 14.71 %). Lowest interspecific distances were found between *Bembidion millerianum* and *Bembidion ruficorne* (4.58 – 4.75 %), indicating a close relationship between both species.

It is very important to mention that DNA barcoding focuses on species delineation and identification, and not on phylogenetic inference. Therefore, the presented topology (Figure 2) does not show reliable phylogenetic relationships between the analysed taxa.

4 Discussion

When it began in 2003, the International Barcode of Life project (www.ibol.org) was conceived as a standard system for a fast, valid and accurate identification of animal species (HEBERT et al. 2003a, MILLER 2007). In this context, DNA barcodes help to assign unidentified specimens to known species as well as to identify new species, especially taxa that have a complex or inaccessible morphology (HEBERT et al. 2003a). One obvious advantage of DNA barcoding comes from the rapid and standardized acquisition of molecular data, using conventional and inexpensive protocols for DNA extraction, amplification and sequencing (BORISENKO et al. 2009, IVANOVA et al. 2009). Most importantly, it facilitates the compilation of a precise, representative open-access reference library (BOLD, www.boldsystems.org). To




Fig. 2: Neighbour-joining topology based on Kimura 2-parameter distances as well as images of selected analysed species. Numbers next to internal branches are bootstrap values [%], which are only given if they had a value of 90 or more.

date, (24.07.2011), 1,312,754 barcodes of 107,794 different species have been published. In contrast to other traditional sequence data libraries (e.g. NCBI or EMBL), BOLD is also an interactive interface where deposited sequences can be revised and taxonomically reassigned (RATNASINGHAM & HEBERT 2007).

Despite the fact that DNA barcoding is still subject to criticism, viewed with reservation and/or simply not understood (e.g. DeSALLE et al. 2005, WILL et al. 2005, KLAUSNITZER 2010, EBACH 2011), barcoding has become an established and highly efficient method for understanding and assessing biodiversity (VALENTINI et al. 2008, JINBO et al. 2011). All around the world, numerous projects have been started to compile reference libraries from specimens of various taxa whose identity is already firmly established (e.g. SMITH & FISHER 2009, BUCKLIN et al. 2010, HAUSMANN et al. 2011, LAKRA et al. 2011). Besides this, various national and regional campaigns, for example NorBoL (Norway, www.dnabarcoding.no), JBoL (Japan, www.jboli.org), or Barcoding Fauna Bavaria (www.faunabavarica.de), as well as global taxa-specific campaigns (e.g. relating to Porifera (www.spongebarcoding.org), fishes (www.fishbol.org), birds (www.barcodingbirds.org), ants (www.formicidaebol.org) or butterflies (www.lepbarcoding.org)) have started to support the iBoL initiative. Initial results show that in many cases there are more species – each more narrowly specialized – than scientists had realized before, e.g. within butterflies (HEBERT et al. 2004, STRUTZENBERGER et al. 2011), molluscs (JOHNSON et al. 2008), amphipods (COSTA et al. 2010), and fishes (ZEMLAJ et al. 2009). A preliminary study of 75 Central European ground beetles revealed the existence of cryptic species within *Nebria hellwigii* (RAUPACH et al. 2010a). However, the use of mitochondrial gene fragments for species identification is not without problems: pseudogenes (numts) (e.g. BENSASSON et al. 2000, KIM et al. 2006, KOUTROUMPA et al. 2009), introgressive hybridization (e.g. SOTA et al. 2001, CROUCHER et al. 2004, ZHANG & SOTA 2007), or maternally inherited α -proteobacteria (e.g. *Wolbachia*) (DURON et al. 2008, WERREN et al. 2008) can influence mitochondrial gene variability. In such cases, the analysis of supplementary nuclear markers is necessary (RAUPACH et al. 2010a).

Besides a valid and accurate identification of animal species, DNA barcodes can also support other scientific domains, for example ecology (VALENTINI

et al. 2008), forensics (DAWNAY et al. 2007, NELSON et al. 2007, FERRI et al. 2009, MEIKLEJOHN et al. 2011), pest biology (ENGSTRAND et al. 2010), evolutionary biology (e.g. CLARE et al. 2008, DA SILVA et al. 2011), invasive species biology (ARMSTRONG & BALL 2005, DARLING & BLUM 2007), biogeography (e.g. LINARES et al. 2009), and conservation biology (WARD et al. 2008, NEIGEL et al. 2008). New insights into ecology and species biology have already emerged from various DNA barcode studies (FRÉZAL & LEBLOIS 2008). For example, the identification of organisms found in stomach extracts allows the elucidation of wild animal diets, especially when behavioural studies are not feasible (see VALENTINI et al. 2008). In this context, newly emerging sequencing techniques such as pyrosequencing (454 Roche, Solexa, SOLID) enable rapid and representative analyses of mixed samples (e.g. stomach contents, food, blood or water columns) (VALENTINI et al. 2008). Moreover, the emerging field of metagenomics will also support future DNA barcoding initiatives (HUDSON 2008).

Our study confirms the usefulness of DNA barcodes for the identification of species of the genus *Bembidion*. We found a clear barcoding gap that allows a firm identification of all analysed species, including closely related species pairs such as *Bembidion lampros/properans* and *Bembidion millerianum/ruficorne* (see Figure 2). We also gained some first impressions of the intraspecific variability of various *Bembidion* species. Of course, it is clear that additional specimens and species have to be analysed to evaluate intra- and interspecific COI variability in more detail in the near future. To summarize the results, our study confirms the high potential of DNA barcodes for a successful species identification of even closely related ground beetle species of the genus *Bembidion* in Central Europe.

Acknowledgments

We thank CHRISTINA BLUME and CLAUDIA ETZBAUER for their laboratory assistance. We are very grateful to BERNHARD MISOF for donating specimens and ORTWIN BLEICH for giving permission to use photos of carabid specimens from www.eurocarabidae.de. MICHAEL J. RAUPACH was supported by a research grant from the German Science Foundation (DFG) for JOHANN-WOLFGANG WÄGELE (INST 427/1-1).

Literature

- ALTSCHUL, S.F., GISH, W., MYERS, W.M.E.W. & D.J. LIPMAN (1990): Basic local alignment search tool. – *Journal of Molecular Biology* 215: 403–410.
- ARMSTRONG, K.F. & S.L. BALL (2005): DNA barcodes for biosecurity: invasive species control. – *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 360: 1813–1823.
- ARNDT, E., BEUTEL, R.G. & K. WILL (2005): Carabidae Latreille, 1802. – In: BEUTEL, R.G. & R.A.B. LESCHEN (Hrsg.) (2005): *Handbuch der Zoologie/Handbook of Zoology*. Vol. IV Arthropoda: Insecta. Part 38. Coleoptera, Beetles. Vol 1: Morphology and Systematics (Archostemata, Adephaga, Myxophaga, Polyphaga partim). – Berlin (deGruyter): 119–146.
- ARNETT, R.H. & M.C. THOMAS (2000): *American Beetles, Volume 1: Archostemata, Myxophaga, Adephaga, Polyphaga: Staphyliniformia*. – xvi + 443 S.; CRC Press LLC, Boca Raton, Florida.
- AVGIN, S.S. & M.L. LUFF (2010): Ground beetles (Coleoptera: Carabidae) as bioindicators of human impact. – *Munis Entomology & Zoology* 5: 209–215.
- BENSASSON, D., ZHANG, D.-X. & G.M. HEWITT (2000): Frequent assimilation of mitochondrial DNA by grasshopper nuclear genomes. – *Molecular Biology and Evolution* 17: 406–415.
- BORISENKO A.V., SONES, J.E. & P.D.N. HEBERT (2009): The front-end logistics of DNA barcoding: challenges and prospects. – *Molecular Ecology Resources* 9: 27–34.
- BUCKLIN, A., HOPCROFT, R.R., KOSOBOKOVA, K.N., NIGRO, L.M., ORTMAN, B.D., JENNINGS, R.M. & C.J. SWEETMAN (2010): DNA barcoding of Arctic Ocean holozooplankton for species identification and recognition. – *Deep-Sea Research Part II: Topical Studies in Oceanography* 57: 40–48.
- CLARE, E.L., LIM, B.K., ENGSTROM, M.D., EGER, J.L. & P.D.N. HEBERT (2007): DNA barcoding of neotropical bats: species identification and discovery within Guyana. – *Molecular Ecology Notes* 7: 184–190.
- CLARE, E.L., KERR, K.C.R., VON KÖNIGSLÖW, T.E., WILSON, J.J. & P.D.N. HEBERT (2008): Diagnosing mitochondrial DNA diversity: applications of a sentinel gene approach. – *Journal of Molecular Evolution* 66: 362–367.
- COSTA, F.O., DEWAARD, J.R., BOUTILLIER, J., RATNASINGHAM, S., DOOH, R.T., HAJIBABAEI, M. & P.D.N. HEBERT (2007): Biological identifications through DNA barcodes: the case of the Crustacea. – *Canadian Journal of Fisheries and Aquatic Sciences* 64: 272–295.
- COSTA, F.O., HENZLER, C.M., LUNT, D.H., WHITELEY, N.M. & J. ROCK (2010): Probing marine *Gammarus* (Amphipoda) taxonomy with DNA barcodes. – *Systematics and Biodiversity* 7: 365–379.
- CROUCHER, P.J.P., OXFORD, G.S. & J.B. SEARLE (2004): Mitochondrial differentiation, introgression and phylogeny in species in the *Tegenaria atrica* group (Araneae: Agelenidae). – *Biological Journal of the Linnean Society* 81: 79–89.
- DARLING, J.A. & M.J. BLUM (2007): DNA-based methods for monitoring invasive species: a review and prospectus. – *Biological Invasions* 9: 751–765.
- DA SILVA, J.M., CREER, S., DOS SANTOS, A., COSTA, A.C., CUNHA, M.R., COSTA, F.O. & G.R. CARVALHO (2011): Systematic and evolutionary insight derived from mtDNA COI barcode diversity in the Decapoda (Crustacea: Malacostraca). – *Public Library of Science ONE* 6: e19449.
- DAWNAY, N., OGDEN, R., MCEWING, R., CARVALHO, G.R. & R.S. THORPE (2007): Validation of the barcoding gene COI for the use in forensic genetic species identification. – *Forensic Science International* 173: 1–6.
- DESALLE, R., EGAN, M.G. & M.E. SIDDALL (2005): The unholy trinity: taxonomy, species delimitation and DNA barcoding. – *Proceedings of the Royal Society of London Series B: Biological Science* 360: 1905–1916.
- DESENDER, K. & D. CRAPPE (1983): Larval and adult morphology and biometry of two sibling species *Bembidion lampros* (Herbst) and *Bembidion properans* Stephens (Coleoptera, Carabidae). – *Biologisch Jaarboek Dodonaea* 51: 36–54.
- DRUMMOND, A.J., ASHTON, B., BUXTON, S., CHEUNG, M., COOPER, A., HELED, J., KEARSE, M., MOIR, R., STONES-HAVAS, S., STURROCK, S., THIERER, T. & A. WILSON (2010): Geneious v5.3, available from www.geneious.com.
- DURON, O., BOUCHON, D., BOUTIN, S., BELLAMY, L., ZHOU, L., ENGELSTÄDTER, J. & G.D.D. HURST (2008): The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. – *BMC Biology* 6: 27.
- EBACH, M.C. (2011): Taxonomy and the DNA barcoding enterprise. – *Zootaxa* 2742: 67–68.
- EDGAR, R.C. (2004): MUSCLE: a multiple sequence alignment method with reduced time and space complexity. – *BMC Bioinformatics* 5: 113.
- ENGSTRAND, R.C., TOVAR, J.C., CIBRIAN-JARAMILLO, A. & S.-O. KOLOKOTRONIS (2010): Genetic variation in avocado stem weevils *Copturus aguacatae* (Coleoptera: Curculionidae) in Mexico. – *Mitochondrial DNA* 21: 38–43.
- FELSENSTEIN, J. (1985): Confidence limits on phylogenies: an approach using the bootstrap. – *Evolution* 39: 783–791.
- FERRI, G., ALU, M., CORRADINI, B. & G. BEDUSCHI (2009): Forensic botany: species identification of botanical trace evidence using a multigene barcoding approach. – *International Journal of Legal Medicine* 123: 395–401.
- FOLMER, O., BLACK, M., HOEH, W., LUTZ, R. & R. VRIEJENHOEK (1994): DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. – *Molecular Marine Biology and Biotechnology* 3: 294–299.
- FREZAL, L. & R. LEBLOIS (2008): Four years of DNA barcoding: current advances and prospects. – *Infection, Genetics and Evolution* 8: 727–736.
- GREENSTONE, M.H., ROWLEY, D.L., HEIMBACH, U., LUNDGREN, J.G., PFANNENSTIEL, R.S. & S.A. REHNER (2005): Barcoding generalist predators by polymerase chain reaction: carabids and spiders. – *Molecular Ecology* 14: 3247–3266.
- HAJIBABAEI, M., JANZEN, D.H., BURNS, J.M., HALLWACHS, W. & P.D.N. HEBERT (2006): DNA barcodes distinguish species of tropical Lepidoptera. – *Proceedings of the National Academy of Sciences of the United States of America* 103: 968–971.
- HAMMER, Ø., HARPER, D.A.T. & P.D. RYAN (2001): PAST: Palaeontological statistics software package for education and data analysis. – *Palaeontologia Electronica* 4: 1–9.
- HAUSMANN, A., HASZPRUNAR, G. & P.D.N. HEBERT (2011): DNA barcoding the Geometrid Fauna of Bavaria (Lepidoptera): successes, surprises, and questions. – *Public Library of Science ONE* 6: e17134.
- HEBERT, P.D.N., CYWINSKA, A., BALL, S.L. & J.R. DEWAARD (2003A): Biological identifications through DNA barcodes. – *Proceedings of the Royal Society of London Series B: Biological Science* 270: 313–321.
- HEBERT, P.D.N., RATNASINGHAM, S. & J.R. DEWAARD (2003B): Barcoding animal life: cytochrome c oxidase subunit I divergences among closely related species. – *Proceedings of the Royal Society of London Series B: Biological Science* 270 (Supplement): S96–S99.
- HEBERT, P.D.N., PENTON, E.H., BURNS, J.M., JANZEN, D.H. & W. HALLWACHS (2004): Ten species in one: DNA barcoding reveals

- cryptic species in the neotropical skipper butterfly *Astrapes fulgerator*. – Proceedings of the National Academy of Science of the United States of America 101: 14812–14817.
- HEBERT, P. D. N., DEWAARD, J.R. & J.-F. LANDRY (2010): DNA barcodes for 1/1000 of the animal kingdom. – *Biology Letters* 6: 359–362.
- HUBERT, N., HANNER, R., HOLM, E., MANDRAK, N.E., TAYLOR, E., BURRIDGE, N.E., WATKINSON, D., DUMONT, P., CURRY, A., BENTZEN, P., ZHANG, J., APRIL, J. & L. BERNATCHEZ (2008): Identifying Canadian freshwater fishes through DNA barcodes. – *Public Library of Science ONE* 3: e2490.
- HUDSON, M.E. (2008): Sequencing breakthroughs for genomic ecology and evolutionary biology. – *Molecular Ecology Resources* 8: 3–17.
- HURKA, K. (1996): Carabidae of the Czech and Slovak Republics. – 565 S.; Kabourek, Zlín.
- IVANOVA, N.V., BORISENKO, A.V. & P.D.N. HEBERT (2009): Express barcodes: racing from specimen to identification. – *Molecular Ecology Resources* 9: 35–41.
- JINBO, U., KATO, T. & M. ITO (2011): Current progress in DNA barcoding and future implications for entomology. – *Entomological Science* 14: 107–124.
- JOHNSON, S.B., WARÉN, A. & R.C. VRIJENHOEK (2008): DNA barcoding of *Lepetodrilus* limpets reveals cryptic species. – *Journal of Shellfish Research* 27: 43–51.
- KIM, J.-H., ANTUNES, A., LUO, S.-J., MENNINGER, J., NASH, W.G., O'BRIEN, S.J. & W.E. JOHNSON (2006): Evolutionary analysis of a large mtDNA translocation (numt) into the nuclear genome of the *Panthera* genus species. – *Gene* 366: 292–302.
- KIMURA, M. (1980): A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. – *Journal of Molecular Evolution* 16: 111–120.
- KLAUSNITZER, B. (2010): Entomologie – quo vadis? – *Nachrichtenblatt der bayrischen Entomologen* 59: 99–111.
- KOUTROUMPA, F.A., LIEUTIER, F. & G. ROUX-MORABITO (2009): Incorporation of mitochondrial fragments in the nuclear genome (numts) of the longhorned beetle *Monochamus galloprovincialis* (Coleoptera, Cerambycidae). – *Journal of Zoological Systematics and Evolutionary Research* 47: 141–148.
- LAKRA, W.S., VERMA, M.S., GOSWAMI, M., LAL, K.K., MOHINDRA, V., PUNIA, P., GOPALAKRISHAN, A., SINGH, K.V., WARD, R.D. & P.D.N. HERBERT (2011): DNA barcoding Indian marine fishes. – *Molecular Ecology Resources* 11: 60–71.
- LINARES, M.C., SOTO-CALDERON, I.D., LEES, D.C. & N.M. ANTHONY (2009): High mitochondrial diversity in geographically widespread butterflies of Madagascar: a test of the DNA barcoding approach. – *Molecular Phylogenetics and Evolution* 50: 485–495.
- LINDROTH, C.H. (1985): The Carabidae (Coleoptera) of Fennoscandia and Denmark. – *Fauna Entomologica Scandinavia* 15: 1–232.
- LÖVEI, G.L. & K.D. SUNDERLAND (1996): Ecology and behaviour of ground beetles (Coleoptera: Carabidae). – *Annual Reviews in Entomology* 41: 231–256.
- MACHIDA, R.J., HASHIGUCHI, Y., NISHIDA, M. & S. NISHIDA (2009): Zooplankton diversity analysis through single-gene sequencing of a community sample. – *BMC Genomics* 10: 438.
- MADDISON, D.R. (2008): Systematics of the North American beetle subgenus *Pseudoperlyphus* (Coleoptera: Carabidae: *Bembidion*) based on morphological, chromosomal, and molecular data. – *Annals of Carnegie Museum* 77: 147–193.
- MEIKLEJOHN, K.A., WALLMAN, J.F. & M. DOWTON (2011): DNA-based identification of forensically important Australian Sarcophagidae (Diptera). – *International Journal of Legal Medicine* 125: 27–32.
- MILLER, S.E. (2007): DNA barcoding and the renaissance of taxonomy. – *Proceedings of the National Academic Sciences of the United States of America* 104: 4775–4776.
- MIN, X.-J. & D.A. HICKEY (2007): DNA barcodes provide a quick preview of mitochondrial genome composition. – *Public Library of Science ONE* 3: e325.
- MÜLLER, G. (1971): *Bembidion lampros* Hrbst. und *B. properans* Steph. – zwei Arten? – *Entomologische Nachrichten* 15: 119–124.
- MÜLLER-MOTZFELD, G. (1982): Zur Trennung von *Bembidion ruficornis* Sturm und *millerianum* Heyden (Col., Carabidae). – *Entomologische Nachrichten und Berichte* 26: 121–125.
- MÜLLER-MOTZFELD, G. (HRSG.) (2004): Bd. 2, Adephega 1: Carabidae (Laufkäfer). – In: FREUDE, H., HARDE, K.W., LOHSE, G.A. & B. KLAUSNITZER: Die Käfer Mitteleuropas. – 521 S.; Spektrum-Verlag (Heidelberg/Berlin), 2. Auflage.
- NEIGEL, J., DOMINGO A. & J. STAKE (2007): DNA barcoding as a tool for coral reef conservation. – *Coral Reefs* 26: 487–499.
- NELSON, L.A., WALLMAN, J.F. & M. DOWTON (2007): Using COI barcodes to identify forensically and medically important blowflies. – *Medical and Veterinary Entomology* 21: 44–52.
- RADULOVICI, A.E., SAINTE-MARIE, B. & F. DUFRESNE (2009): DNA barcoding of marine crustaceans from the Estuary and Gulf of St Lawrence: a regional-scale approach. – *Molecular Ecology Resources* 9: 181–187.
- RAINIO, J. & J. NIEMELÄ (2003): Ground beetles (Coleoptera: Carabidae) as bioindicators. – *Biodiversity and Conservation* 12: 487–506.
- RATNASINGHAM, S. & P.D.N. HEBERT (2007) BOLD: The Barcode of Life Data Systems. – *Molecular Ecology Notes* 7: 355–364.
- RAUPACH, M.J., ASTRIN, J.J., HANNIG, K., PETERS, M.K., STOECKLE, M.Y. & J.-W. WÄGELE (2010A): Molecular species identifications of Central European ground beetles (Coleoptera: Carabidae) using nuclear rDNA expansion segments and DNA barcodes. – *Frontiers in Zoology* 7: 26.
- RAUPACH, M.J., HANNIG, K. & J.-W. WÄGELE (2010B): DNA Barcoding – Perspektiven und Chancen: Eine Fallstudie am Beispiel der Laufkäfer (Coleoptera: Carabidae). – *Entomologie heute* 22: 171–190.
- REITTER, E. (1908): *Fauna Germanica. Die Käfer des Deutschen Reiches*. 1. Band – Stuttgart.
- RIVERA, J. & D.C. CURRIE (2009): Identification of Nearctic black flies using DNA barcodes (Diptera: Simuliidae). – *Molecular Ecology Resources* 9 (Suppl. 1): 224–236.
- ROBINSON, E.A., BLAGOEV, G.A., HEBERT, P.D.N. & S.J. ADAMOWICZ (2009): Prospects for using DNA barcoding to identify spiders in species-rich genera. – *ZooKeys* 16: 27–46.
- SAITOU, N. & M. NEI (1987): The neighbour-joining method: a new method for reconstructing phylogenetic trees. – *Molecular Biology and Evolution* 4: 406–425.
- SMITH, M.A. & B.L. FISHER (2009): Invasions, DNA barcodes, and rapid biodiversity assessment using ants of Mauritius. – *Frontiers in Zoology* 6: 31.
- SOTA, T., ISHIKAWA, R., UJIE, M., KUSUMOTO, F. & A.P. VOGLER (2001): Extensive trans-species mitochondrial polymorphisms in the carabid beetles *Carabus* subgenus *Ohomopterus* caused by repeated introgressive hybridization. – *Molecular Ecology* 10: 2833–2847.
- STOECKLE, M.Y. & P.D.N. HEBERT (2008): Barcode of Life. – *Scientific American* 299: 82–88.
- STRUTZENBERGER, P., BREHM, G. & K. FIEDLER (2011): DNA barcoding-based species delimitation increases species count of *Eois* (Geometridae) moths in a well-studied tropical mountain forest up to 50%. – *Insect Science* 18: 349–362.

- TAMURA, K., PETERSON, D., PETERSON, N., STECHER, G., NEI, M. & S. KUMAR (2011): MEGA5: Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and Maximum Parsimony Methods. – *Molecular Biology and Evolution* (in press): doi:10.1093/molbev/msr121).
- TAUTZ, D., ARCTANDER, P., MINELLI, A., THOMAS, R.H. & A.P. VOGLER (2003): A plea for DNA taxonomy. – *Trends in Ecology and Evolution* 18: 70–74.
- UTHICKE, S., BYRNE, M. & C. CONAND (2010): Genetic barcoding of commercial Bêche-de-mer species (Echinodermata: Holothuroidea). – *Molecular Ecology Resources* 10: 634–646.
- VALENTINI, A., POMPANON, F. & P. TABERLET (2008): DNA barcoding for ecologists. – *Trends in Ecology and Evolution* 24: 110–117.
- VOGLER, A.P. & M.T. MONAGHAM (2007): Recent advantages in DNA taxonomy. – *Journal of Systematics and Evolutionary Research* 45: 1–10.
- WARD, R.D., HOLMES, B.H., WHITE, W.T. & P.R. LAST (2008): DNA barcoding Australasian chondrichthyan: results and potential uses in conservation. – *Marine and Freshwater Research* 59: 57–71.
- WEI, S.-J., SHI, M., CHEN, X.-X., SHARKEY, M.J., VAN ACHTERBERG, C., YE, G.-Y. & J.-H. HE (2010): New views on strand asymmetry in insect mitochondrial genomes. – *Public Library of Science ONE* 5: e12708.
- WERREN, J.H., BALDO, L. & M.E. CLARK (2008): *Wolbachia*: master manipulators of invertebrate biology. – *Nature Reviews* 6: 741–751.
- WILL, K.P., MISHLER, P.D. & Q.D. WHEELER (2005): The perils of DNA barcoding and the need for integrative taxonomy. – *Systematic Biology* 54: 844–851.
- ZEMLAJ, T.S., WARD, R.D., CONNELL, A.D., HOLMES, B.H. & P.D.N. HEBERT (2009): DNA barcoding reveals overlooked marine fishes. – *Molecular Ecology Resources* 9: 237–241.
- ZHANG, A.-B. & T. SOTA (2007): Nuclear gene sequences resolve species phylogeny and mitochondrial introgression in *Leptocarabus* beetles showing trans-species polymorphisms. – *Molecular Phylogenetics and Evolution* 45: 534–546.
- ZHOU, X., ADAMOWICZ, S.J., JACOBUS, L.M., DEWALT, R.E. & P.D.N. HEBERT (2009): Towards a comprehensive barcode library for arctic life – Ephemeroptera, Plecoptera, and Trichoptera of Churchill, Manitoba, Canada. – *Frontiers in Zoology* 6: 30.

Manuskripteingang: 2. 5. 2011

Author's addresses

Dr. Michael J. Raupach
Senckenberg am Meer
Deutsches Zentrum für Marine Biodiversitätsforschung
AG Molekulare Taxonomie mariner Organismen
Südstrand 44
26382 Wilhelmshaven
e-mail: mraupach@senckenberg.de

Dipl.-Biol. Karsten Hannig
Dresdener Straße 6
45731 Waltrop
e-mail: karsten.hannig@gmx.de

Prof. Dr. Johann-Wolfgang Wägele
Zoologisches Forschungsmuseum Alexander Koenig
Adenauerallee 160-162
53113 Bonn
e-mail: w.waegel.zfmk@uni-bonn.de

Tab. 1: Species identification, voucher code, location and year of collection as well as GenBank accession numbers of the 101 analysed *Bembidion* specimens. Country codes: GER (Germany): NW (North-Rhine Westphalia), NI (Lower Saxony), SH (Schleswig-Holstein), RP (Rhineland-Palatinate), ST (Saxony-Anhalt); AU (Austria): K (Carinthia).

Species	ZFMK voucher code	Locality	COI accession number
<i>Bembidion articulatum</i> (Panzer, 1796)	ZFMK_COL_2008_44	Bienen, NW, GER, 2006	GU347065
<i>Bembidion articulatum</i> (Panzer, 1796)	ZFMK_COL_2008_99	Waltrop, NW, GER, 1999	GU347068
<i>Bembidion articulatum</i> (Panzer, 1796)	ZFMK_COL_2008_119	Waltrop, NW, GER, 1999	GU347069
<i>Bembidion articulatum</i> (Panzer, 1796)	ZFMK_COL_2008_159	Waltrop, NW, GER, 1999	GU347067
<i>Bembidion articulatum</i> (Panzer, 1796)	ZFMK_COL_2009_38	Schmeddehausen, NW, GER, 2008	GU347066
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2009_350	Meindorf, NW, GER, 2008	JF895158
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2009_351	Meindorf, NW, GER, 2008	JF895159
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2009_352	Neheim, NW, GER, 2008	JF895160
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2011_292	Arnsberg, NW, GER, 2008	JF895155
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2011_293	Arnsberg, NW, GER, 2008	JF895156
<i>Bembidion atrocaeruleum</i> (Stephens, 1828)	ZFMK_COL_2011_294	Arnsberg, NW, GER, 2008	JF895157
<i>Bembidion aspericolle</i> (Panzer, 1796)	ZFMK_COL_2008_23	Teutschenthal, ST, GER, 2005	GU347070
<i>Bembidion aspericolle</i> (Panzer, 1796)	ZFMK_COL_2008_24	Teutschenthal, ST, GER, 2005	GU347072
<i>Bembidion aspericolle</i> (Panzer, 1796)	ZFMK_COL_2008_25	Teutschenthal, ST, GER, 2005	GU347073
<i>Bembidion aspericolle</i> (Panzer, 1796)	ZFMK_COL_2008_26	Teutschenthal, ST, GER, 2005	GU347074
<i>Bembidion aspericolle</i> (Panzer, 1796)	ZFMK_COL_2008_27	Teutschenthal, ST, GER, 2005	GU347071
<i>Bembidion cruciatum</i> Dejean, 1831	ZFMK_COL_2009_355	Feistritz, K, AU, 2008	JF895169
<i>Bembidion cruciatum</i> Dejean, 1831	ZFMK_COL_2009_356	Feistritz, K, AU, 2008	JF895170
<i>Bembidion cruciatum</i> Dejean, 1831	ZFMK_COL_2009_357	Feistritz, K, AU, 2008	JF895171
<i>Bembidion cruciatum</i> Dejean, 1831	ZFMK_COL_2009_358	Maria Elend, K, AU, 2008	JF895172
<i>Bembidion decoratum</i> (Dufschmid, 1812)	ZFMK_COL_2009_361	Maria Elend, K, AU, 2008	GU347075
<i>Bembidion decoratum</i> (Dufschmid, 1812)	ZFMK_COL_2009_362	Maria Elend, K, AU, 2008	GU347076
<i>Bembidion decoratum</i> (Dufschmid, 1812)	ZFMK_COL_2009_363	Maria Elend, K, AU, 2008	GU347077
<i>Bembidion decoratum</i> (Dufschmid, 1812)	ZFMK_COL_2009_364	Maria Elend, K, AU, 2008	GU347078
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2008_12	Neheim, NW, GER, 2004	GU347079
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2008_13	Neheim, NW, GER, 2004	GU347081
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2008_14	Neheim, NW, GER, 2004	GU347082
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2008_16	Bienen, NW, GER, 2004	GU347083
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2008_42	St. Augustin-Meindorf, NW, GER, 2008	GU347080
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2009_360	St. Augustin-Meindorf, NW, GER, 2008	GU347085
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2009_400	St. Augustin-Meindorf, NW, GER, 2008	GU347084
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2009_498	St. Augustin-Meindorf, NW, GER, 2008	GU347086
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2009_499	St. Augustin-Meindorf, NW, GER, 2008	GU347087
<i>Bembidion decorum</i> (Panzer, 1799)	ZFMK_COL_2009_500	Bienen, NW, GER, 2008	GU347088
<i>Bembidion deletum</i> Audinet-Serville, 1821	ZFMK_COL_2009_366	Hochobir, K, AU, 2008	JF895161
<i>Bembidion deletum</i> Audinet-Serville, 1821	ZFMK_COL_2009_367	Hochobir, K, AU, 2008	JF895162
<i>Bembidion deletum</i> Audinet-Serville, 1821	ZFMK_COL_2009_368	Hochobir, K, AU, 2008	JF895163
<i>Bembidion elongatum</i> Dejean, 1831	ZFMK_COL_2009_40	Schmeddehausen, NW, GER, 2008	GU347089
<i>Bembidion elongatum</i> Dejean, 1831	ZFMK_COL_2009_41	Schmeddehausen, NW, GER, 2008	GU347090
<i>Bembidion elongatum</i> Dejean, 1831	ZFMK_COL_2009_42	Schmeddehausen, NW, GER, 2008	GU347091
<i>Bembidion elongatum</i> Dejean, 1831	ZFMK_COL_2009_43	Schmeddehausen, NW, GER, 2008	GU347092
<i>Bembidion incognitum</i> Müller, 1931	ZFMK_COL_2009_370	Gurktaler Alpen, K, AU, 2008	JF895164
<i>Bembidion incognitum</i> Müller, 1931	ZFMK_COL_2009_371	Gurktaler Alpen, K, AU, 2008	JF895165
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2008_122	Borkum, NI, GER, 2005	GU347094
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2008_142	Arnsberg, NW, GER, 2007	GU347096
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2008_146	Haltern-Lavesum, NW, GER, 1999	GU347097
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_53	Schmeddehausen, NW, GER, 2008	GU347101
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_54	Schmeddehausen, NW, GER, 2008	GU347102
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_611	Teutschenthal, ST, GER, 2008	GU347098
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_612	Teutschenthal, ST, GER, 2008	GU347093
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_613	Teutschenthal, ST, GER, 2008	GU347099
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_614	Teutschenthal, ST, GER, 2008	GU347100
<i>Bembidion lampros</i> (Herbst, 1787)	ZFMK_COL_2009_644	Wyk, SH, GER, 2003	GU347095
<i>Bembidion litorale</i> (Olivier, 1790)	ZFMK_COL_2009_59	Greven-Bockholt, NW, GER, 2008	GU347103
<i>Bembidion litorale</i> (Olivier, 1790)	ZFMK_COL_2009_60	Greven-Bockholt, NW, GER, 2008	GU347105
<i>Bembidion litorale</i> (Olivier, 1790)	ZFMK_COL_2009_61	Greven-Bockholt, NW, GER, 2008	GU347106
<i>Bembidion litorale</i> (Olivier, 1790)	ZFMK_COL_2009_62	Greven-Bockholt, NW, GER, 2008	GU347104
<i>Bembidion millerianum</i> Heyden, 1883	ZFMK_COL_2009_372	Neheim, NW, GER, 2008	JF895177
<i>Bembidion millerianum</i> Heyden, 1883	ZFMK_COL_2009_373	Neheim, NW, GER, 2008	JF895178
<i>Bembidion millerianum</i> Heyden, 1883	ZFMK_COL_2009_374	Neheim, NW, GER, 2008	JF895180
<i>Bembidion millerianum</i> Heyden, 1883	ZFMK_COL_2009_375	Neheim, NW, GER, 2008	JF895179

Species	ZFMK voucher code	Locality	COI accession number
<i>Bembidion pallidipenne</i> (Illiger, 1802)	ZFMK COL 2009 67	Fehmarn, SH, GER, 2008	GU347107
<i>Bembidion pallidipenne</i> (Illiger, 1802)	ZFMK COL 2009 68	Fehmarn, SH, GER, 2008	GU347108
<i>Bembidion pallidipenne</i> (Illiger, 1802)	ZFMK COL 2009 69	Fehmarn, SH, GER, 2008	GU347109
<i>Bembidion pallidipenne</i> (Illiger, 1802)	ZFMK COL 2009 70	Fehmarn, SH, GER, 2008	GU347110
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 75	Castrop-Rauxel, NW, GER, 2008	GU347116
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 76	Castrop-Rauxel, NW, GER, 2008	GU347117
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 77	Castrop-Rauxel, NW, GER, 2008	GU347112
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 615	Teutschenthal, ST, GER, 2008	GU347111
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 616	Teutschenthal, ST, GER, 2008	GU347113
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 617	Teutschenthal, ST, GER, 2008	GU347114
<i>Bembidion properans</i> (Stephens, 1828)	ZFMK COL 2009 618	Teutschenthal, ST, GER, 2008	GU347118
<i>Bembidion punctulatum</i> Drapiez, 1820	ZFMK COL 2008 39	Bienen, NW, GER, 2006	GU347136
<i>Bembidion punctulatum</i> Drapiez, 1820	ZFMK COL 2009 382	Dessau, SA, GER, 2008	GU347139
<i>Bembidion punctulatum</i> Drapiez, 1820	ZFMK COL 2009 383	Dessau, SA, GER, 2008	GU347138
<i>Bembidion punctulatum</i> Drapiez, 1820	ZFMK COL 2009 384	Neheim, NW, GER, 2008	GU347137
<i>Bembidion pygmaeum</i> (Fabricius, 1792)	ZFMK COL 2009 385	Halle an der Saale, ST, GER, 2008	JF895173
<i>Bembidion pygmaeum</i> (Fabricius, 1792)	ZFMK COL 2009 386	Halle an der Saale, ST, GER, 2008	JF895174
<i>Bembidion pygmaeum</i> (Fabricius, 1792)	ZFMK COL 2009 387	Halle an der Saale, ST, GER, 2008	JF895175
<i>Bembidion pygmaeum</i> (Fabricius, 1792)	ZFMK COL 2009 388	Halle an der Saale, ST, GER, 2008	JF895176
<i>Bembidion ruficornne</i> Sturm, 1825	ZFMK COL 2009 391	Feistritz, K, AU, 2008	GU347118
<i>Bembidion ruficornne</i> Sturm, 1825	ZFMK COL 2009 392	Feistritz, K, AU, 2008	GU347119
<i>Bembidion ruficornne</i> Sturm, 1825	ZFMK COL 2009 393	Feistritz, K, AU, 2008	GU347120
<i>Bembidion ruficornne</i> Sturm, 1825	ZFMK COL 2009 394	Feistritz, K, AU, 2008	GU347121
<i>Bembidion semipunctatum</i> (Donovan, 1806)	ZFMK COL 2008 43	Bienen, NW, GER, 2006	GU347123
<i>Bembidion semipunctatum</i> (Donovan, 1806)	ZFMK COL 2009 396	Bienen, NW, GER, 2008	GU347122
<i>Bembidion semipunctatum</i> (Donovan, 1806)	ZFMK COL 2009 397	Bienen, NW, GER, 2008	GU347124
<i>Bembidion semipunctatum</i> (Donovan, 1806)	ZFMK COL 2009 398	Bienen, NW, GER, 2008	GU347125
<i>Bembidion testaceum</i> (Duftschmid, 1812)	ZFMK COL 2008 40	Bienen, NW, GER, 2006	JF895166
<i>Bembidion testaceum</i> (Duftschmid, 1812)	ZFMK COL 2009 401	Feistritz, K, AU, 2008	JF895167
<i>Bembidion testaceum</i> (Duftschmid, 1812)	ZFMK COL 2009 402	Feistritz, K, AU, 2008	JF895168
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 90	Schmeddehausen, NW, GER, 2008	GU347126
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 91	Schmeddehausen, NW, GER, 2008	GU347130
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 92	Schmeddehausen, NW, GER, 2008	GU347127
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 93	Schmeddehausen, NW, GER, 2008	GU347128
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 94	Schmeddehausen, NW, GER, 2008	GU347129
<i>Bembidion tetracolum</i> Say, 1823	ZFMK COL 2009 601	Remagen, RP, GER, 2008	GU347131
<i>Bembidion tibiale</i> (Duftschmid, 1812)	ZFMK COL 2009 404	Nachrode-Wiblingwerde, NW, GER, 2008	GU347132
<i>Bembidion tibiale</i> (Duftschmid, 1812)	ZFMK COL 2009 405	Nachrode-Wiblingwerde, NW, GER, 2008	GU347133
<i>Bembidion tibiale</i> (Duftschmid, 1812)	ZFMK COL 2009 406	Zell-Mitterwinkel, K, AU, 2008	GU347134
<i>Bembidion tibiale</i> (Duftschmid, 1812)	ZFMK COL 2009 407	Zell-Mitterwinkel, K, AU, 2008	GU347135

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Angewandte Carabidologie](#)

Jahr/Year: 2012

Band/Volume: [9](#)

Autor(en)/Author(s): Hannig Karsten, Raupach Michael J., Wägele Johann Wolfgang

Artikel/Article: [Identification of Central European ground beetles of the genus *Bembidion* \(Coleoptera: Carabidae\) using DNA barcodes: A case study of selected species 63-72](#)