

Zodarion ohridense (Araneae: Zodariidae) – a new record for Central Europe

Tomáš Krejčí, Milan Řezáč & Tomáš Kadlec



doi: 10.5431/aramit5402

Abstract. *Zodarion ohridense* Wunderlich, 1973 was found to be introduced in an abandoned stone quarry near Kolín in the Czech Republic. The Central European species of *Zodarion* are discussed.

Keywords: Czech Republic, faunistics, introduced species, new record, quarry

Zusammenfassung. *Zodarion ohridense* (Araneae: Zodariidae) – ein Neunachweis für Mitteleuropa. *Zodarion ohridense* Wunderlich, 1973 wurde in einen aufgelassenen Steinbruch bei Kolín, Tschechische Republik, eingeschleppt. Die mitteleuropäischen *Zodarion*-Arten werden diskutiert.

The family Zodariidae comprises 1126 species in 84 genera. The most species-rich genus, *Zodarion*, comprises 158 species, 100 species of which occur in Europe (World Spider Catalog 2017). Representatives of *Zodarion* are obligatory ant-eaters (each species being specialized on a certain group of ants), shelter themselves in silken retreats camouflaged by soil particles and morphologically, chemically and behaviourally mimic ants (Pekár et al. 2005). The centre of the distribution area for this genus is the Mediterranean: so far only four species – *Zodarion germanicum* (C. L. Koch, 1837), *Zodarion hamatum* Wiehle, 1964, *Zodarion italicum* (Canestrini, 1868) and *Zodarion rubidum* Simon, 1914 – are known to extend into Central Europe (Nentwig et al. 2017). Here we present the record of a fifth species of *Zodarion* in Central Europe.

Material and methods

Zodarion ohridense Wunderlich, 1973. CZECH REPUBLIC, Elbe Lowland, Nová Ves I, 3 km northwest of Kolín, natural monument Lom u Nové Vsi (abandoned stone quarry), 50.0549°N, 15.1323°E (grid no. 5956), 220 m a.s.l., 1 male, pitfall trap, 1–30 June 2015, leg. Jiří Skala, det. Milan Řezáč & Tomáš Krejčí. The male was preserved in 75 % ethanol and photographed using a HIROX RH-2000 digital microscope. The material is deposited in the collection of the second author at the Crop Research Institute, Prague.

Results

Main morphological features. The male of *Zodarion ohridense* is characterised by the following morphological features (Bosmans 2009, Nentwig et al. 2017): Body length 2.5–3.2 mm. Prosoma (Fig. 1a) reddish brown, fovea and margin darker. Legs yellowish brown, femora dark brown. Opisthosoma dorsally dark sepia brown, ventrally with whitish oval spot. Palp (Fig. 1b, c) with rather short tibial apophysis, with broad base, terminally pointed; tegulum with angularity at base; median apophysis with two large prolateral teeth; embolus linear, with subterminal bend.

The colouration of *Z. ohridense* is very similar to *Z. germanicum*, *Z. italicum* and *Z. hamatum* but it is easy to distinguish these species according to the shape of the copulatory organs

(Bosmans 1997, 2009). Only *Z. rubidum* distinctly differs in colouration from other species. It has a light brown prosoma and legs, and the opisthosoma is brown-violet (Bosmans 1997).

Natural history. *Zodarion ohridense* was described from the Ohrid region in Macedonia (Wunderlich 1973) and was considered an endemic species of the Balkan Peninsula. So far it is known from Croatia (Bosmans 2009), Bulgaria (Blagoev et al. 2001, Tzonev & Lazarov 2001, Deltshev 2004, Lazarov 2007), Greece (Bosmans 2009) and Macedonia (Komnenov 2002, 2003, Lazarov 2004, Bosmans 2009) (Fig. 2). In Macedonia it was found in the Shar Mountains (1200–1700 m) in *Acer* sp. and *Quercus cerris* forest and in *Picea abies* and *Fagus sylvatica* forests (Komnenov 2002) and in the Jakupica Mountains (1400–1900 m) on high mountain pastures or in *Fagus sylvatica* forests (Komnenov 2003). Deltshev (2004) found this species in southwestern Bulgaria in a *Pinus* forest (200–250 m). In Greece Bosmans (2009) found it in a rive-rine (430 m) and deciduous forest (600 m) and in grassland (1910 m).

Discussion

In Central Europe four species of the genus *Zodarion* have been found so far. The first, *Z. germanicum*, was described from Germany (Bosmans 1997) and is probably the only *Zodarion* species which can be considered an autochthonous component of the Central European fauna. The other three species are probably not native to Central Europe and because both ballooning ability and terrestrial migration are very low in *Zodarion* spiders, these species are believed to have been introduced into Central Europe via traffic (Bönsel et al. 2000, Pekár 2002, Pekár et al. 2005, Komposch 2009). The genus *Zodarion* is well preadapted for passive human transport thanks to its tolerance for dry environments and especially by attaching the silken shelters masked by soil particles (so-called igloos) to solid objects on the ground (Jocqué 1991). Spiders are thus transported with these objects and, if there are suitable conditions at their final destination, the spiders can establish new populations far from their continuous distribution area (Pekár 2002).

The most frequently found species in the vicinity of railways tracks is *Zodarion rubidum* (Pekár 2002). It was described from the south-eastern French Pyrenees (Simon 1914) and in 1971 it was found for the first time outside France, in central Austria (Wunderlich 1973). In 1979 this species

Tomáš KREJČÍ & Tomáš KADLEC, Faculty of Environmental Sciences, Czech University of Life Sciences Prague, Kamýcká 129, Praha 6 – Suchdol, CZ-165 21, Czech Republic; E-mails: tomesko@seznam.cz; kadlect@fzp.czu.cz
Milan ŘEZÁČ, Biodiversity Lab, Crop Research Institute, Drnovská 507, Praha 6 – Ruzyně, CZ-161 06, Czech Republic; E-mail: rezac@urv.cz

submitted 3.2.2017, accepted 18.4.2017, online 30.4.2017




Fig. 1: *Zodarion ohridense* Wunderlich, 1973; **a.** Male, habitus, dorsal view; **b.** Male left palp, ventral view; **c.** Male left palp, retrolateral view. Scales: 500 µm (a), 100 µm (b, c)

was recorded in the Czech Republic (Kůrka 1981) and in the 1980s also in Germany (Broen & Moritz 1987, Renner 1992, Bosmans 1994), Belgium (Bara 1984, Hermanns & Bastin 1989, Couvreur 1990) and Spain (Bosmans 1994). Thereafter it has been spread across the other western, central and eastern European countries (Pekár 2002) up to the Baltic Sea in the north (Scharff et al. 2007) and Ukraine in the east (Fedorak et al. 2010). It has been also introduced into the USA (Vogel 1968) and Canada (Paquin & Dupérré 2006).

Zodarion italicum and *Z. hamatum* are also often found close to railways (Horak & Kropf 1999, Pekár et al. 2005). *Zodarion hamatum* occurs in Italy, Slovenia and Croatia (Bosmans 1997) and in 1995 it was found for the first time in the southern part of Austria (Horak & Kropf 1999). *Zodarion italicum* was described from Italy (Canestrini 1868) from where it expanded into Western and Central Europe (Bosmans 1997). In the 1970s it was found in Germany (Misioch 1977) and in 2000 on the western boundary of the distribution area,

in the Czech Republic (Řezáč 2002, Pekár et al. 2005). The distribution area of this species is still enlarging and nowadays it is also known from Spain, the Netherlands and Denmark (Nentwig et al. 2017). East European records from Bulgaria (Drensky 1936), Ukraine and Azerbaijan (Dunin 1984, Mikhailov 1997) are erroneous (Deltchev 1987, Bosmans pers. com.). It often occurs in abandoned stone quarries and sand pits (Harvey & Murphy 1985, Řezáč 2002, Heneberg & Řezáč 2014).

Acknowledgements

Tomáš Kadlec was supported by Faculty of Environmental Sciences, CULS Prague (EHP-CZ02-OV-1-027-2015), Milan Řezáč was supported by the Ministry of Agriculture of the Czech Republic (institutional support MZe RO0415).

References

- Bara L 1984 Six nouvelles espèces pour l'aranéofaune belge découvertes dans la région de Treignes. – Bulletin et Annales de la Société Royale d'Entomologie de Belgique 120: 271–276
- Blagoev G, Lazarov S & Deltchev C 2001 Spiders (Araneae in Kresna Gorge (SW Bulgaria). In: Beron P (ed.) Biodiversity of Kresna Gorge (SW Bulgaria). National Museum of Natural History, Sofia. pp. 103–107
- Bönsel D, Malten A, Wagner S & Zizka G 2000 Flora, Fauna und Biotoptypen von Haupt- und Güterbahnhof in Frankfurt am Main. – Kleine Senckenberg-Reihe 38: 1–63
- Bosmans R 1994 Revision of the genus *Zodarion* Walckenaer, 1833 in the Iberian Peninsula and Balearic Islands (Araneae, Zodariidae). – Eos 69: 115–142
- Bosmans R 1997 Revision of the genus *Zodarion* Walckenaer, 1833, part II. Western and Central Europe, including Italy (Araneae: Zodariidae). – Bulletin of the British Arachnological Society 10: 265–294
- Bosmans R 2009 Revision of the genus *Zodarion* Walckenaer, 1833, part III. South East Europe and Turkey (Araneae: Zodariidae). – Contributions to Natural History 12: 211–295
- Broen B & Moritz M 1987 Zum Vorkommen von *Zodarion rubidum* Simon, 1914 im Berliner Gebiet. – Deutsche Entomologische Zeitschrift N.F. 34: 155–159 – doi: [10.1002/mmnd.19870340117](https://doi.org/10.1002/mmnd.19870340117)




Fig. 2: Records of *Zodarion ohridense* Wunderlich, 1973. The new record is marked by asterisk. BG – Bulgaria, CZ – Czech Republic, GR – Greece, HR – Croatia, MK – Macedonia. Base map adopted from Bosmans (1997)

Zodarion ohridense – new for Central Europe

- Canestrini G 1868 Nuove aracnidi italiani. – Annuario della Società dei Naturalisti in Modena 3: 190-206
- Couvreur JM 1990 Quelques aspects de la biologie de *Zodarion rubidum* Simon, 1918. – Nieuwsbrief van de Belgische Arachnologische Vereniging 5: 7-15
- Deltshev CD 1987 A critical review of genus *Zodarion* Walckenaer (Araneae: Zodariidae) in Bulgaria. – Acta Zoologica Bulgaria 33: 19-25
- Deltshev CD 2004 Spiders (Araneae) from Sandanski-Petrich Valley (SW Bulgaria). – Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe 80: 71-76 – doi: [10.1002/mmnz.20040800106](https://doi.org/10.1002/mmnz.20040800106)
- Drensky P 1936 Katalog der echten Spinnen (Araneae) der Balkanhalbinsel. – Journal of the Bulgarian Academy of Sciences 32: 1-223
- Dunin PM 1984 Fauna and ecology of the spiders (Aranei) of the Apsheron Peninsula (Azerbaijanian SSR). In: Utochkin A (Ed.) Fauna i ekologija paukoobrasnykh. University, Perm. pp. 45-49
- Fedorik MM, Rudenko SS, Marusik YM & Brushnivska LV 2010 Spiders-gerpetobiontes of Chernivtsi city parks. – Nature Reserves in Ukraine 16: 64-71
- Harvey P & Murphy J 1985 *Zodarion italicum* (Araneae: Zodariidae), a species newly recorded in Britain. – Newsletter of the British Arachnological Society 44: 4
- Heneberg P & Řezáč M 2014 Dry sandpits and gravel-sandpits serve as key refuges for endangered epigaeic spiders (Araneae) and harvestmen (Opiliones) of Central European steppes aeolian sands. – Ecological Engineering 73: 659-670 – doi: [10.1016/j.ecoleng.2014.09.101](https://doi.org/10.1016/j.ecoleng.2014.09.101)
- Hermanns JF & Bastin J 1989 Rapport de l'excursion d'Arabel du 25 juin 1988 aux carrières de Montfort (prov. de Liège). – Nieuwsbrief van de Belgische Arachnologische Vereniging 10: 11-12
- Horak P & Kropf C 1999 Landeskundlich bedeutsame Spinnenfunde in der Steiermark (Arachnida: Araneae). – Mitteilungen des Naturwissenschaftlichen Vereines für Steiermark 129: 253-268
- Jocqué R 1991 A generic revision of the spider family Zodariidae (Araneae). – Bulletin of the American Museum of Natural History 201: 1-160
- Komnenov M 2002 Contribution to the study of spiders (Araneae) on Šar Planina Mountain, north-western Macedonia. – Bulletin of the Biology Students' Research Society 2: 103-110
- Komnenov M 2003 Contribution to the study of spiders (Araneae) on Jakupica Mountain, Macedonia. – Bulletin of the Biology Students' Research Society 3: 45-49
- Komposch C 2009 Spinnentiere: Spinnen, Weberknechte, Pseudoskorpione, Skorpione (Arachnida: Araneae, Opiliones, Pseudoscorpiones, Scorpiones). In: Essl F & Rabitsch W (eds.) Neobiota in Österreich. Umweltbundesamt/Federal Environment Agency, Wien. pp. 250-262
- Kůrka A 1981 Spiders of the Baba State Natural reserve in Prague. – Časopis Národního Muzea, Řada přírodrovědná 150: 70-88
- Lazarov S 2004 A contribution to the study of spiders (Araneae) in Macedonia. – Acta Zoologica Bulgarija 56: 155-166
- Lazarov S 2007 Haplogyne spiders (Araneae) in Bulgaria: faunistic and zoogeographical analysis. In: Fet V & Popov A (eds.) Biogeography and ecology of Bulgaria. Series: Monographiae Biologicae. Springer, Dordrecht/Netherlands. pp. 481-492 – doi: [10.1007/978-1-4020-5781-6_16](https://doi.org/10.1007/978-1-4020-5781-6_16)
- Mikhailov KG 1997 Catalogue of the spiders of territories of the former Soviet Union (Arachnida, Aranei). Zoological Museum of the Moscow State University, Moscow. 416 pp.
- Misioch M 1977 Zur Spinnenfauna des Kaiserstuhl (Arach. Araneae). – Abhandlungen und Verhandlungen des Naturwissenschaftlichen Vereins in Hamburg 20: 133-149
- Nentwig W, Blick T, Gloor D, Hänggi A & Kropf C 2017 araneae – Spiders of Europe, version 4.2017. – Internet: <http://www.araneae.unibe.ch> (April 12, 2017)
- Paquin P & Dupérré N 2006 The spiders of Quebec: update, additions and corrections. – Zootaxa 1133: 1-37
- Pekár S 2002 *Zodarion rubidum* Simon, 1914: railroad riders? – Newsletter of the British Arachnological Society 95: 11-12
- Pekár S, Král J, Malten A & Komposch C 2005 Comparison of natural histories and karyotypes of two closely related ant-eating spiders, *Zodarion hamatum* and *Z. italicum* (Araneae, Zodariidae). – Journal of Natural History 39: 1583-1596 – doi: [10.1080/00222930400016762](https://doi.org/10.1080/00222930400016762)
- Renner F 1992 Liste der Spinnen Baden-Württembergs (Araneae). Teil 2. Liste der Spinnen Baden-Württembergs excl. Linyphiidae, Nesticidae, Theridiidae, Anapidae und Mysmenidae. – Arachnologische Mitteilungen 4: 21-55 – doi: [10.5431/aramit0402](https://doi.org/10.5431/aramit0402)
- Řezáč M 2002 *Sitticus distinguendus* and *Zodarion italicum*, two spider species recently found in Bohemia, Czech Republic (Araneae: Salticidae, Zodariidae). – Acta Universitatis Carolinae Biologica 45: 295-298
- Scharff N, Schmidt JB & Pedersen J 2007 Edderkoppen *Zodarion rubidum* Simon, 1914 - ny art og familie for Danmark. – Entomologiske Meddelelser 75: 65-70
- Simon E 1914 Les arachnides de France. Synopsis générale et catalogue des espèces françaises de l'ordre des Araneae. Tome VI. 1re partie. Roret, Paris. pp. 1-308
- Tzanev G & Lazarov S 2001 A contribution to the study of spiders (Araneae) in Osogovo Mountain, south-west Bulgaria. – Acta Zoologica Bulgarija 53: 67-78
- Vogel BR 1968 A zodariid spider from Pennsylvania (Araneida: Zodariidae). – Journal of the New York Entomological Society 76: 96-100
- World Spider Catalog 2017 World spider catalog, version 18.0. Natural History Museum Bern. – Internet: <http://wsc.nmbe.ch> (April 12, 2017)
- Wunderlich J 1973 Beschreibung einiger bisher unbekannter Arten der Gattung *Zodarion* Walckenaer aus Südeuropa (Arachnida: Araneae: Zodariidae). – Senckenbergiana Biologica 54: 171-176

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Arachnologische Mitteilungen](#)

Jahr/Year: 2017

Band/Volume: [54](#)

Autor(en)/Author(s): Krejci Tomas, Rezac Milan, Kadlec Tomas

Artikel/Article: [Zodarion ohridense \(Araneae: Zodariidae\) – a new record for Central Europe 5-7](#)