

## *Prinerigone vagans* new to Poland (Araneae: Linyphiidae), with comments on taxonomy and distribution

Jürgen Guttenberger, Luis Guttenberger & Tobias Bauer



doi: 10.30963/aramit5513

**Abstract.** The first record of the hygrophilous linyphiid spider *Prinerigone vagans* (Audouin, 1826) for Poland, collected on a bank of the River Olza near the Czech Border, is presented together with a discussion of known habitats of the species. A review of the taxonomic literature revealed that some characters of the male palp are highly variable within the distribution and even single populations of *P. vagans*. Future investigations should clarify the status of populations in Europe, the Near and Middle East as well as North Africa including Macaronesia.

**Keywords:** Erigoninae, new record, pedipalpus, spider

**Zusammenfassung.** *Prinerigone vagans* neu für Polen (Araneae: Linyphiidae), mit Kommentaren zur Taxonomie und Verbreitung. Der Erstnachweis der hygrophilen Linyphiide *Prinerigone vagans* (Audouin, 1826) für Polen, gesammelt am Ufer des Flusses Olza nahe der tschechischen Grenze, wird vorgestellt und zusammen mit den bekannten Habitaten diskutiert. Eine Durchsicht der taxonomischen Literatur zeigte auf, dass verschiedene Merkmale des männlichen Pedipalpus innerhalb des Verbreitungsgebiets und selbst einzelner Populationen sehr variabel sind. Zukünftige Untersuchungen sollten daher den Status der Populationen in Europa, dem Nahen und Mittleren Osten sowie Nordafrika inklusive Makaronesien klären.

Reporting the newly revealed presence of a species in a given country is one of the first steps towards overcoming the “Wallacean Shortfall” (Lomolino 2004) and represents important information for local biodiversity conservationists and nature conservation authorities. However, the distribution of many European spider species is still insufficiently known, especially in parts of Eastern and Southern Europe. While the biodiversity of Poland may be the highest in Central Europe (Convention on Biological Diversity 2017), currently only about 845 spider species are listed for the country (Nentwig et al. 2017). However, new country records of widespread European species are frequently published (e.g. Kronestedt 2006, Hajdamowicz 2009, Hajdamowicz et al. 2014, Rozwałka & Stachowicz 2015, Wiśniewski & Wesółowska 2015, Rozwałka et al. 2016, Wiśniewski & Dawidowicz 2017), indicating a large knowledge gap concerning the distribution of sometimes even relatively widespread species. Over 300 spider species currently known from Poland are Linyphiidae, which is comparable to the number in adjacent countries like Germany, the Czech Republic or Ukraine (Nentwig et al. 2017). *Prinerigone vagans* (Audouin, 1826), a linyphiid repeatedly found in Germany (Arachnologische Gesellschaft 2017) and distributed in the area of Berlin (Kielhorn 2010, 2016), was never collected in Poland before.

### Material and methods

The specimen was collected by hand and preserved in 75% ethanol. Photographs were made with a NikonD300 attached to a Novex RZ stereomicroscope. The map was created using the mapping system of the Arachnologische Gesellschaft (Arachnologische Gesellschaft 2017) and the records presented on the recording scheme of the Czech Arachnological Society and the cited literature (Czech Arachnological Society 2017). The drawing was made by TB, the material is deposited at the private collection of JG.

### Results

*Prinerigone vagans* (Audouin, 1826) (Figs 1–3)

Material. POLAND, Silesia, The Silesian Beskids mountain range, Istebna, bank of the river Olza, 49.57397° N, 18.90317° E (WGS 84), 1♂, 552 m a.s.l., collected by hand, 16.viii.2015, leg. Luis Guttenberger.




**Fig. 1:** Live male of *Prinerigone vagans* (Audouin, 1826) from Poland, dorsal view



**Fig. 2:** Male of *Prinerigone vagans* (Audouin, 1826) from Poland, dorsal view (Scale line = 0.5 mm)

Jürgen GUTTENBERGER, Luis GUTTENBERGER, Kurfürstenstr. 4, 92283 Lauterhofen, Germany; E-mail: juergenguttenberger@web.de  
Tobias BAUER, Staatliches Museum für Naturkunde Karlsruhe, Erbprinzenstr. 13, 76133 Karlsruhe, Germany; E-mail: tobias.bauer@smnk.de



**Fig. 3:** *Prinerigone vagans* (Audouin, 1826) from Poland, left male pedipalp, retrolateral view (Scale line = 0.2 mm; PA = patellar apophysis)

**Habitat.** The male was found on a bank of the river Olza (Figs 4–5) near the village of Istebna on wet mud with small water-filled pits and without vegetation. The river is enclosed by shrubbery, tree lines and a street. The landscape is dominated by forested areas along the hillsides and agriculture in the valleys. Other linyphiids found at the bank of the river were *Agyneia rurestris* (C. L. Koch, 1836) and *Bathyphantes gracilis* (Blackwall, 1841).

#### Habitat and distribution

*Prinerigone vagans* prefers humid, open habitats (Entling et al. 2007) and is often found at ground level near water (e.g. our record, Helsdingen 1997, Manderbach & Framenau 2001, Armbruster 2003, Bosmans 2007). A very detailed review of the habitat affinities in Europe can be found in Knülle (1954: p. 101), who also mentioned a strong association of *P. vagans* with small and saturated or still water-filled ground pits (“[...] Solche kleinen Bodenauskolbungen von 3–5 cm Tiefe, oft noch mit Wasserresten gefüllt, sind die Vorzugshabitate der Art.”), which could also be found at the river Olza (Fig. 5). However, several other records in Central Europe were made on annual cropland with pitfall traps (Blick et al. 2000), possibly due to aeronautic activities. It is questionable whether this species is able to build larger populations in cropland dominated landscapes, since harvest, tillage and crop rotation induce fast changes in abiotic conditions, which lead to a very ephemeral distribution of suitable habitats for this species (e.g. areas with waterlogging; Kielhorn 2016) and often to domination by only a few agrobiont species (Blick et al. 2000, Samu & Szinetár 2002). In the Mediterranean, *P. vagans* was collected near ponds (Morano et al. 2012), but also in a variety of other habitats (Buchholz 2013). In the Maghreb, *P. vagans* was recorded in similar habitats with temporary or permanent water, e.g. on stones along a river bed, marshy areas or in an irrigated garden (Bosmans 2007).




**Fig. 4:** Locality of *Prinerigone vagans* (Audouin, 1826), Isdebna, Bank of river Olza, Poland

*Prinerigone vagans* has been frequently found in Great Britain, France, Germany and the Benelux (Le Peru 2007, Tutelaers 2012, Arachnologische Gesellschaft 2017, British Arachnological Society 2017), and records exist from nearly all European Mediterranean countries (Nentwig et al. 2017), but the species seems to be rare in the Czech Republic, from which only two localities are known (Czech Arachnological Society 2017). Our single male was found in a typical habitat, which supports the hypothesis that (at least along the river Olza) Polish populations of the species exist. The species is probably absent from Scandinavia (except Denmark, Vangsgård et al. 1997) and other northern parts of Eastern Europe (Nentwig et al. 2017). It seems to be widely distributed in North Africa (Audouin 1826, Jocqué 1981, Bosmans 2007) and the Near East through to Iran (Pickard-Cambridge, 1872, Tanasevitch 2009). Interestingly, *P. vagans* is considered as the most common linyphiid in the Maghreb by Bosmans (2007). Other records were made, e.g., in Chinese parts of Central Asia (Zhou et al. 1983) and Marion Island in the southern Indian Ocean (Lawrence 1971).

Based on the known distribution in Germany, the Czech Republic and Poland (Fig. 5) it is possible that *P. vagans* is sensitive to continental climates with low winter temperatures, as already pointed out by Knülle (1954), and that the species already benefits or will benefit from climate change in Central Europe. However, this remains speculative since wide parts of eastern parts of Europe can still be seen as arachnological “Terra incognita”.

#### Taxonomic notes

*Prinerigone vagans* was described by Audouin (1826, sub. *Erigone*) from Egypt, Northern Africa, based on a male. Denis (1948) noted that the drawings of the male pedipalp in Audouin (1826) fit relatively well to an Algerian specimen, especially in the length of the tibia and patella, but not to his French specimens, which all had a longer and more slender patella and tibia and a differing patellar apophysis. He used a younger synonym, *Erigone spinosa* O. Pickard-Cambridge, 1872, to name this variation and delimit it from the variation described by Audouin (1826). Unfortunately, *E. spinosa* was originally used for specimens collected from a variety of localities in different countries (Egypt, Palestine, Italy) and it is not clear on which specimen the original drawing by O. Pickard-



**Fig. 5:** Records of *Prinerigone vagans* (Audouin, 1826) in Poland (red), Germany (black) and the Czech Republic (blue)

Cambridge (1872) was based. Denis (1948), not satisfied with the quality of the drawings in the first description, referred to the drawings of *E. spinosa* in O. Pickard-Cambridge (1910), which on the other hand were based on English specimens and correspond well to his French males. Our male from Poland possesses a long and slender patella and tibia and the typical patellar apophysis described by O. Pickard-Cambridge for *E. spinosa* (O. Pickard-Cambridge 1872, 1910). However, the length of the palpal segments seems to vary considerably within populations in Europe, as Knülle (1954) and Locket & Millidge (1953) pointed out. This hypothesis is also supported by assemblages from Hautes-Pyrénées (France) by Denis, which contained specimens of both variants (Denis 1950). Knülle (1954) mentioned two specimens from Northern Germany which fit the original drawings by Audouin (1826), without having visible differences in the more difficult structures of the bulbus compared to numerous specimens of the other variant. Locket & Millidge (1953) explained the variations in the segment lengths with the presence of allometric growth in males of *P. vagans*. Bosmans (2007) illustrated a male from the Maghreb, which seems to present a more intermediate form between the two variations of Denis (1948). However, the accompanying illustration of a male *P. vagans* palp (sub. *P. vagans vagans*) in Jocqué (1981) based on a specimen from the central Sahara is partially similar to the *spinosa*-variety, possessing a long and slender patella and tibia, but also a shorter and more robust patellar apophysis, better fitting the drawing by Audouin (1826) and hardly explained by allometric growth (Locket & Millidge 1953). Already Jocqué (1981) pointed out that there are considerable morphological variations between isolated populations in the Sahara region, which can be considered as relicts of a once vast distribution in a more humid past. On the other hand, Tanasevitch (2009) demonstrated that even within single populations in Iran very noticeable variations of the teeth on the palpal tibia occur. Future investigations should therefore target the gene-

tic diversity of *P. vagans* throughout its distribution and clarify the situation and the relationships especially between the North African, Near/Middle East and European populations including *Prinerigone pigra* (Blackwall, 1862) from Madeira, which seems to be only separable by the length of the patella and tibia of the male palp and possesses no visible differences in the structure of the bulbus (Wunderlich 1995). If some populations will be revealed as unrecognized species, their names must be chosen with care, since the younger synonym *Erigone spinosa* O. Pickard-Cambridge, 1872 refers to specimens from Europe and the Near East. However, it seems possible that *E. vagans* is a single polymorphic species and includes *P. vagans arabica* and *P. pigra*, as already pointed out by Tanasevitch (2009).

#### Acknowledgements

We are grateful to Konrad Wiśniewski for information on the Polish spider fauna and a constructive review. We would also like to thank Karl-Hinrich Kielhorn and Michael Schäfer for help with literature, Florian Raub for proofreading an earlier draft of the manuscript, the members of [spinnen-forum.de](http://spinnen-forum.de) for fruitful arachnological discussions, Marzena Stańska for reviewing the manuscript and Theo Blick for valuable comments. JG would like to thank fam. Kaluza und fam. Urbaczka for their support and the pleasant accommodation during the collecting activities in Poland.

#### References

- Arachnologische Gesellschaft 2017 Atlas der Spinnentiere Europas. – Internet: <http://atlas.arages.de> (15.ix.2017)
- Armbruster J 2003 Untersuchungen zur Spinnenfauna (Araneae) an Mittel-Gebirgsbächen und zur Besiedlung neu entstandener Uferstrukturen. – Arachnologische Mitteilungen 25: 17–37 – doi: [10.5431/aramit2502](https://doi.org/10.5431/aramit2502)
- Audouin V 1826 Explication sommaire des planches d'araignées de l'Égypte et de la Syrie. In: "Description de l'Égypte, ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expédition de l'armée française, publié par les ordres de sa Majesté l'Empereur Napoléon le Grand." – Histoire Naturelle 1(4): 99–186

- Blick T, Pfiffner L & Luka H 2000 Epigäische Spinnen auf Äckern der Nordwest-Schweiz im mitteleuropäischen Vergleich (Arachnida: Araneae). – Mitteilungen der Deutschen Gesellschaft für allgemeine und angewandte Entomologie 12: 267-276
- Bosmans R 2007 Contribution to the knowledge of the Linyphiidae of the Maghreb. Part XII. Miscellaneous erigonine genera and additional records (Araneae: Linyphiidae: Erigoninae). – Bulletin & Annales de la Société Entomologique de Belgique 143: 117-163
- British Arachnological Society 2017 Spider and Harvestman Recording Scheme. British Arachnological Society. – Internet: <http://srs.britishtspiders.org.uk/portal.php/p/Summary/s/Prinerigone+vagans> (10.x.2017)
- Buchholz S 2013 Spider records from East Macedonia and Thrace (NE Greece). – Arachnologische Mitteilungen 45: 45-53 – doi: [10.5431/aramit4510](https://doi.org/10.5431/aramit4510)
- Convention on Biological Diversity 2017 Poland – country profile. – Internet: <https://www.cbd.int/countries/profile/default.shtml?country=pl#facts> (20.x.2017)
- Czech Arachnological Society 2017 Distribution maps of arachnids in Czechia. – Internet: <http://www.arachnology.cz> (10.ix.2017)
- Denis J 1948 A new fact about *Erigone vagans* Aud. and Sav. – Proceedings of the Zoological Society of London 118: 588-590 – doi: [10.1111/j.1096-3642.1948.tb00400.x](https://doi.org/10.1111/j.1096-3642.1948.tb00400.x)
- Denis J 1950 Araignées de la région d'Orédon (Hautes-Pyrénées). – Bulletin de la Société d'Histoire Naturelle de Toulouse 85: 77-113
- Entling W, Schmidt MH, Bacher S, Brandl R & Nentwig W 2007 Niche properties of Central European spiders: shading, moisture and the evolution of the habitat niche. – Global Ecology and Biogeography 16: 440-448 – doi: [10.1111/j.1466-8238.2006.00305.x](https://doi.org/10.1111/j.1466-8238.2006.00305.x)
- Hajdamowicz I 2009 *Tetragnatha shoshone* Levi, 1981, a new spider species of longjawed orbweavers (Araneae, Tetragnathidae) in Poland. – Polish Journal of Entomology 78: 169-175
- Hajdamowicz I, Stańska M & Rutkowski T 2014 *Walckenaeria incisa* (O. P.-Cambridge) – a rare European species, new to Poland (Araneae: Linyphiidae). – Genus 25: 357-363
- Helsdingen PJ van 1997 Floodplain spider communities. – Proceedings of the 16th European Colloquium of Arachnology: 113-126
- Jocqué R 1981 Arachnids of Saudi Arabia: Fam. Linyphiidae. – Fauna of Saudi Arabia 3: 111-113
- Kielhorn K-H 2010 Neu- und Wiederfunde von Webspinnen (Araneae) in Berlin und Brandenburg, Teil 3. – Märkische Entomologische Nachrichten 12(1): 133-142
- Kielhorn K-H 2016 Beitrag zur Kenntnis der Webspinnen und Weberknechte in Berlin und Brandenburg. – Märkische Entomologische Nachrichten 17(2): 261-286
- Knülle W 1954 Zur Taxonomie und Ökologie der norddeutschen Arten der Spinnen-Gattung *Erigone* Aud. – Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 83: 63-110
- Kronstedt T 2006 On *Pardosa schenkeli* (Araneae, Lycosidae) and its presence in Germany and Poland. – Arachnologische Mitteilungen 32: 31-37 – doi: [10.5431/aramit3206](https://doi.org/10.5431/aramit3206)
- Lawrence R F 1971 Araneida. In: Zinderen Bakker EM van, Winterbottom JM & Dyer RA (eds.) Marion and Prince Edward Islands. Cape Town. pp. 301-313
- Le Peru B 2007 Catalogue et répartition des araignées de France. – Revue Arachnologique 16: 1-473
- Lockett GH & Millidge AF 1953 British spiders. Vol. II. Ray Society, London. 449 pp.
- Lomolino MV 2004 Conservation Biogeography. In: Lomolino MV & Heaney LR (eds.) Frontiers of biogeography: new directions in the geography of nature. Sinauer Associates, Sunderland/Massachusetts. pp. 293-296
- Manderbach R & Framenau VR 2001 Spider (Arachnida: Araneae) communities of riparian gravel banks in the northern parts of the European Alps. – Bulletin of the British Arachnological Society 12: 1-9
- Morano E, Pérez-Bilbao A, Benetti CJ & Garrido J 2012 Arañas (Arachnida: Araneae) en lagunas de la Red Natura 2000 de Galicia (Noroeste de España). – Revista Ibérica de Aracnología 20: 71-83
- Nentwig W, Blick T, Gloor D, Hänggi A & Kropf C 2017 Spinnen Europas. Version 10.2017. – Internet: <http://www.araneae.unibe.ch> – doi: [10.24436/1](https://doi.org/10.24436/1) (10.x.2017)
- Pickard-Cambridge O 1872 General list of the spiders of Palestine and Syria, with descriptions of numerous new species, and characters of two new genera. – Proceedings of the Zoological Society of London 40(1): 212-354, Pl. XIII-XVI
- Pickard-Cambridge O 1910 On British Arachnida noted and observed in 1908. – Proceedings of the Dorset Natural History and Antiquarian Field Club 30(for 1909): 97-115
- Rozwałka R & Stachowicz J 2015 The first record of *Clubiona saxatilis* L. Koch 1867, (Araneae: Clubionidae) in Poland. – Fragmenta faunistica 58: 59-63 – doi: [10.3161/00159301FF2015.58.1.059](https://doi.org/10.3161/00159301FF2015.58.1.059)
- Rozwałka R, Rutkowski T, Sienkiewicz P & Zawal A 2016 *Zelotes erebeus* (Thorell, 1871) (Araneae: Gnaphosidae) in Poland and its distribution in Europe. – Entomologica Fennica 27: 1-7
- Samu F & Szinetar C 2002 On the nature of agrobi-ont spiders. – Journal of Arachnology 30: 389-402 – doi: [10.1636/0161-8202\(2002\)030\[0389:otnoas\]2.0.co;2](https://doi.org/10.1636/0161-8202(2002)030[0389:otnoas]2.0.co;2)
- Tanasevitch AV 2009 The linyphiid spiders of Iran (Arachnida, Araneae, Linyphiidae). – Revue Suisse de Zoologie 116: 379-420 – doi: [10.5962/bhl.part.81325](https://doi.org/10.5962/bhl.part.81325)
- Tutelaers P 2012 Benelux spider distribution maps. – Internet: <http://www.knnv.nl/eindhoven/iwg/Araneae/SpiBenelux> (16.iii.2018)
- Vansgård C, Reinke H-D, Schultz W & Helsdingen PJ van 1997 Red List of spiders (Araneae) of the Wadden Sea Area. – Helgoländer Meeresuntersuchungen 50, Suppl.: 77-82
- Wiśniewski K & Dawidowicz A 2017 *Uloborus walckenaerius* and *Oxyopes heterophthalmus* in Poland (Araneae: Uloboridae, Oxyopidae). – Arachnologische Mitteilungen 54: 48-51 – doi: [10.5431/aramit5411](https://doi.org/10.5431/aramit5411)
- Wiśniewski K & Wesolowska W 2015 *Maro leptineni* (Araneae: Linyphiidae) – a spider species new to the fauna of Poland. – Arachnologische Mitteilungen 50: 81-84 – doi: [10.5431/aramit5011](https://doi.org/10.5431/aramit5011)
- Wunderlich J 1995 Zu Ökologie, Biogeographie, Evolution und Taxonomie einiger Spinnen der Makaronesischen Inseln (Arachnida: Araneae). – Beiträge zur Araneologie 4: 385-439
- Zhou, NL, Wang H & Zhu CD 1983 [New records of spiders from Uygur Autonomous Region and Heilongjiang Province, China]. – Journal of the Bethune Medical University 9 (suppl.): 153-160 [in Chinese]

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Arachnologische Mitteilungen](#)

Jahr/Year: 2018

Band/Volume: [55](#)

Autor(en)/Author(s): Guttenberger Jürgen, Guttenberger Luis, Bauer Tobias

Artikel/Article: [Prinerigone vagans new to Poland \(Araneae: Linyphiidae\), with comments on taxonomy and distribution 75-78](#)