

Beitrag zur Insektenfauna von Zanzibar.

III. Coleoptera.

(Fortsetzung von Jahrg. XXXIII. p. 49.)

Von

A. Gerstaecker.

Fam. Carabidae.

92. (1 a). *Cicindela neglecta* Dejean, Spec. gén. I. p. 114, Nr. 96. — Von der Insel Sansibar.

93. (1 b). *Cicindela congrua* * Klug, Insekt. v. Mosamb. p. 148. Taf. IX, fig. 2. — Von Mombas.

94. (11 a). *Pheropsophus marginatus* Dej., Spec. gén. I. p. 309, No. 16. — Von der Insel Sansibar.

95. (22 a). *Tefflus juvenilis*, n. sp. *Minor, niger, setosus, prothoracis vix hexagoni angulis lateralibus obtuse rotundatis, marginibus deplanatis: coleopteris oblongo-ovatis, octocostatis, interstitiis angustis, transverse crenulatis.* Long. 33, lat. elytr. 12 mill. ♂. — Aus der Tula Colonie, 196 s. Br.

96. (24 a). *Chlaenius posticus* Fab., Entom. syst. suppl. I. p. 75, No. 102—103. — Von der Insel Sansibar.

97. (24 b). *Chlaenius eineticollis* * Bohem., Insect. Caffrar. I. p. 148, Nr. 156. — Von der Insel Sansibar.

98. (27 a). *Hypolithus holosericeus* Dej., Spec. gén. IV. p. 171, Nr. 3. — Von der Insel Sansibar.

99. (29 a). *Hypolithus venustulus* Bohem.,
Insect. Caffrar. I. p. 202, No. 211. — Von der Insel Sansibar.

Fam. Dyticidae.

100. (31 a). *Cybister tripunctatus* Oliv., Entom.
III, 40. p. 14. pl. 3, fig. 24 (= *Dyt. lateralis* Fab. = *Cyb.*
Africanus Lap.). Von der Insel Sansibar.

101. (31 b). *Cybister auritus*, n. sp. *Parvus,*
nigro-piceus, nitidus, supra aeneo-micans, antennis, labro,
clypei fascia anteriore, prothoracis margine laterali testaceis
vel ferrugineis, pedibus anterioribus elytrorumque macula
subapicali (interdum obsoleta) rufis: clytris perspicue trise-
riatim punctatis, tibiarum posticarum calcare externo bre-
viusculo, acute lunceolato. Long. $15\frac{1}{2}$ —17 mill. ♂♀.

Var. ♀. *Elytrorum disco obsolete et minus confertim*
pustuloso.

Auf der Insel Sansibar und auf Madagascar einheimisch.

102. (31 c). *Cybister immarginatus* Fab., En-
tom. syst. IV. p. 444, No. 4—5 (= *Cyb. bimaculatus* Aubé
= *Cyb. binotatus* Bohem. = *Cyb. Owas* Lap. Aubé).
Auf der Insel Sansibar gefunden.

Fam. Palpicornia.

103. (38 a). *Cyclonotum abdominale* * Fab.
Entom. syst. I. p. 79, No. 10 (= *Sphaeridium depressum*
* Klug = *Cyclonotum nitidum* * Bohem.). Von Mombas
und der Insel Sansibar vorliegend.

Fam. Staphylinidae.

104. (44 a). *Paederus aestuans* * Erichson, Gen.
et spec. Staphyl. p. 655, No. 10. — Von der Insel Sansibar.

Fam. Palpatores.

105. (46 a). *Scydmaenus pinguiculus*, n. sp.
Palporum maxillarium articulo quarto obsoleto, mesosterno
carinato, tarsi anticis basin versus dilatatis: alatus, nigro-
piceus, griseo-pilosus, nitidus, antennis, palpis pedibusque
rufis, prothorace rotundato-quadrato, satis convexo, colco-

pteris ovatis, basi subtruncatis, obsolete punctatis. Long. $2\frac{3}{4}$ mill. — Bei Mbaramu aufgefunden.

106. (52 a). *Hister nigrita* * Erichs., Jahrb. d. Insektenk. p. 131, No. 7 (= *Hist. caffer* Bohem.) — Von der Insel Sansibar.

107. (52 b). *Hister striolatus* * Mars., Annal. soc. entom. de France 3. sér. II. p. 200, No. 21. pl. 6, fig. 21. — Von der Insel Sansibar.

Fam. Phalacridae.

108. *Phalacrus aethiops*, n. sp. *Breviter ovatus, convexus, niger, subaeneo-micans, nitidissimus, elytris obsolete sulcatis, disperse et subtilissime punctulatis, antennis piceis, clava oblonga, obtusiuscula.* Long. 2 mill. — Von Mombas.

Fam. Nitidulariae.

109. *Brachypeplus depressus* * Erichs. in Germar's Zeitschrift f. d. Entom. IV. p. 247, No. 3. — Bei Mbaramu in Menge gefunden.

110. *Carpophilus hemipterus* Lin. Syst. nat. I. p. 565, No. 30. — Von Mbaramu.

111. *Carpophilus fumatus* Bohem. Insect. Caffr. I. p. 564, Nr. 618. — Häufig auf den Ugono-Bergen gefunden.

112. *Carpophilus obsoletus* * Erichs. in Germar's Zeitschrift f. d. Entom. IV. p. 259, Nr. 13. — Von Mombas.

113. *Aethina combusta*, n. sp. *Obtuse ovata, subparallela, confertim punctulata, ferruginea, prothorace, scutello elytrorumque basi flavescenti-sericeis, his ceterum brunneis et latera versus infuscatis, setulis nigris dense obsitis.* Long. 4—6 mill. — Von Mombas und der Insel Sansibar.

114. *Meligethes heteropus*, n. sp. *Niger, subaeneo-micans, confertim punctulatus, sericeus, antennis pedibusque anticis rufo-brunneis, tarsorum articulis 1.—4. testaceis: tibiis anticis latis, margine externo serrato, ante apicem*

bidentato, unguiculis simplicibus. Long. $1\frac{3}{4}$ mill. — Von Kisuani.

115. *Lordites claudus*, n. sp. *Nigro-piceus, confertim punctatus, pubescens, ore, antennarum basi, prothoracis elytrorumque lateribus, coxis nec non femorum dimidio apicali rufis, elytrorum interstitiis 3., 5., 7. costatim elevatis, seriatim sanguineo-maculatis: ♂ tibiis anterioribus apice fortiter incurvis, posticis arcuatis, apice dilatatis.* Long. $4\frac{1}{2}$ (♀) — 6 (♂) mill. — Von den Ugono-Bergen.

116. *Trogosita mauritanica* Lin. Syst. nat. I. p. 674, No. 4. — Von der Insel Sansibar.

Fam. Colydi.

117. *Cerylon pygmaeum*, n. sp. *Deplanatum, rufo-ferrugineum, nitidum, prothorace subquadrato, apicem versus angustato, supra sat confertim punctato, elytris punctato-striatis, striis apicem versus obsoletioribus.* Long. $1\frac{3}{4}$ mill. — Von den Ugono-Bergen (und Madagascar).

Fam. Cucujini.

Laemotmetus, nov. gen. (Trib. Passandrii). *Tarsorum articulus primus brevissimus. Processus jugulares late truncati. Labrum obtectum. Mandibulae obtusae. Caput declive, tumidum, haud sulcatum. Oculi fere obtecti, rotundato-triangulares. Antennae breviusculae, moniliformes.*

118. *Laemotmetus ferrugineus*, n. sp. *Elongatus, parallelus, rufo-ferrugineus, nitidus, glaber, capite prothoraceque sat confertim punctatis, elytris dilutioribus.* Long. $3\frac{2}{3}$ mill. — Von Mombas.

Fam. Dermestini.

119. *Dermestes vulpinus* Fab. Spec. Insect. I. p. 64, No. 9. — Häufig auf dem Festlande und auf der Insel Sansibar.

120. *Dermestes cadaverinus* Fab. Syst. Entom. p. 55, Nr. 3. — Von Mombas und der Insel Sansibar.

121. *Trogoderma haemorrhoea*, n. sp. *Antennarum clava oblonga, triarticulata, corpore breviter ovato, pubescente, nigro, antennarum basi, pedibus intermediis ely-*

trorumque parte postica rufo-brunneis. Long. 1 $\frac{3}{4}$ mill. — Von Endara.

Hypocenthes, nov. gen. *Ab Anthreno differt corpore glabro, confertissime et profunde punctato, prothoracis lobo scutellari retrorsum magis producto et acutiore, oculis minoribus.*

122. *Hypocenthes aterrimus*, n. sp. *Breviter ovatus, fortiter punctatus, glaber, niger, subnitidus, prothoracis margine basali, elytrorum apice et margine laterali obscure rufo-brunneis. Long. 2 $\frac{1}{3}$ mill. — Von Kisuani.*

Fam. Lamellicornia.

123. *Figulus sublaevis* Palis. Insect. rec. en Afrique p. 3, pl. 1, fig. 3 (= *Fig. anthracinus* * Klug = *Fig. ebenus* Westw.). — Von der Insel Sansibar.

124. (64 a). *Pachnoda euparypha*, n. sp. *Ob longo-ovata, subparallela, viridis, infra nitida, albo-maculata, supra opaca, tomentosa, fronte, pronoti marginibus, elytrorum limbo lato pygidioque croceis, hoc albo-viridique signato. Long. 18 $\frac{1}{2}$ —21 mill. ♂♀.*

Var. *Elytrorum disco toto lacte rufo, vitta exteriori tantum viridi.* — Von der Insel Sansibar (und aus Mosambik).

125. (66 a). *Diplognatha hebraea* Oliv., Entom. I, 6. p. 89, No. 116. pl. 12, fig. 115. — Von der Insel Sansibar.

126. (66 b). *Cymophorus undatus* Kirby, Zool. Journ. III. p. 271, pl. 5, fig. 6. — Vom Küstenstrich des Festlandes.

127. (71 a). *Rhizoplatys ambiguus*, n. sp. *Mandibulis angustis, tuberculo frontali pyramidalis, clypeo acuminato, mento apicem versus angustato et profunde excavato: subdepressus, niger, nitidulus, prothorace fortiter marginato, basin versus canaliculato, inter puncta majora confertim punctulato, elytris subcostatis, subtiliter cicatricoso-punctatis. Long. 26 mill. ♀ — Von der Insel Sansibar.*

128. (71 b). *Cyphonistes vallatus* Wiedem., Zoolog. Magaz. II, 1. p. 4, No. 1. — Von der Insel Sansibar.

129. (70 a). *Oryctes monoceros* Oliv. Entom. I, 3. p. 37, No. 40. pl. 13, fig. 123. — Von der Insel Sansibar.

130. (70 b). *Heteronychus atratus* * Klug, Monatsber. Akad. d. Wiss. Berlin 1855. p. 657. Vom Küstenstrich des Festlandes.

131. (74 a). *Leucopholis lepidota* * Klug, Insekt. v. Mosamb. p. 256. Taf. 15. fig. 5. — Vom Küstenstrich des Festlandes und von der Insel Sansibar.

Empycastes, nov. gen. (Trib. Melolonthidae).
A Psilonycho Burm. *differt antennarum clava graciliore, clypei excavati margine apicali profunde trisinuato ideoque quadridentato, prothoracis angulis anticis auriculato-recurvis, ejusdem margine apicali incrassato, scutello cordiformi, tibiis anticis extus fortiter tridentatis, intermediis bispinosis, tarsis anterioribus validis, elongatis, pedibus posticis breviusculis, robustis, tarsis dilatatis, depressis.*

132. (74 b). *Empycastes coronatus*, n. sp. *Rufopiceus, parum nitidus, supra fere nudus, infra dense fulvo-villosus, antennis ferrugineis, prothorace rufescente, confertim cicatricoso-punctato callisque tribus basalibus instructo, elytris parallelis, rugoso-punctatis, parce albo-squamulosis.* Long. $17\frac{1}{2}$ — $18\frac{1}{2}$ mill. ♂. — Von der Insel Sansibar.

133. (75 a). *Schizonycha tumida* (* Illig.) Casteln. Hist. nat. d. Coléopt. II. p. 133, No. 7. — Von der Insel Sansibar.

134. (80 a). *Hoplia retusa* * Klug, Insekt. von Madag. p. 82, No. 109. Taf. 3, fig. 9. — Vom Küstenstrich des Festlandes.

135. *Aphodius marginicollis* (Dej.) * Klug, Insekt. v. Mosamb. p. 244. — Von der Insel Sansibar.

136. *Aphodius moestus* Fab. Syst. Eleuth. I. p. 78, No. 45 (= *Aphod. mutans* Walk.). — Von der Insel Sansibar.

137. *Aphodius anthrax*, n. sp. *Niger, glaber, nitidus, capitis prothoracisque marginibus, tibiis tarsisque rufopiceis, illo irregulariter disperse punctato, elytris profunde punctato-sulcatis, interstitiis apicem versus costatim elevatis.* Long. $3\frac{1}{2}$ mill. — Von Uru.

138. *Ateuchus prodigiosus* * Erichs., Archiv

f. Naturgesch. IX, 1. p. 231, No. 51. — Vom Küstenstreich des Festlandes und von der Insel Sansibar.

139. *Ateuchus Aegyptiorum* Latr. in Caillaud, Voyage à Méroé II. Ins. p. 9.

Var. *purpurascens*. *Supra purpureo-cupreus, infra cum pedibus aterrimus, rarius pro parte viridi-micans*. — Von Mombas.

140. *Ateuchus aeratus*, n. sp. *Fronte tuberculata, elytrorum margine laterali bicarinato, tibiaram anticarum dente superiore fere obsoleto: obscure aeneus, subnitidus, pronoti confertim granoso-punctati linea media areisque duabus antebasalibus laevigatis, elytrorum interstitiis cicatricoso-punctatis, disperse setulosis, sutura elevata, nitida*. Long. 23—29 mill. ♂♀.

Fem. *Tibiaram posticarum scopa nigra, abbreviata*.

Var. *minor*. *Pronoti areis antebasalibus obsoletis*. Long. 20 mill. — Von Mombas und Moschi.

141. *Ateuchus pustulosus*, n. sp. *Fronte vix tuberculata, elytrorum margine laterali bicarinato, tibiaram anticarum dente superiore fere obsoleto: parum convexus, niger, subnitidus, pronoti discum versus laxius punctati linea media apice abbreviata laevi, elytrorum interstitiis sat fortiter disperse punctatis, punctis opacis, setigeris*. Long. 22—32 mill. ♂♀.

Fem. *Tibiaram posticarum scopa fusca, elongata*. — Von Moschi.

142. *Ateuchus (Actinophorus) catenatus*, n. sp. *Fronte haud tuberculata, elytrorum margine laterali dilatato, acutissimo, tibiis anticis acute quadridentatis, mesosterno lanceolato, apice recurvo, femoribus anticis fortiter dilatatis, basi bidentatis, posticis basi excisis: niger, subopacus, pronoto confertim punctato, foveis majoribus setigeris disperse obsito, elytris subsulcatis, interstitiis interrupte unicastis*. Long. 27—29 mill. — Von Aruscha und Endara.

143. *Ateuchus (Pachylomera) femoralis* Kirby, Zool. Journ. III. p. 520, Taf. 14, fig. 1 (= *Ateuch. horridus* * Bohem.). — Von Dschagga.

144. *Gymnopleurus splendidus* Bertol. Illustr. rer. nat. Mozamb., Coleopt. I. p. 15, No. 10 (= *Gymn. pro-*

fanus Latr. = Gymn. Wahlbergi * Bohem.). — Von Endara.

145. *Gymnopleurus umbrinus*, n. sp. *Clypeo sexdentato, mesosterno obtuse tuberculato, fusco-aeneus, supra confertim granulatus ibique breviter fulvo-pilosus: pronoti basi profunde biimpressa, disco posteriore obsolete tricalloso, elytrorum interstitiis alternis maculatim denudatis*. Long. $10\frac{1}{2}$ — $11\frac{1}{2}$ mill. ♂♀.

Mas. *Tibiarum anticarum calcare obtuso, apice incurvo, dentibus tribus externis brevioribus: femoribus anticis minus dilatatis, muticis*.

Fem. *Tibiarum anticarum calcare acuminato, dentibus tribus externis longioribus: femoribus anticis fortiter dilatatis, unidentatis*. — Von Endara.

146. *Gymnopleurus virens* * Erichs. Archiv f. Naturgesch. IX, 1. p. 231, No. 52 (= Gymn. laevicollis Casteln. = Gymn. pumilus Reiche = Gymn. gibbosus Roth). — Von Mombas.

147. *Sisypheus nodifer*, n. sp. *Parvulus, capite prothoraceque acneo-micantibus, hoc basin versus canaliculato, elytris catenato-striatis, seriatim ferrugineo-setulosus, trochanteribus posticis appendice brevi, apice dilatato instructis*. Long. 5 mill. ♂. — Von Mbaramu.

148. *Sisypheus seminulum*, n. sp. *Minutus, niger, acneo-micans, fere opacus, elytrorum margine pedibusque sanguineis: capite prothoraceque disperse punctatis et pallide setulosus, elytris subtilissime aciculatis, catenato-striatis, interstitiis uniserialim punctatis*. Long. $3\frac{1}{2}$ —4 mill. ♀. — Von der Insel Sansibar.

149. *Anachaleos procerus*, n. sp. *Oblongus, cupreo-aeneus, subopacus, coleopteris latitudine perspicue longioribus, evidenter striatis, pygillii apice fortiter reflexi superficie ventrali laevi, lucida, tibiis anterioribus apicem versus acute tridentatis, posticis longissimis, arcuatis*. Long. 31 mill. — Von Moschi.

150. *Copris Neptis* Reiche, Voyage en Abyssin. III. p. 314, No. 2. pl. 19, fig. 3. — Von der Insel Sansibar.

151. *Onitis Inuus* Oliv. Entom. I, 3. p. 138,

No. 165. pl. 14, fig. 135 (= Onit. Alexis Dej. * Klug). — Von der Insel Sansibar.

152. *Onthophagus lanista* Casteln. Hist. nat. d. Coléopt. II. p. 84, No. 3 (= Onth. aulicus Dej. Bohem.). — Vom See Jipe.

153. *Onthophagus loricatus* * Klug, Monatsber. Akad. d. Wiss. Berlin 1855. p. 653 (= Onth. collaris Bohem.). — Vom See Jipe.

154. *Onthophagus catta* Fab. Mant. Insect. I. p. 12, No. 115 (= Copr. gazella et metallicus Fab. ♂). — Von Mombas und der Insel Sansibar.

155. *Onthophagus pinguis*, n. sp. *Obscure viridi-acneus, parum nitidus, clypeo transverse ruguloso nec non verticis cornu erecto subviolaceis: prothorace confertim granoso-punctato, ante apicem bicalloso, clytris gemellato-striatis, interstitiis internis parcius, externis confertim granulosis, perspicue pubescentibus.* Long. 12—13½ mill. ♂. — Von der Insel Sansibar.

156. *Onthophagus pugionatus* Bohem. Ins. Caffrar. II. p. 286, No. 969. — Von der Insel Sansibar.

157. *Onthophagus exasperatus*, n. sp. *Ovatus, fusco-acneus, subnitidus, setulosus, clypeo subtruncato, vertice cornu erecto, triangulari instructo, prothorace aequaliter scabroso-punctato, ante apicem leviter bituberculato, clytris gemellato-striatis, interstitiis disperse granoso-punctatis.* Long. 5 mill. ♂. — Von den Bura-Bergen.

158. *Onthophagus laceratus*, n. sp. *Clypeo amplo, biapiculato, niger, parum nitidus, parce flavo-setulosus, antennis, palpis femoribusque posterioribus rufis, pronoti macula laterali postica clytrisque aurantiacis, harum plaga magna discali, multifariam excisa atra.* Long. 7—9 mill. ♂♀.

Mas. *Clypeo acute bidentis, prothorace latiore, convexiore.*

Fem. *Clypeo sinuato, prothorace angustiore, subdepresso.* — Von der Insel Sansibar.

159. *Onthophagus hinnulus* * Klug, Insekt. v. Madag. p. 75, No. 93. — Von der Insel Sansibar.

160. *Onthophagus vinetus* * Erichs., Archiv f.

Naturgesch. IX, 1. p. 234, No. 58. — Von der Insel Sansibar.

161. *Onthophagus picticollis*, n. sp. *Frontis excavatae margine antico elevato, eum carina clypei angulari transversa connexo: oblongo-ovatus, glaber, nitidus, capite prothoraceque piceis, aeneo-micantibus, huius disco et lateribus flavo-maculatis, elytris prothorace subtilius punctatis, testaceis, fusco-pictis.* Long. 7 mill. — Von Uru.

162. *Onthophagus aterrimus*, n. sp. *Niger, glaber, nitidus, prothorace disperse punctato, elytris vix brevior, his subtilissime aciculatis, catenato-striatis, in interstitiis parce punctulatis.* Long. $8\frac{1}{2}$ mill. ♂♀.

Mas. *Clypeo breviusculo, subtruncato, frontis excavatae carina arcuata, acuta, prothorace gibboso, apicem versus utrinque retuso et tuberculato.*

Fem. *Clypeo longiore, transverse ruguloso, frontis carina transversa obtusiuscula, prothorace simplice, fortius punctato.* — Vom See Jipe und den Bura-Bergen.

163. *Onthophagus tumidulus*, n. sp. *Nigro-piceus, aeneo-micans, glaber, nitidus, clypeo rotundato, capite prothoraceque inermibus, hoc disperse punctato, elytris subsulcatis, sulcis subtilius, interstitiis fortius et parce punctatis.* Long. $8\frac{1}{2}$ — $9\frac{1}{2}$ mill. ♂♀.

Mas. *Clypeo breviusculo, subrugose punctato, frontis excavatae carina arcuata, acuta, prothorace apicem versus leviter gibboso et subtilius punctato.*

Fem. *Clypeo longiore, confertim transverse rugoso, frontis densius punctatae carina transversa, obtusior, prothorace aequaliter convexo.* — Vom See Jipe.

164. *Onthophagus ovulum*, n. sp. *Ovatus, nitidus, supra glaber, capite prothoraceque cupreo-micantibus, disperse et sat fortiter punctatis, illius vertice excavato carinaque transversa terminato: elytris nigro-aeneis, subsulcatis, in sulcis geminato-striatis et punctatim impressis, interstitiis fortiter biserialim punctatis.* Long. $3\frac{1}{4}$ mill. — Vom See Jipe.

165. *Onthophagus sculptilis*, n. sp. *Ovatus, aeneus, setulosus, nitidus, capite prothoraceque subcupreis, illius vertice bicorni, hoc rude punctato, ante apicem unitu-*

berculato, clytris punctatissimis, suturam versus costatis, costis laevigatis. Long. $5\frac{1}{2}$ mill. ♂. — Von Mbaramu.

166. *Onthophagus morosus*, n. sp. *Nigro-aeneus, opacus, confertim punctatus, breviter setulosus, clypeo subtruncato, fronte bicarinata, vertice bituberculato, prothorace apicem versus binodoso, utrinque leviter retuso, clytris catenato-striatis, interstitiis parce granoso-punctatis.* Long. $6\frac{1}{4}$ mill. ♂. — Vom See Jipe.

167. *Onthophagus lamelliger*, n. sp. *Ovatus, niger, subaeneo-micans, fere opacus, setulosus, clypeo breviusculo, vertice lamina erecta subquadrata, apice bifida instructo, prothorace fortiter punctato, antice profunde excavato, supra angulatim producto, clytrorum macula humerali margineque apicali rufo-ferrugineis, interstitiis biscriatim punctatis.* Long. $5\frac{1}{3}$ mill. ♂.

168. *Oniticellus nasieornis* Reiche in: Voyage en Abyssinie III. p. 339, No. 3. pl. 10, fig. 7 (= *Onit. pallens* Casteln.). — Von der Insel Sansibar.

169. *Oniticellus planatus* (* Illig.) Casteln., Hist. nat. d. Coléopt. II. p. 91, No. 3. — Vom See Jipe.

170. *Oniticellus humilis*, n. sp. *Oblongus, depressus, subtus testaceus, supra fuscus, opacus, ferrugineo-varius, setulosus, capite prothoraceque confertim cicatricoso-punctatis, illius clypeo simplice, aeneo-micante, hoc basin versus sulcato: clytris subtilissime aciculatis, rufo limbatis, apice setulis bacilliformibus obsitis.* Long. $5\frac{1}{3}$ — $6\frac{1}{3}$ mill. — Von Mbaramu.

171. *Oniticellus caelatus*, n. sp. *Capite utrinque auriculato, clypeo amplo, apice bilobo, fronte basin versus acute carinata: niger, opacus, setulosus, prothorace confertim ocellato-punctato, basin versus profunde, in disco levius complurics impresso, clytris obsolete quadricostatis.* Long. 5 mill. — Von Aruscha.

172. *Drepanocerus laticollis* Bohem. Insect. Caffrar. II. p. 325, No. 1009. — Von der Insel Sansibar.

Fam. Buprestidae.

173. *Steraspis fastuosa*, n. sp. *Infra cum pedibus cyanca, maculatim albido-tomentosa, supra aurco-viri-*

dis, elytrorum margine laterali purpurascenti-cupreo: pronoti profunde sulcati foveis lateralibus ante basin abbreviatis. Long. 45 mill. ♀. — Von Kiriama.

174. *Sphenoptera neglecta* * Klug in Erman's Reise p. 30, No. 41. — Von Endara.

175. *Sphenoptera quadraticollis*, n. sp. *Elytris apice trispinosis, prothorace quadrato, utrinque ante basin subsinuato, supra disperse punctato: angusta, aenea, nitida, elytris prothorace obscurioribus.* Long. 8½ mill. — Von Mombas.

176. *Belionota canaliculata* Fab. Mant. Insect. I. p. 181, No. 58. — Von der Insel Sansibar.

177. *Chrysobothris serrata* Fab. Entom. syst. I, 2. p. 200, No. 61. — Von Mombas.

178. *Chrysobothris empyrea*, n. sp. *Supra viridi-cyanea, disperse punctata, nitida, infra cum pedibus cupreo-metallica, utrinque maculatim albo-pilosa: capite aureo-viridi, vertice prothoracisque trapezoidei lateribus purpurascenti-cupreis: femoribus anticis incrassatis, unidentatis, segmento ventrali ultimo apice exciso, fortiter carinato.* Long. 7 mill. ♂. — Von der Insel Sansibar.

179. *Pseudagrilus zonatus* Roth, Archiv f. Naturgesch. XVII, 1. p. 119. No. 27. — Von der Insel Sansibar.

180. *Aphanisticus nodosus*, n. sp. *Infra obscure aeneus, supra cupreus, elytris apice excepto nigro-cyaneis: capite bicornuto, prothoracis disco fortiter quadri-nodoso, basi utrinque profunde foveolata, elytris interrupte costulatis, apice unidentatis.* Long. 4⅓ mill. — Von Mombas.

Fam. Elateridae.

181. *Agrypnus maculicollis*, n. sp. *Elytris apice integris, fuscus vel rufo-brunneus, flavescenti-pubescens, prothorace discum versus laxius punctato, linea media plagisque duabus anterioribus laevibus, his pube densiore circumdatis.* Long. 22—25 mill. — Von der Insel Sansibar.

182. *Alaus atropos*, n. sp. *Scutello plano, oblongo-quadrato, elytrorum angulo suturali submucronato:*

infra niger, supra rufo-brunneus, flavescenti-tomentosus, prothoracis disco denudato: latus maculis duabus ovatis elytrorumque fasciis duabus, altera post medium, altera ante apicem sita, utro-tomentosis. Long. 18 mill. ♀. — Vom See Jipe.

183. *Lacon occidentalis*, Caud., Monogr. d. Elatér. I. p. 130, No. 48. — Vom See Jipe.

184. *Ischiodontus pedestris*, n. sp. *Supra nigro-fuscus, infra cum antennis pedibusque rufo-castaneus, cervino-pubescent, prothorace apicem versus sensim angustato, punctis umbilicatis laxè obsito, sat nitido, angulis posticis acute carinatis, rufo-brunneis. Long. 11 mill. — Von der Insel Sansibar.*

185. *Monoerepidius advena*, n. sp. *Elytris apice rotundatis, tarsorum articulo quarto anguste lamellato: breviusculus, nigro-piceus, nitidus, subtiliter cervino-pubescent, antennis pedibusque ferrugineis, prothoracis basi fortiter declivi, rufescente. Long. 4 mill. — Von Aruscha.*

186. *Heteroderes acutangulus*, n. sp. *Tarsorum articulo quarto anguste lamellato, prothorace basi vix tuberculato, longiore quam latiore, angulis posticis acutissimis, carinatis: fuscus, flavescenti-sericeus, antennis pedibusque testaceis. Long. 8—8½ mill. — Von Mbaramu.*

187. *Melanoxanthus melanocephalus* Fab. Spec. Insect. I. p. 272, No. 38. — Von der Insel Sansibar.

188. *Cardiophorus hoploderus* Cand., Monogr. d. Elatér. III. p. 175, No. 83. — Von Mombas.

189. *Cardiophorus obsoletus*, n. sp. *Unguiculis simplicibus, piceus, cinereo-pubescent, antennis, pedibus elytrisque rufo-brunneis, his suturam versus infuscat: prothorace oblongo, parce et obsolete punctato, in medio bascos unituberculato. Long. 8 mill. — Von Endara.*

190. *Melanotus umbilicatus* Schönh., Synon. Insect. III. Append. p. 137, No. 188 (= *Cratonychus Africanus* Bohem.). — Von der Insel Sansibar.

191. *Ludius penicillatus*, n. sp. *Fuscus, umbrino-pubescent, antennis, pedibus, scutello, elytris abdomine-*

que rufo-brunneis: prothoracis oblongi angulis posticis acute carinatis, apice penicillatis. Long. 15 mill. — Von Mbaramu.

Fam. Malacoderma.

192. *Lampyris amplicolis*, n. sp. *Prothorace amplo, reflexo-marginato, basin versus acute carinato, ante apicem utrinque pellucido: aurantiaca, antennis, prothoracis macula didyma elytrorumque disco nigricantibus, tibiis tarsisque piceis.* Long. 10—11, lat. 4 mill. ♂. — Von der Insel Sansibar.

193. *Lampyris vidua*, n. sp. *Aptera, angusta, subparallela, pallide testacea, opaca, prothorace apicem versus angustato, rosco-tincto, plaga disci gemina notato, arcis hyalinis nullis.* Long. 14—17, lat. $3\frac{1}{2}$ —4 mill. ♀. — Vom See Jipe.

194. *Luciola cisteloides* * Klug, Insekt. v. Mosamb. p. 202. — Von Mombas.

195. *Luciola bimaculicollis* Bohem., Insect. Caffrar. I. p. 445, No. 487. — Von Mombas und dem See Jipe.

196. *Luciola linearis*, n. sp. *Elongata, parallela, dilute fusca, fronte tota prothoracisque aurantiaci macula oblonga subapicali nigris, scutello, elytrorum limbo pedumque basi testaccis.* Long. $7\frac{1}{2}$ mill. ♂. — Vom See Jipe.

197. *Luciola cineticollis* * Klug, Insekt. v. Mosamb. p. 203. — Vom See Jipe.

198. *Luciola lacta*, n. sp. *Nigra, supra fere glabra, prothorace lacte miniacco, profunde sulcato, reflexo-marginato, basi truncato, disperse punctato, nitido: pectore, ventre, coxis femoribusque aurantiacis.* Long. $6\frac{1}{2}$ mill. ♀. — Von Mombas.

199. *Lycus constriectus* Bohem., Insect. Caffrar. I. p. 434, No. 476 (= ♂ *Lyc. cuspidatus* * Klug). — Von Mombas, dem See Jipe und der Insel Sansibar; in copula gefangen.

200. *Lycus congener*, n. sp. *Niger, prothoracis*

abdominisque lateribus nec non elytrorum dimidio maiore aurantiacis: prothoracis lobo apicali acuminato, angulis posticis divaricatis, elytrorum interstitiis quatuor internis aequaliter latis. Long. 19, lat. 8 mill. ♀. — Von Mombas.

201. *Lycus latissimus* Lin. Syst. nat. I. 2. p. 646, No. 14. — Von Mombas.

202. *Lycus gravidulus*, n. sp. *Oblongo-ovatus, niger, nitidus, prothoracis abdominisque limbo laterali nec non elytris — macula apicali bifida excepta — aurantiacis, horum interstitio secundo ceteris latiore. Long. 15, lat. 8 mill. ♀. — Vom See Jipe.*

203. *Eros favosus*, n. sp. *Antennis breviter flabellatis, pedibus compressis: clongatus, niger, opacus, antennarum basi, prothoracis circuitu elytrorumque dimidio maiore flavis: pronoto triareolato, elytris costatis, interstitiis biserialiter foveatis. Long. 7—8 mill. — Von Endara und dem See Jipe.*

204. *Melyris nobilis*, n. sp. *Supra glabra, coerulea, antennarum basi pedibusque cinnabarinis, pectoris lateribus abdominisque basi viridi-micantibus: prothorace confertim cicatricoso-punctato, utrinque carinato, elytris alte costatis, interstitiis quadriseriatim profunde punctatis. Long. 9—11 mill. — Vom See Jipe.*

205. *Melyris parvula*, n. sp. *Viridis vel coerulea, breviter nigro-setulosa, antennarum basi pedibusque rufis: prothorace confertim cicatricoso punctato, utrinque carinato, medio profunde sulcato, elytris alte costatis, interstitiis triseriatim ruguloso-punctatis. Long. 4½—5 mill. — Von Endara.*

206. *Prionocerus dimidiatus*, n. sp. *Ater, prothorace, scutello elytrorumque dimidio basali maiore aurantiacis, apicali cyanescenti: oculis distantibus, antennis breviusculis, articulis 4.—10. subdilatis. Long 12—15 mill. — Von Mombas.*

207. *Prionocerus (Idgia) apicalis*, n. sp. *Aurantiacus, capite posteriore elytrorumque apice nigris: oculis approximatis, antennis filiformibus. Long. 10 mill. — Von Mombas und der Insel Sansibar.*

Fam. Clerii.

208. *Phloeocopus vinctus*, n. sp. *Niger, griseo-hirtus, antennis pedibusque concoloribus, clytris basi sanguineis, fascia pone medium sita, angusta eburnea: prothorace nitido, profunde punctato, ante medium perspicue bicalloso.* Long. 14 mill. ♀. — Vom See Jipe.

209. *Necrobia rufipes* de Geer, Mémoires V. p. 165, No. 1. pl. 15, fig. 4. — Von Wanga und der Insel Sansibar.

Fam. Xylophaga.

210. *Atractocerus brevicornis* Lin. Syst. nat. I, 2. p. 643, No. 11 (= *Lymexylon abbreviatum* Fab. = *Atract. necydaloides* Palis. = *Atract. Madagascariensis* Lap. = *Atract. Africanus* Bohem. = var. *Atract. frontalis* * Klug). — Von der Insel Sansibar.

211. *Xylopertha crinitarsis* Imh., Ber. Verh. naturf. Gesellsch. in Basel V. p. 177 (= *Xylop. sericea* Muls. Wach.) — Von der Insel Sansibar.

212. *Xylographus perforatus*, n. sp. *Prothoracis sinuati angulis anticis rotundatis, niger, glaber, fortiter punctatus, clytris ultra medium usque adscendentibus.* Long. $3\frac{1}{2}$ —4 mill. ♂♀.

Mas maj. *Mandibula sinistra cornuta, vertice profunde excavato.*

Mas min. *Mandibula sinistra subcornuta, vertice plano, disperse punctato.*

Fem. *Mandibula sinistra incrimata, vertice plano, saepe confertim punctato.* — Von der Insel Sansibar.

Fam. Melasoma.

213. *Zophosis sculptilis*, n. sp. *Oblongo-ovata, apicem versus acuminata, capite prothoraceque confertim punctatis, clytris basi subtiliter coriaceis, deinde nitidissimis, parce punctulatis, utrinque fortiter et oblique rugoso-plicatis.* Long. $5\frac{1}{3}$ mill. — Von Mbaramu.

214. *Zophosis congesta*, n. sp. *Acuminato-ovata,*

plerumque aeneo-micans, capite prothoraceque confertim et subtilius, clytris fortius punctatis, tricostatis: fronte leviter impressa, clypei margine subincrassato. Long. 6½—7½ mill.
— Von Mombas, Mbaramu, dem See Jipe u. s. w.

215. *Zophosis alternata*, n. sp. *Obtuse ovata, aterrima, nitida, fronte prothoraceque subtilissime, clypeo distincte separato multo fortius punctato, elytrorum costis tribus latis, laevigatis, interstitiis distincte punctatis. Long. 9 mill.* — Vom See Jipe.

216. *Zophosis funerea*, n. sp. *Oblongo-ovata, atra, subopaca, prothorace subtilissime, capite clytrisque distinctius punctatis, his obsolete tricostatis: fronte clypeum versus leviter bimpressa. Long. 8 mill.* — Mbaramu.

217. *Zophosis convexiuscula** Gerst., Ber. Akad. Wissensch. Berlin 1854, p. 530. — Von der Insel Sansibar.

218. *Diodontes arcolatus*, n. sp. *Oblongo-ovatus, fronte pronotoque rube reticulato-punctatis, hoc late sulcato: elytrorum costis acutis, ramis transversis areolatim inter se conjunctis. Long. 9 mill.* — Von Endara.

219. *Adesmia baccata*, n. sp. *Mandibulis excavatis, capite rugoso-, prothorace fortius reticulato-punctato, huius disco bicalloso, clytris acuminato-ovatis, confertim et subaequaliter seriato-tuberculatis. Long. 16 mill.* — Vom See Jipe.

220. *Rhytidonota stupida*, n. sp. *Antennarum articulo tertio brevi, prothoracis rotundato-quadrati nec-non elytrorum basi marginatis, his illo parum latioribus, secundum suturam fortiter depressis: scutello elevato, transverso. Long. 12—13½, lat. 4—4⅓ mill. ♂♀.* — Von Mombas und dem See Jipe.

221. *Rhytidonota ventricosa*, n. sp. *Antennarum articulo tertio longiore, prothoracis transversim lateribus rotundatis, marginatis, angulis anticis productis, margine basali tenui: scutello punctiformi, clytris acuminato-ovatis, suturam versus depressis, angulis humeralibus reflexis, tuberculatis. Long. 12½—15, lat. 4½—6 mill. ♂♀.* — Von den Bura-Bergen und dem See Jipe.

222. *Rhytidonota gravidula*, n. sp. *Antennarum articulo tertio longiore, prothoracis marginati angulis anticis haud productis, margine basali incrassato: scutello punctiformi, clytris acuminato-ovatis, humeris tuberculatis.* Long. 14—15, lat. 6 mill. — Von Aruscha.

223. *Rhytidonota gracilis*, n. sp. *Prothoracis subcordati margine laterali ante medium obsoleto, processu prosternali brevi, obtuso, clytris oblongo-ovatis, angulis humeralibus simplicibus, rotundatis.* Long. $10\frac{1}{2}$ — $11\frac{1}{2}$, lat. 3 — $3\frac{1}{2}$ mill. — Von Aruscha und Endara.

224. *Pogonobasis platessa*, n. sp. *Antennarum articulo tertio parum longiore quam latiore, nigra, nitida, prothorace clytrisque aequaliter latis, disperse punctatis et fulvo-pilosis, marginibus lateralibus acutis, serratis.* Long. $6\frac{1}{2}$ — $7\frac{1}{2}$ mill. — Vom See Jipe.

225. *Machla hamaticollis*, n. sp. *Prothoracis margine laterali tenui, reflexo, longe ante basin truncato, clytris acuminato-ovatis, cribrato-punctatis, suturam versus scabris, in margine laterali et disco seriatim tuberculatis.* Long. 13 mill. — Vom See Jipe.

226. *Psammodes (Phanerotoma) carbonarius* * Gerst., Ber. Akad. Wissensch. Berlin 1854. p. 532. — Von der Insel Sansibar.

227. *Phrynocolus petrosus*, n. sp. *Robustus, niger, opacus, indumento rufo-brunneo tectus, prothorace utrinque fortiter binodoso, supra deplanato et rube rugoso-punctato, clytris acuminato-ovatis, alte bicostatis, inter suturam et costam superiorem depressis, ubique irregulariter tuberculatis.* Long. 27 mill. — Vom See Jipe.

228. *Phrynocolus frondosus*, n. sp. *Gracilior, niger, cinereo-tomentosus, prothorace utrinque calloso, lineis tribus disci elevatis instructo, clytris supra tricostatis, costis pinnatis, interna abbreviata, media apice hamata calloque semilunari terminata.* Long. 15 mill. — Vom See Jipe.

229. *Phrynocolus plicatus*, n. sp. *Robustus, niger, prothorace rotundato-quadrato, supra confertissime et aequaliter rugoso-punctato, clytris nitidissimis, disperse pun-*

ctulatis, rufescenti-setulosis, latera versus sensim fortius longitudinaliter plicatis. Long. 22—25 mill. ♂♀.

Mas. *Abdominis segmento secundo plaga media rufotomentosa instructo.*

Fem. *Abdominis segmentis duobus primis in medio confertim punctatis et setulosis.* Von Aruscha, Endara und dem See Jipe.

230. *Sepidium muscosum*, n. sp. *Antennis perspicue 11-articulatis, dense cervino-tomentosum, nigro-canoque setosum, prothoracis processu apicali bilobo, dentibus lateralibus magnis, hamatis, elytris subcaudatis, ante apicem fortiter obtuse cornutis.* Long. 18½—20 mill. — Von Endara und dem See Jipe.

231. *Anchophthalmus clathratus*, n. sp. *Obtuse ovatus, niger, parum nitidus, glaber, prothoracis margine deplanato sat lato, extus ruguloso, disco convexiusculo, subtiliter punctato, elytris seriatim foveato-punctatis, interstitiis 3., 5., 7. costatim elevatis.* Long. 13, lat. 6 mill. — Von See Jipe.

232. *Anchophthalmus scutatus*, n. sp. *Oblongo-ovatus, niger, subnitidus, glaber, prothoracis subtiliter punctati margine laterali explanato, apicem versus angustato, elytris punctato-sulcatis, alternatim subcostatis.* Long. 10, lat. 4½ mill. — Vom See Jipe.

233. *Opatrinus trivialis*, n. sp. *Oblongo-ovatus, ater, opacus, prothorace deplanato, elytris anguste et subtiliter sulcatis, interstitiis fere planis, disperse punctulatis.* Long. 11—12 mill. ♂♀.

Mas. *Tibiis anticis arcuatis, intus bisinuatis.* — Von Mombas und der Insel Sansibar.

234. *Opatrinus elevatus*, n. sp. *Oblongo-ovatus, ater, parum nitidus, prothorace convexiusculo, subsulcato, elytris late et profunde punctato-sulcatis, interstitiis convexis, disperse punctulatis.* Long. 8½—10 mill. ♂♀.

Mas. *Tibiis anticis arcuatis, intus emarginatis ibique flavo-fimbriatis.* — Von Kisuani und Endara.

235. *Anomalipus heraldicus*, n. sp. *Oblongus, deplanatus, niger, opacus, prothorace ante medium subhamato-*

dilatato, hexagono, suprabicalloso, elytris sulcatis, interstitiis convexis, subtiliter granulosis, post medium tuberculis magnis, laevibus, in series duas transversas dispositis ornatis, margine laterali fortiter undulato: femoribus simplicibus, tibiis angustis, intermediis extus emarginatis. Long. 26½ mill. ♂.

— Vom See Jipe.

236. *Opatrum contractum*, n. sp. *Alatum, latiusculum, fusco-setulosum, prothoracis lateribus leviter reflexis, basin versus vix sinuatis, elytris obsolete punctato-sulcatis, interstitiis convexis, subtiliter granulatis. Long. 9—11, lat. 4¾—5¾ mill. — Vom See Jipe.*

237. *Opatrum patrule** Erichs., Archiv f. Naturgesch. IX, 1. p. 248, No. 84. — Von Mombas.

238. *Opatrum dermestoides*, n. sp. *Scutello elevato, glabro, oblongum, alatum, fusco-setulosum, prothorace apicem versus sat fortiter angustato, angulis posticis depressis, elytris distincte punctato-sulcatis, interstitiis rugulosis. Long. 7⅔—8, lat. 3¾ mill. — Von Aruscha.*

239. *Opatrum debile*, n. sp. *Oblongum, coriaceum, opacum, parce fulvo-setulosum, antennis, palpis, tibiis tarsisque rufo-brunneis: prothoracis transversus angulis anticis obtusis, haud productis, tibiis anticis apice dilatatis, subuncinatis. Long. 5 mill. — Vom See Jipe.*

Cyptus, nov. gen. (Trib. Opatrini). *Antennae breviusculae, articulis 7.—10. transversis. Oculi divisi. Palporum maxillarium articulus ultimus truncato-ovatus. Prothorax transversus, in medio bascos productus, angulis anticis obtusis, marginibus lateralibus breviter ciliatis. Processus prosternalis lanceolatus. Scutellum transverse triquetrum. Coleoptera ovata, basi angulariter excisa, lateribus ciliatis. Abdominis processus intercoxalis triqueter. Tibiarum calcaria breviuscula. Tibiae anticae digitatae, apice late truncatae.*

240. *Cyptus scabrosus*, n. sp. *Ovatus, convexus, nigro-fuscus, opacus, confertim scabroso-punctatus, prothorace elytrisque rufo-ciliatis, his aequaliter granulosis et sulcatis. Long. 9 mill. — Von der Insel Sansibar.*

241. *Platydemia variipenne* Gemm., Coleopt.

Hefte VI, p. 122 (= *Diaperis variegata* * Klug). — Vom See Jipe.

242. *Uloma pusilla*, n. sp. *Rufo-brunnea, vertice, pronoti disco clytrisque margine basali excepto piceis: prothorace transverso, apicem versus angustato, utrinque fortiter, per discum subtilius et disperse punctato, clytris anguste sulcatis, abdominis lateribus rude eribrato-punctatis.* Long. 6—6½ mill. ♀. — Von der Insel Sansibar.

243. *Alphitobius ovatus* * Herbst, Käfer VIII. p. 16, No. 40. Taf. 118, fig. 8 (= *Tenebr. fagi* Panz.). — Von Mombas, Moschi und Aruscha.

244. *Alphitobius piceus* Oliv. Entom. III, 58. p. 17, No. 22. pl. 2, fig. 13 (= *Tenebr. diaperinus* Panz. = *Tenebr. oryzae* * Herbst). — Von Mombas und der Insel Sansibar.

245. *Alphitobius parallelus* Thoms. Archiv. entom. II. p. 87, No. 125. — Von Urn.

246. *Alphitobius opacus*, n. sp. *Oblongo-ovatus, convexus, niger, opacus, ore, antennis pedibusque rufis: prothorace basi profunde bisinuato, utrinque sulcato, supra subtilissime punctulato, clytris suturam versus obsolete striato-punctatis, iridescentibus.* Long. 4⅔ mill. — Von den Ugono-Bergen.

247. *Hypophloeus volvulus*, n. sp. *Elongatus, rufo-brunneus, capite prothoraceque castaneis, hoc apicem versus sensim dilatato, utrinque subtiliter marginato, clytris striato-punctulatis, seriebus alternis paullo subtilioribus.* Long. 5 mill. — Von den Ugono-Bergen.

Platyotus, nov. gen. (Trib. Ulomidae). *Antennae breviusculae, apicem versus sensim dilatatae. Frons bicornis. Prothorax subquadratus, apice trilobus. Scutellum brevissimum, transversum. Corpus oblongum, parallelum, subdepressum.*

248. *Platyotus glabratus*, n. sp. *Rufo-piceus, glaber, nitidus, antennis pedibusque dilute castaneis, prothorace obscuriore, perspicue punctato, basi lateribusque marginato, clytris distincte punctato-striatis, interstitiis punctulatis.* Long. 3¼ mill. — Von den Ugono-Bergen.

249. *Tribolium ferrugineum* Fab. Spec. Insect. I. p. 324, No. 12. (= *Dermestes navalis* Fab. = *Colydium castaneum* * Herbst.) — Von Mombas.

250. *Cossyphus dentiventris*, n. sp. *Alatus, oblongo-ovatus, prothoracis elytrorumque lateribus fusco-conspersis reflexo-marginatis, horum disco bicostato et utrinque profunde crenato-punctato: femoribus anticis dilatatis, abdominis segmentis duobus ultimis fortiter bidentatis.* Long. 12, lat. 6 mill. ♂. — Vom See Jipe.

251. *Nyctobates brachialis*, n. sp. *Antennis breviusculis, subclavatis, clypeo transversim impresso, prosterno inter coxas dilatato, apice acuminato, pronoti lateribus immarginatis: angustulus, niger, opacus, elytris sat fortiter striato-punctatis, pedibus crassiusculis.* Long. $11\frac{2}{3}$ mill. — Von den Ugono-Bergen.

252. *Lypros breviusculus*, n. sp. *Nigro-piceus, pubescens, punctatissimus, fere opacus, antennarum articulis 5.—10. subquadratis, prothorace transverso, lateribus rotundato, coleopteris ovatis, apicem versus dilatatis.* Long. 7 mill.

Var. *Corpore infra elytrisque dilute brunneis.* — Von Wanga.

Diachastops, nov. gen. *A Lypro differt oculis divisis, antennis validioribus, processu prosternali latiore, prothorace brevi, transverso, coleopteris multo angustiore.*

253. *Diachastops subaeneus*, n. sp. *Rufopiceus, aeneo-micans, punctatissimus, griseo-pubescens, antennis nigris, elytris badiiis, latera versus subrugose punctatis.* Long. 10— $11\frac{1}{2}$ mill. — Von Endara und der Insel Sansibar.

254. *Himatismus trivialis*, n. sp. *Fuscus, confertim punctatus, supra maculatim cervino-pubescens, oculis parum prominentibus, prothorace transverso.* Long. 11—13 mill. ♂♀.

Mas. *Palpis maxillaribus securiformibus, trochanteribus anticis in spinam brevem productis.* — Von Wanga und der Insel Sansibar.

255. *Himatismus Senegalensis* Haag, Coleopt. Hefte VI. p. 89, No. 6. — Von Endara.

256. *Pycnocerus Passerini* Bertol. *Illustr. rer. nat. Mozamb.* II. p. 40, No. 30. fig. 7. — Von Mbaramu.

257. *Aspidosternum festivum*, n. sp. *Viridi-aeneum, glabrum, confertim punctatum, coleopteris purpurascenti-cupreis, obsolete unicostatis, ante apicem ampliatis, prothorace his multo angustiore: antennis validis, cribrato-punctatis, tibiis posticis rectis.* Long. 20 mill. — Von Kiriamu.

258. *Micrantereus variolosus*, n. sp. *Prothorace obsolete punctulato, coleopteris breviter ovatis, supra irregulariter tuberculatis, inter tubercula parce punctatis, opacis pilisque minutis adpressis ferrugineis obtectis: pedibus simplicibus, tibiis rectis.* Long. 15, lat. $9\frac{1}{2}$ mill. ♀. — Von Endara.

259. *Micrantereus femoratus*, n. sp. *Capite prothoraceque distincte punctulatis, nitidis, coleopteris oblongo-ovatis, glabris, aterritis, irregulariter profunde punctatis et verrucosis: pedibus elongatis, femoribus incrassatis, anterioribus subtus dentato-dilatatis, tibiis arcuatis, intermediis brevissimis.* Long. $13\frac{1}{2}$ —15, lat. $7\frac{1}{2}$ —8 mill. ♂♀. — Von Mbaramu und Endara.

260. *Eupezus longipennis*, n. sp. *Ater, subnitidus, glaber, coleopteris oblongo-ovatis, striato-punctatis.* Long. $25\frac{1}{2}$ mill.

Mas. *Tibiis anterioribus intus rufo-barbatis, abdomine glabro, nitidissimo, longitudinaliter plicato, segmento ultimo apice excavato.* — Vom See Jipe.

261. *Gonocnemis brevicollis*, n. sp. *Aterrime, nitida, antennis validis, apicem versus serratis, oculis permagnis, prothorace fortiter transverso, lateribus distincte marginato, inaequaliter disperse punctato, elytris profunde punctato-striatis, interstitiis externis subcarinatis.* Long. 17 mill. — Vom See Jipe.

262. *Praogena nobilis* Mäklin, *Act. soc. scient. Fenn.* VII. p. 568, No. 10. — Von der Insel Sansibar.

263. *Dysgena scabripennis*, n. sp. *Rufobrunnea, flavescenti-pilosa, subopaca, capite prothoraceque fusciscentibus: antennarum articulis 3. et 4. subaequalibus,*

pronoto confertim cicatricoso, elytrorum interstitiis parcius granoso-punctatis. Long. 14—15 mill. — Von der Insel Sansibar.

Miltoprepes, nov. gen. A Praogena differt fronte luta, convexa, a clypeo vix discreta, sulcis intraocularibus nullis: palporum maxillarium articulo terminali vix dilatato: antennarum articulis 4.—10. brevioribus, ovatis: prothorace subcordato, lacvi, coleopteris magis parallelis, subtiliter striato-punctatis.

264. *Miltoprepes laetus, n. sp. Cyaneus, lacvis, nitidus, antennis, tibiis tarsisque nigris, elytris lacte rubris, punctato-striatis, interstitiis planis, disperse subtilissimeque punctulatis. Long. 11—12½ mill. — Vom See Jipe.*

265. *Monomma triplacinum, n. sp. Ovatum, parum convexum, supra nigro-piceum, infra cum pedibus rufo-brunneum, capite prothoraceque confertim, elytris parcius et subtilius punctatis, distincte punctato-striatis, stria saturali obsoletiore. Long. 5½ mill.*

Fam. Lagriariae.

266. *Lagria villosa* Fab. Spec. Insect. I. p. 160, No. 9. — Von Mombas und der Insel Sansibar.

267. *Lagria mollis, n. sp. Antennis gracilibus, pedibus validis, tibiis arcuatis: nigra, dense griseo-pilosa, capite, prothorace abdomineque rufo-brunneis, coleopteris testaceis, confertim punctatis, oblongis, fere parallelis, parum convexis. Long. 10 mill. — Vom See Jipe.*

268. *Lagria semicyanea, n. sp. Antennis apicem versus incrassatis, pedibus gracilibus, nigro-cyanea, griseo-pilosa, supra confertim punctata, elytris dilute brunneis: capite et prothorace subquadratis. Long. 7—8 mill. ♂♀.*

Mas. Prothorace angustiore, elytris aurichalco-micantibus, antennarum articulo ultimo oblongo, 6. et 7. ceteris latioribus.

Fem. Prothorace transverso, antennarum articulo ultimo acuminato-ovato. — Von der Insel Sansibar (und Mosambik).

269. *Lagria plebeja, n. sp. Antennis apicem*

versus incrassatis, pedibus gracilibus, picca, grisco-pilosa, supra confertim punctata, scutello elytrisque rufescenti-testaceis: capite et prothorace subquadratis. Long. 8—9 $\frac{1}{3}$ ♂♀.

Mas. Prothorace angustiore, antennarum articulo ultimo elongato, 8.—10. transversis.

Fem. Prothorace latiore, antennarum articulo ultimo oblongo, acuminato. — Von Mombas und der Insel Sansibar.

270. *Lagria pulverulenta*, n. sp. *Antennis apicem versus incrassatis, pedibus gracilibus, rufo-brunnea, flavescenti-pubescentis: antennis, capite prothoraeque nigro-piceis, hoc transverse quadrato, subsulcato. Long. 7 $\frac{1}{2}$ mill. ♀. — Von der Insel Sansibar.*

Entypodera, nov. gen. *Corpus gracile, glabrum, nitidum. Antennae apicem versus sensim validiores, articulo ultimo oblongo. Caput liberum, rotundato-quadratum. Prothorax capite angustior, oblongus, pone medium fortiter constrictus, bipartitus. Scutellum minutum, oblongo-triquetrum. Coleoptera subparallela, depressa, punctato-sulcata. Femora clavata: tarsorum articulus penultimus bilobus.*

271. *Entypodera anthicoides*, n. sp. *Rufobrunnea, glabra, nitida, capite, antennarum apice pedibusque nigro-piceis, elytris aeneo-fuscis: prothoracis parte antica profunde foveata, postica binodosa. Long. 4 $\frac{1}{2}$ —5 $\frac{1}{2}$ mill. — Von den Ugono-Bergen.*

Fam. Rhipiphoridae.

272. *Rhipiphorus bipunctatus* * Fab. Syst. Eleuth. II. p. 120, No. 17. (= *Rhip. apicalis* Westw. = *Rhip. tropicus* * Nietner). — Von Mombas, Wanga und der Insel Sansibar.

273. *Rhipiphorus flabellatus* * Fab. Spec. Insect. II. Append. p. 501, No. 3—4. (= *Rhip. ferrugineus* * Fab.). — Von Mombas und der Insel Sansibar.

Fam. Vesicantia.

274. *Horia cephalotes* Oliv. Entom. III, 53 bis. p. 5, No. 3. pl. 1, fig. 3. — Von der Insel Sansibar.

275. *Mylabris bizonata* * Gerst., Ber. Akad. Wiss. Berlin 1854. p. 694, No. 3. — Von Wanga und der Insel Sansibar.

276. *Mylabris bifasciata* Oliv. Entom. III, 47. p. 5, No. 3. pl. 1, fig. 10 u. pl. 2, fig. 10 b. — Von Uru und der Insel Sansibar.

277. *Mylabris praestans*, n. sp. *Antennis croceis, capite prothoraceque praeter hirsutiem atram albo-pilosis, elytris glabris, stramineis, summa basi, macularum paribus duobus fasciaeque apicali, antice sinuata nigris.* Long. 23 mill. — Von Endara.

278. *Mylabris amplexens*, n. sp. *Antennis croceis, corpore elytrisque praeter hirsutiem nigram cinereo-pubescentibus, his rufo-brunneis, fasciis tribus flavis, anteriore basali, ampla, humeros amplexente, posterioribus duabus angustis, subundulatis ornatis.* Long. 7—13½ mill. — Von Uru, Wanga und der Insel Sansibar.

279. *Mylabris callicera*, n. sp. *Antennis nigris, femoribus tibiisque rufis, corpore dense cano-sericeo, elytrorum vitta suturae approximata abbreviata, altera laterali angustiore maculaque apicali aurantiacis.* Long. 9—13 mill. ♂♀.

Mas. *Antennis longioribus, fortiter serratis.* — Von Mombas.

280. *Mylabris (Dices) Kersteni*, n. sp. *Antennis croceis, corpore elytrisque praeter hirsutiem nigram cinereo-pubescentibus, his rufo-brunneis, punctatissimis, maculis sex flavis in series duas longitudinales dispositis ornatis.* Long. 13 mill. — Von Wanga.

281. *Mylabris (Dices) Deckeni*, n. sp. *Antennis croceis, corpore elytrisque praeter hirsutiem nigram cinereo-pubescentibus, his rufo-brunneis, punctatissimis, fasciis duabus anterioribus amplis, confluentibus, tertia in maculas duas dissoluta, flavis ornatis.* Long. 14 mill. — Von den Ugono-Bergen.

282. *Mylabris (Dices) dorsalis*, n. sp. *Antennis croceis, corpore elytrisque praeter hirsutiem nigram cinereo-pubescentibus, his punctatissimis, distincte tricostatis, rufo-brunneis, apice nigris, maculis duabus basalibus, anteriore transversa, posteriore longitudinali, fasciisque duabus*

• *angustis, undulatis flavis ornatis.* Long. 9—12 mill. — Von Uru.

283. *Mylabris (Dices) ambigua*, n. sp. *Antennis croceis, corpore elytrisque dense flavescenti-sericeis, prothorace ante basin profunde sulcato, elytris confertim rugoso-punctatis, obsolete tricostatis, vittis duabus, altera dorsali ante medium abbreviata, altera marginali longiore, intus incisa, maculisque tribus dimidii posterioris flavis ornatis.* Long. $8\frac{1}{2}$ — $10\frac{1}{2}$ mill. — Von Mombas und der Insel Sansibar.

284. *Mylabris (Dices) parenthesis*, n. sp. *Antennis croceis, corpore elytrisque dense flavescenti-sericeis, prothorace ante basin sulcato, elytris confertim rugoso-punctatis, obsolete tricostatis, maculis tribus dorsalibus (antere elongata, arcuata) totidemque marginalibus, omnibus secretis, flavis ornatis.* Long. 10 mill. — Von Endara und dem See Jipe.

285. *Lytta strangulata* * Gerst., Ber. Akad. Wiss. Berlin 1854. p. 695, No. 15. — Von der Insel Sansibar.

Fam. Oedemeridae.

286. *Ditylus vittaticollis*, n. sp. *Pubescens, testaceus, capite prothoraceque fere glabris, confertim et parum profunde punctatis, fusco-vittatis, elytris tricostatis, punctatissimis.* Long. 11 mill. — Von der Insel Sansibar.

Fam. Curculionina.

287. *Episus tuberosus*, n. sp. *Prothorace inermi, profunde trisulcato: squamulis griseis undique dense tectus, elytris prope suturam undulato-sulcatis, tuberculis disci duobus, apicis obtusi quinque obsitis.* Long. $12\frac{1}{2}$ mill. — Von Uru.

288. *Microcerus spiniger* * Gerst., Monatsber. Akad. Wiss. Berlin 1855. p. 83, No. 6. — Von Mombas.

289. *Blosyrus angulatus*, n. sp. *Fronte binodosa, prothorace basin versus angustato, supra vermiculoso calloque longitudinali instructo, coleopteris rotundato-quadratis, pone basin oblique truncatis, sutura fasciisque duabus abbreviatis cinereo-squamulosis, interstitio secundo ante medium,*

quarto ante apicem altius elevatis. Long. 9 mill. — Von Wanga.

Synaptoplus, nov. gen. (Trib. Brachyderidae). *Maxillae obtectae. Prothorax haud lobatus. Sulci antennales laterales, lineares. Antennae breviusculae, scapo marginem oculorum anticum vix attingente. Corbiculae tibiatarum posticarum obtectae. Unguiculi omnino coaliti. Scutellum minutum, coleoptera breviter ovata. Corpus dense squamulosum.*

290. *Synaptoplus cervinus*, n. sp. *Corpore elytrorumque lateribus cano-, horum dorso cervino-squamosis, prothoracis linea media denudata, elytris sat profunde striato-punctatis, ante apicem leviter trinodosus, sutura retrorsum subcarinata.* Long. 10 mill. — Von Mombas und Wanga.

291. *Mitophorus aeneipennis*, n. sp. *Antennarum scapo gracili, recto, prothorace oblongo, ante medium leviter dilatato, transverse rugoso, coleopteris oblongo-ovatis, aeneo-micantibus, subtilissime fusco-pubescentibus, profunde punctato-sulcatis.* Long. 7 mill. — Von Endara.

292. *Mitophorus inflatus*, n. sp. *Antennarum scapo valido, arcuato, prothorace subquadrato, granuloso, coleopteris subglobosis, profunde sulcatis, aeneo-micantibus, griseo-pubescentibus.* Long. 7 mill. — Von Endara.

293. *Siderodactylus falciger*, n. sp. *Mandibulis elongatis, falciformibus, breviusculus, niger, subnitidus, squamulis viridi-coerulcis parce obsitus: prothorace basin versus rotundato-dilatato, confertim tuberculato, profunde sulcato, elytris punctato-sulcatis.* Long. 8½ mill. ♂. — Von Endara.

Diatmetus, nov. gen. (Trib. Otiorrhynchidae). *Maxillae obtectae. Prothorax haud lobatus. Scrobiculi antennales superi, arcuati. Antennarum scapus validus, prothoracem medium attingens, funiculus illo longior, septemarticulatus, articulis duobus primis elongatis. Rostrum apice alatum, a fronte separatum. Oculi prominuli. Scutellum inconspicuum. Coleoptera ovata, apice singulatim rotundata, angulis basalibus productis. Corbiculae tibiatarum posticarum obtectae. Unguiculi basi coaliti. Abdominis segmentum secundum sequentibus multo longius. Corpus subdepressum, dense squamulosum.*

294. *Diatmetus praemorsus*, n. sp. *Cervino-squamosus*, *antennarum funiculo tarsisque coerulescentibus*, *elytrorum sutura denudata, aterrima, nitidissima, basin versus tuberculata*. Long. $12\frac{1}{2}$ mill. — Von Kisuani.

Chaunoderus, nov. gen. (Trib. Otiorrhynchidae). *Maxillae obtectae*. *Prothorax haud lobatus*. *Scrobiculi antennales superi, recti*. *Antennarum scapus validus, oculorum marginem posticum attingens, funiculus breviusculus, septemarticulatus, articulis duobus primis oblongis*. *Rostrum subparallelum, a fronte separatum*. *Oculi prominuli*. *Scutellum apertum*. *Coloptera acuminato-ovata, angulis basalibus subproductis*. *Corbiculae tibiarum posticarum obtectae*. *Unguiculi basi coaliti*. *Abdominis segmentum secundum sequentibus longius*. *Corpus parcius squamulosum*.

295. *Chaunoderus stupidus*, n. sp. *Niger, nitidus, cinereo-squamulosus, prothorace alutaceo, elytris profunde striato-punctatis, apicem versus sulcatis*. Long. $9-10\frac{1}{2}$ mill.

Mas. *Corpore densius squamuloso, elytris tuberculo humerali instructis, sutura posteriore squamis magnis, verruciformibus obtecta: abdominis segmento ultimo acuminato, ante apicem calloso*.

Fem. *Corpore parcius squamuloso, elytris simplicibus: abdominis segmento ultimo plano, obtuso*. — Von Endara.

Sphrigodes, nov. gen. (Trib. Otiorrhynchidae). *Maxillae obtectae*. *Prothorax haud lobatus*. *Scrobiculi antennales superi, apicales*. *Antennae elongatae, filiformes, scapo prothoracis medium attingente, funiculi articulo secundo longissimo, clava oblonga, acuminata*. *Rostrum a fronte separatum, apicem versus dilatatum*. *Scutellum inconspicuum, coleoptera rotundata*. *Corbiculae tibiarum posticarum obtectae*. *Unguiculi basi coaliti*. *Abdominis segmentum secundum sequentibus duobus parum longius*.

296. *Sphrigodes margaritaceus*, n. sp. *Fronte profunde sulcata, corpore elytrisque dense cretaceo-, horum dorso interdum cinereo-squamoso, punctorum striis apicem versus subsulcatis, 4.—7. abbreviatis*. Long. $6\frac{2}{3}-7$ mill. — Von Endara.

297. *Pyctoderes plicatus*, n. sp. *Niger, dense viridi-squamosus, antennis pedibusque ferrugineis: prothorace transverso, apicem versus fortiter angustato, elytrorum interstitiis 5. et 7. basi costatim elevatis. Long. 5 mill. — Von Moschi.*

298. *Embrithes muscosus*, n. sp. *Infra cinereo-, supra fusco-squamosus, antennis setosis, capite et prothorace supra pilis erectis, squamiformibus dense vestitis, elytris punctato-sulcatis, tubercula dua basalia denudata gerentibus. Long. 9½ mill. — Vom See Jipe.*

Systates, nov. gen. (Trib. Otiorrhynchidae). *Maxillae obtectae. Prothorax haud lobatus. Scrobiculi antennales superi, ampli. Antennae elongatae, filiformes, scapo prothoracis apicem superante, funiculi articulis duobus primis longioribus, clava angusta, acuminata. Rostrum a fronte separatum. Scutellum inconspicuum, coleoptera ovata. Corbiculae tibiarum posticarum apertae. Unguiculi basi coaliti. Abdominis segmenta tria intermedia subaequalia.*

299. *Systates pollinosus*, n. sp. *Antennis longissimis, scapo tenui, clavato, prothoracis medium fere attingente: niger, subopacus, tenuiter cano-pubescent, prothorace varioloso, elytris ovatis, fortiter seriato-punctatis, interstitiis transverse rugosis. Long. 8—11½ mill. ♂♀.*

Mas. Angustior, tibiis intus emarginatis ibique canopilosus, abdominis segmento secundo basi, quinto apice impresso. — Von Endara, Mombas und der Insel Sansibar.

300. *Systates seminudus*, n. sp. *Antennis breviusculis, niger, subnitidus, parce albido-setulosus, elytrorum apice densius cervino-villoso: prothorace confertim varioloso, elytris basi marginatis, supra profunde seriato-punctatis, interstitiis subtiliter coriaceis. Long. 7—8½ mill. ♂♀.*

Mas. Angustior, tibiis arcuatis et intus longe pilosis, abdominis basi impressa.

301. *Systates amplicollis*, n. sp. *Antennis filiformibus, scapo tenui, clavato, elongatus, niger, dense cervino-squamulosus, prothorace fortiter rotundato-ampliato, disperse tuberculato, coleopteris oblongo-ovatis, punctato-sulcatis, interstitiis dorsalibus breviter fusco-setulosus. Long. 8 mill. ♂.*

Mas. *Tibiis anticis apice fortiter incurvis, intermediis intus incisus, posticis arcuatis, ante apicem appendiculatis.* — Von Endara.

302. *Systates hystrix*, n. sp. *Antennis gracilibus, niger, cinereo-squamulosus, prothorace angusto, supra tuberculato, coleopteris ovatis, setis sat longis albidis seriatim obsitis.* Long. 8 mill. ♀. — Von Mombas.

303. *Systates amoenulus*, n. sp. *Antennis filiformibus, fronte latissima, niger, dense albo-griseoque squamulosus, prothorace subcylindrico, coleopteris breviter ovatis, cervino-maculatis, interstitiis seriatim setulosis.* Long. 4 — $4\frac{3}{4}$ mill. ♂♀.

Mas. *Angustior, abdominis segmento basali leviter impresso.* — Von der Insel Sansibar.

304. *Systates hirtus*, n. sp. *Niger, dense cinereo-squamulosus, capite prothoraceque brevius, elytris longius setosis, his sat fortiter striato-punctatis: antennarum scapo incrassato.* Long. $5\frac{2}{3}$ — $6\frac{1}{3}$ mill. ♂♀.

Mas. *Angustior, abdominis segmento basali deplanato, tibiis intus parce pilosis, intermediis spinulosis.* — Von Mombas und der Insel Sansibar.

Peribrotus, nov. gen. (Trib. Otiorrhynchidae). *Maxillae obtectae. Prothorax haud lobatus. Scrobiculi antennales superi, profundi. Rostrum apice dilatatum, a fronte separatum. Antennae breviusculae, setosae, scapo valido, funiculi articulis 3.—7. abbreviatis, clava acuminato-ovata, solida. Oculi superi, rotundati. Scutellum inconspicuum, coleoptera ovata. Abdominis segmentum secundum sequentibus vix longius. Tibiarum posticarum corbiculae apertae. Unguiculi basi coaliti.*

305. *Peribrotus pustulosus*, n. sp. *Niger, opacus, hic inde squamulis cervinis variegatus, prothorace transverso, supra tuberculato, coleopteris acuminato-ovatis, punctorum seriebus ultra medium usque tubercula laevia, nitida gerentibus.* Long. 8 mill. ♀. — Von Mombas.

306. *Brachycerus atrox*, n. sp. *Oblongus, squamulis rufescenti-fuscis indutus, rostro utrinque biplicato, orbitis planis, prothorace transverse hexagono, supra fortiter*

bicalloso, elytris rude reticulatis, supra biscriatim tuberculatis, tuberculis exterioribus conicis, acutis: femoribus tibiisque annulatim cervino-squamosis. Long. 20 mill. — Von der Insel Sansibar.

307. *Cleonus sannio* * Hbst., Käfer VI. p. 104, No. 65. Taf. 66. fig. 9 (= *Cl. mucidus* Germ. = *Cl. velatus* Schönh.). — Vom See Jipe.

308. *Cleonus mitis*, n. sp. *Rostro obtuse angulato, supra profunde bisulcato, prothorace fortiter constricto, scutello vix conspicuo: cinereo-tomentosus, prothoracis plaga magna media triquetra fasciisque duabus lateralibus, elytrorum maculis duabus dorsalibus obliquis femorumque annulo antecapicali fuscis.* Long. 17—20 mill. — Vom See Jipe (und aus Sennaar).

309. *Lixus haerens* Schönh., Gen. et spec. Curcul. VII. p. 462, No. 103. (? = *Lix. aethiops* Hbst.). — Von Aruscha.

310. *Lixus coarctatus* * Klug, Insekt. v. Madag. p. 110, No. 170 (= *Lix. rhomboidalis* Schönh. = *Lix. compressicollis* Thoms.). — Von Endara.

311. *Lixus sulcirostris*, n. sp. *Elongatus, rostro basin versus bisulcato, prothoracis conoidei dorso confertim rugoso, elytris profunde striato-punctatis, cinereo-nebulosis, apice leviter oblique truncatis.* Long. 11½ mill. — Vom See Jipe.

312. *Lixus pinguis*, n. sp. *Oblongo-ovatus, rostro breviusculo, fronte foveolata, prothorace confertim punctulato punctisque majoribus obsito, coleopteris cuneatis, fortiter striato-punctatis, basi profunde impressis, nitidis, fasciis duabus — anteriore laterali — apiceque densius cinereo-pubescentibus.* Long. 10½, lat. 4⅔ mill. — Von Aruscha.

313. *Lixus baculus*, n. sp. *Elongatus, subcylindricus, rostro angusto, carinato, prothorace profunde sulcato, apicem versus attenuato, elytris apice rotundatis, supra distincte sulcatis, maculatim cinereo-pubescentibus, interstitiis convexis, coriaceo-rugulosis.* Long. 18 mill. — Vom Kilimandscharo, 8000' hoch.

314. *Apoderus submarginatus* Schönh., Gen. et spec. Curcul. V. p. 282, No. 11. — Von Wanga.

315. *Apoderus nigripennis* Fab., Entom. syst. I, 2. p. 386, No. 8. — Von der Insel Sansibar.

316. *Apion nasua*, n. sp. *Rostro valido, utrinque ante oculos sulcato, frontem versus deplanato, opaco, prothorace conoideo, ante medium constricto, basin versus leviter foveato, supra disperse punctato, subnitido, coleopteris obtuse ovatis, alte convexis, ad humeros angulatis, distincte sulcatis, interstitiis planis, subtiliter alutaceis.* Long. (c. rostr.) 3 mill. — Von Mombas.

317. *Apion gallinula* n. sp. *Rostro valido, punctatissimo, utrinque subsulcato, fronte impressa, prothorace constricto, apicem versus leviter attenuato, supra transversim ruguloso, longitudinaliter sulcato, coleopteris oblongo-ovatis, anguste sulcatis, humeris parum prominentibus.* Long. (c. rostr.) 3 mill. — Von Mombas.

318. *Alcides convexus* Oliv., Encycl. méth. V. p. 507, No. 167. (= *Rhynch. excavatus* Oliv. = *Curc. gibbus* Fab. = *Alcid. gibbus* Klug). — Von Aruscha.

319. *Alcides exilis* Schönh., Gen. et spec. Curcul. III. p. 613, No. 8. — Von Endara.

320. *Alcides obsoletus*, n. sp. *Oblongus, niger, opacus, subtus cano-squamosus, femorum basi elytrisque sanguineis, his fortiter striato-punctatis, interstitiis 3., 5., 7. ante apicem strigatim albo-squamulosis: rostro supra disperse punctato, nitido, prothorace brevissimo, ante apicem constricto, confertim rugoso-punctato.* Long. (c. rostr.) 8 mill. ♀. — Von Mombas.

321. *Phaenomerus leucogrammus*, n. sp. *Elongatus, niger, subnitidus, antennis clara excepta, tibiis tarsisque rufo-brunneis: rostro cylindrico, aciculato, prothorace punctatissimo, huius fascia media angulisque posticis, elytrorum basi, fascia media arcuata suturaque posteriore, scapulis pectoreque nigro-pilosis.* Long. $4\frac{1}{2}$ mill. — Von den Ugono-Bergen.

322. *Sphadasmus figuratus*, n. sp. *Ovatus, niger, confertim punctatus, supra maculatim, infra aequaliter et densius cervino-squamulosus, antennis rufo-brunneis, rostri linea media laevi, prothorace transverso, apicem versus*

distincte carinato, elytris punctato-sulcatis, interstitiis 1., 3., 5. elevatis. Long. 5 $\frac{1}{3}$ mill. — Vom See Jipe.

Psalistus, nov. gen. (Trib. Zygopidae). *Antennarum funiculus 7-articulatus, articulo primo crassiusculo, secundo oblongo, ceteris brevissimis. Prosternum ante coxas distincte canaliculatum. Pleurae metathoracicae parallelae. Abdominis segmenta 1. et 2. ampla, medio connata. Pygidium fere obtectum. Femora compressa, fortiter dentata, postica haud elongata.*

323. *Psalistus sordidus*, n. sp. *Oblongo-ovatus, niger, opacus, pilis crassiusculis supra cervinis, infra cinereis vestitus, antennis rufis, clava fusca. Long. 4 mill. — Von Mbaramu.*

324. *Baridius Wahlbergi* Schönh., Gen. et spec. Curcul. VIII. p. 123, No. 26. — Von Endara.

325. *Baridius speciosus*, n. sp. *Ovatus, laetecyanus, punctatissimus, glaber, subnitidus, rostro confertim rugoso-punctato, prothorace transverso, apice fortiter constricto, elytris profunde punctato-sulcatis, sulcis viridi-micantibus, interstitiis planis, seriatim punctatis. Long. 6 mil. — Von den Bura-Bergen.*

326. *Baridius sculptilis*, n. sp. *Oblongus, niger, punctatissimus, supra in punctis subtilissime setulosus, infra plagiatis cano-squamosus, antennis tibiisque rufo-piceis: prothorace oblongo, apicem versus angustato, elytris profunde punctato-sulcatis, interstitiis uniseriatim punctatis, basin versus sulcis vix angustioribus. Long. 5 $\frac{1}{2}$ mill. — Von Mombas.*

327. *Calandra oryzae* Lin. Cent. Insect. p. 12, No. 19. — Von den Bura-Bergen.

328. *Calandra linearis* Hbst., Käfer VII. p. 5, No. 23. Taf. 100, fig. 1. — Vom See Jipe und von den Ugono-Bergen.

329. *Cossonus abscissus* Schönh., Gen. et spec. Curcul. IV. p. 1036, No. 43. — Von Aruscha.

330. *Cossonus procerus*, n. sp. *Rostro carinato, prothoracis confertim punctati linea media laevi, basi bisinuata, disco postice profunde bimpresso: abdomine, pedibus elytrisque rufis, his nigro-limbatis, profunde striato-*

punctatis, interstitiis angustis, uniseriatim punctulatis. Long. $7\frac{1}{2}$ —8 mill. — Von Endara.

Fam. Brentidae.

331. *Ceoecephalus pieipes* Oliv., Entom. V, 84. p. 442, No. 15. pl. 2, fig. 18. — Von Uru, Kisuni und Endara.

Fam. Bostrichidae.

332. *Bostrichus ferrugineus* * Fab. Syst. Eleuth. II. p. 388, No. 20. (= *Bostr. xylographus* Say). — Von Mombas.

333. *Diamerus pulverulentus*, n. sp. *Ob-longo-ovatus, niger, punctatissimus, cinereo-setulosus, antennis tarsisque rufis: clypeo impresso, vertice subtiliter sulcato, prothorace transverse quadrato, basin versus subcarinato, coleopteris fere parallelis, latera versus profundius sulcatis.* Long. $3\frac{1}{2}$ — $4\frac{1}{4}$ mill. — Von den Bura-Bergen, dem See Jipe und Endara.

Fam. Anthribidae.

334. *Phloeobius pustulosus*, n. sp. *Cervino-cinereoque tomentosus, prothorace ante basin quadrifariam impresso, fasciculis setarum rufarum quinque ornato: elytris fasciculos binos basales fuscos maiores, antecapicales minores miniaceos septem gerentibus, tibiis tarsisque rufescentibus.* Long. 9—10 mill. ♂. — Vom See Jipe.

Fam. Longicornia.

335. *Cantharoctenus insignis*, n. sp. *Rufopiceus, supra glaber, nitidus, antennis 20-articulatis, mandibulis basin versus bidentatis, coxis, femorum basi tibiisque intus dense fulvo-pilosis.* Long. 28 mill. ♂. — Von Endara.

336. *Macrotoma palmata* Fab., Entom. syst. 1, 2. p. 249, No. 26. (= *Prion. Senegalensis* Oliv.) — Von der Insel Sansibar.

337. *Macrotoma Natala* (Dej.) Thoms., Essai d'une classif. d. Céramb. p. 314, No. 177. — Von den Bura-Bergen.

338. *Macrotoma micros* White, Catal. Longicorn. Brit. Mus. I. p. 39, No. 15. — Von der Insel Sansibar.

339. *Xystrocera globosa* Oliv., Entom. IV, 67. p. 27, No. 30. pl. 12, fig. 81. — Von Mombas.

340. *Cordylomera spinicornis* Fab., Syst. Entom. p. 167, No. 13 (= *Ceramb. torridus* Oliv.). Var. *nitidipennis* (Dej.) Serville, Annal. soc. ent. de France III. p. 24. No. 1. — Von der Insel Sansibar.

341. *Compsomera fenestrata*, n. sp. *Capite prothoraceque rubris, hoc obsolete nigro-signato, elytris obscure cyaneis, nigro-pictis, macula obliqua triquetra testacea ornatis.* Long. 18 mill. ♀. — Von Endara.

342. *Rhopalizus Sansibarius*, n. sp. *Aurco-vel cyanescenti-viridis, subnitidus, antennis pedibusque nigris, femoribus basi apiceque exceptis laete rubris: prothorace minus confertim plicato, elytris alutaceis, scutellum versus distincte punctatis.* Long. 23—26 mill. ♂♀. — Von der Insel Sansibar.

343. *Mastododera nodicollis* * Klug, Insekt. v. Madag. p. 121. No. 196. Taf. V. fig. 9 (= *Toxotus basalis* Dej. = *Toxot. lateralis* Guér.). — Von Mombas.

344. *Diatomocephala simplex* Gyllenh. in Schönh. Synon. Insect. III. Append. p. 178, No. 248 (= *Diat. maculaticollis* Blanch. = *Callid. vile* Newm. = *Arhopalus ambiguus* Newm. = *Hesperophanes guttaticollis* Fairm.). — Von der Insel Sansibar.

345. *Phantasis gigantea* Guér. Iconogr. règne anim., Insect. p. 241. — Vom See Jipe.

346. *Coptops aedificator* Fab. Entom. syst. I, 2. p. 275, No. 33 (= *Ceramb. fuscus* Oliv.). — Von Mombas und der Insel Sansibar.

347. *Sternotomis Bohemani* Westw., Arcan. entom. II. p. 154. pl. 85, fig. 3 (= *Sternot. Ferreti* Westw., Reiche). — Von der Insel Sansibar und Mombas.

348. *Phryneta spinator* Fab. Entom. syst. I, 2. p. 276, No. 36. — Von der Insel Sansibar.

349. *Ceratites jaspidea* Serv., Ann. soc. entom. de France IV. p. 35, No. 1. — Vom See Jipe.

350. *Tragocephala variegata* Bertol., Illustr.

rer. nat. Mozamb. I. p. 25, No. 21. tab. 1, fig. 9. (= *Trag. venusta* Thoms.) — Von Mombas und der Insel Sansibar.

351. *Frea marmorata*, n. sp. *Fronte angusta, capite prothoraceque cano-tomentosis, hoc supra nigro-consperso, elytris basin versus fortiter punctatis, fusco-cinereoque nebulosis, plagis duabus, altera basali, altera antecapicali dense flavescenti-tomentosis.* Long. 14 mill. — Von Mbaramu.

352. *Crossotus barbatus*, n. sp. *Antennarum articulis tribus primis infra dense fimbriatis, prothorace brevi, apicem versus fortiter angustato: cano-tomentosus, elytris cinereis, cervino-strigosis, basin versus profunde seriatim punctatis, litura albida tuberculoque fusco signatis.* Long. 11 mill. ♀. — Vom See Jipe.

353. *Niphona appendiculata*, n. sp. *Subdepressa, cinerco-tomentosa, prothorace quinqueseriatim tuberculato, elytris basin versus cristatis, apice excisis, angulo suturali appendiculato.* Long. 14—16 mill. — Vom See Jipe.

354. *Sthenias cylindrator* Fab. Syst. Eleuth. II. p. 292, No. 57. — Von Mbaramu.

Phoryctus, nov. gen. (Trib. Apomecynidae). *Corpus latiusculum, subdepressum. Caput angustum, vertice declivi, a fronte vix separato. Antennae corpore dimidio breviores, articulo tertio subarcuato. Prothorax transversus, muticus, apicem versus angustatus. Elytra prothorace latiora, apice truncata et quasi appendiculata, supra multifariam tuberculata. Prosternum retrorsum fortiter dilatatum, mesosternum transversum, trapezoideum. Pedes breviusculi, validi, tibiis posterioribus arcuatis, compressis, apicem versus extus ciliatis et dilatatis, intermediis intus angulatis: femoribus posterioribus subdentatis.*

355. *Phoryctus mucoreus*, n. sp. *Cervino-tomentosus, albo-variegatus, elytrorum tuberculis rufo-sectosis.* Long. 12 mill. — Von Mombas.

356. *Praonetha melanura* Pasc., Transact. entom. soc. of London 2. ser. IV. p. 106. — Von der Insel Sansibar.

357. *Praonetha scripta*, n. sp. *Cervino-pubescentis, prothorace fortiter punctato, transverse calloso, elytris latera versus albo-fuscoque pictis, supra fasciculorum seriebus duabus longitudinalibus ornatis.* Long. $6\frac{2}{3}$ mill. — Von Endara.

358. *Hippopsicon virgatum*, n. sp. *Nigro-piceum, confertim punctatum, cinereo-pubescentis, prothoracis lateribus elytrorumque margine suturali densius albo-tomentosis, his rufo-piceis, trifariam flavescenti-lineatis.* Long. $10\frac{1}{2}$ mill. — Vom See Jipe.

359. *Hippopsicon rusticum*, n. sp. *Antennarum articulis 3.—6. fimbriatis, breviusculum, piceum, sericeum, vertice bicalloso, prothorace disperse sat fortiter punctato, ante basin constricto, elytris submaculatis pubescentibus.* Long. 8 mill. — Vom See Jipe.

360. *Tetraglenes phantoma*, n. sp. *Antennis corpore brevioribus, parce fimbriatis, picea, confertim punctata, maculatis cano-tomentosa, coleopteris lanceolatis, apice profunde excisis.* Long. 10 mill. — Von der Insel Sansibar.

Fam. Chrysomelinae.

361. *Lema hottentota* Lacord., Monogr. Coléopt. subpent. I. p. 326, No. 19. — Von der Insel Sansibar.

362. *Lema foveicollis*, n. sp. *Prothorace fortiter constricto et transverse sulcato, apice utrinque tuberculato, ante medium profunde bifoveato, basin versus plicato: nigro-aenea, griseo-pubescentis, elytris cyaneis vel purpurascenti-cupreis.* Long. $6\frac{1}{2}$ mill.

Var. a. *Capite prothoraceque viridi-aeneis.*

Var. b. *Occipite, vertice prothoraceque sanguineis.* — Von Mombas und Mbaramu.

363. *Lema morosa*, n. spec. *Breviuscula, nigra, subopaca, dense cinereo-pubescentis, prothorace elytrisque rufo-brunneis, illo ante basin tenuiter sulcato, subtiliter et sat dense, in dorso medio fortius punctato, elytris convexiusculis, fortiter striato-punctatis, interstitiis subtilissime coriaceis.* Long. $5\frac{1}{2}$ mill. — Von der Insel Sansibar.

364. *Lema nigri ventris*, n. sp. *Supra cum pedibus antennisque testacea, capitis rufo-ferruginei macula ver-*

ticali, lubro, pectore abdomineque nigris, griseo-pubescentibus: prothorace elytrisque nitidissimis, illo disperse punctato, ante basin sulcato. Long. 4½ mill. — Von der Insel Sansibar.

365. *Diapromorpha haemorrhagica*, n. sp. *Subtus dense cano-sericea, supra glabra, disperse punctata, nitida, nigra, prothoracis fascia subapicali utrinque ad basin usque producta, elytrorum duobus — altera basali, altera submedia ad humeros usque extensa — saturate flavis, horum maculis duobus subapicalibus lacte rubris: tibiis tarsisque rufis, abdomine cyanescente. Long. 11—14 mill. ♂♀. — Von der Insel Sansibar.*

366. *Clythra Wahlbergi* Lacord., Monogr. d. Coléopt. subpent. II. p. 197, No. 5. — Von Mombas.

367. *Melitonomalitigiosa* Lacord. ibid. II. p. 378, No. 7. — Von Mombas.

368. *Gynandrophthalma asphaltina*, n. sp. *Aterrima, infra albido-sericea, supra glabra, nitidissima, antennarum basi, genibus tarsisque piceis, elytris violaceomicantibus, disperse et subtilius punctatis: capite inter oculos transverse impresso, fronte bifoveolata. Long. 4—4⅔ mill. ♂♀. — Von der Insel Sansibar.*

369. *Gynandrophthalma amoenula*, n. sp. *Oblonga, apicem versus leviter dilatata, rufo-ferruginea, supra glabra, nitida, elytris disperse punctatis, testaceis, sutura, vitta humerali ante apicem abbreviata, marginis lateralis parte posteriore nec non scutello pleurisque nigris. Long. 4 mill. — Von Mombas und der Insel Sansibar.*

370. *Damia* (?) *confusa*, n. sp. *Clypeo truncato, oblonga, subparallela, nigra, subtus griseo-pubescentibus, supra glabra, nitida, antennis tibiisque basi ferrugineis, prothorace elytrisque lacte flavis, hoc maculis duabus magnis basalibus, inter se connexis, his punctis tribus anterioribus (1. 2.) fasciaque posteriore utrinque abbreviata nigris. Long. 7—7½ mill. — Von der Insel Sansibar.*

371. *Cryptocephalus tabidus*, n. sp. *Vertice angustato, antennarum articulis sex ultimis elytrorumque limbo basali nigro-piceis, ceterum dilute testaceus, nitidus, elytris stramineis, fortiter striato-punctatis, interstitiis disperse punctulatis. Long. 4⅔ mill. ♂. — Von der Insel Sansibar.*

372. *Cryptocephalus Kersteni*, n. sp. *Ferrugineus, elytris dilutioribus, subtiliter acieulatis, pectore, abdomine, prothoracis maculis duabus rotundatis, scutello, elytrorum limbo vittaque discoidali, ante apicem abbreviata, nigris.* Long. $3\frac{1}{3}$ mill. ♀. — Von Mombas.

373. *Cryptocephalus callias* * Suffr., Linn. entom. XI. p. 122, No. 30.

Var. ♀. *Elytris violaceis, macula apicali aurantiaca.* — Von der Insel Sansibar.

374. *Cryptocephalus apertus*, n. sp. *Corpore pedibusque nigris, prothoracis lucidi macula laterali elytris-que — puncto humerali nec non signatura communi cruciata exceptis — laete aurantiacis.* Long. 6— $6\frac{1}{3}$ mill. ♂♀. — Von der Insel Sansibar.

375. *Pachnephorus conspersus*, n. sp. *Oblongus, fusco-aeneus, disperse albo-squamulosus, prothorace truncato-ovato irregulariter, elytris fortius et striatim punctatis, interstitiis suturam et apicem versus latioribus, punctulatis.* Long. 3 mill. — Von Wanga.

376. *Corynodes Dejeani* * Gerst., Insekt. v. Mossamb. p. 336. — Von Mombas.

377. *Scelodonta Sansibarica*, n. sp. *Cupreo-aenea, infra dense, supra variegatim albido-setulosa, prothorace transverso, confertim rugoso-punctato, elytris rude seriatim et basin versus reticulato-punctatis.* Long. 4— $4\frac{1}{2}$ mill. — Von der Insel Sansibar.

378. *Pseudocolaspis chrysites*, n. sp. *Cupreo-aenea, confertim punctata, pilis decumbentibus aureo-micantibus variegatim obsita, prothorace subglobozo, scutello apice trilobo, femoribus omnibus unispinis.* Long. $3\frac{1}{2}$ — $4\frac{1}{2}$ mill. — Von der Insel Sansibar.

379. *Rhyparida collaris*, n. sp. *Rufo-brunnea, glabra, nitida, antennarum basi palpisque testaceis, prothorace infuscato, coleopteris multo angustiore, disperse et subtilissime punctulato, elytris fere parallelis, regulariter striato-punctatis, femoribus omnibus subtus dentatis.* Long. $5\frac{1}{2}$ mill. — Von der Insel Sansibar.

380. *Rhyparida cyanipennis*, n. sp. *Ovata, glabra, nitida, laete ferruginea, capite, prothorace scutelloque*

rufescentibus, antennarum articulis quatuor ultimis nigropiceis, elytris obscure cyaneis, fortiter striato-punctatis, femoribus omnibus acute dentatis. Long. $3\frac{2}{3}$ mill. — Von Mombas.

381. *Rhyparida micans*, n. sp. *Ovata, glabra, nitida, rufo-brunnea, prothorace latiusculo, supra infusato et aeneo-micante, disperse punctato: pedibus testaceis, femoribus omnibus dentatis. $3\frac{1}{4}$ mill. — Von Mombas und der Insel Sansibar.*

382. *Rhyparida trivialis*, n. sp. *Ovata, glabra, nitida, rufo-brunnea, prothorace angustiore, fortius punctato, antennarum articulis quatuor ultimis nigris: pedibus concoloribus, femoribus acute dentatis. Long. 3— $3\frac{2}{3}$ mill. — Von Mombas und der Insel Sansibar.*

383. *Rhyparida obscurella*, n. sp. *Ovata, glabra, nitida, fusco-aenea, antennis, tibiis tarsisque ferrugineis: prothorace coleopteris multo angustiore, sat fortiter disperse punctato, humeris callosis, rufo-piceis. Long. $2\frac{3}{4}$ mill. — Von Mombas.*

384. *Colasposoma subcostatum*, n. sp. *Tibiis anticis rectis, antennarum basi ferruginea, aurichalceum vel metallico-viride, glabrum, nitidum, capite confertim rugoso, prothorace elytrisque laxius profunde punctatis, his obsolete quinquecostatis. $5\frac{2}{3}$ — $6\frac{1}{3}$ mill. — Von Aruscha und der Insel Sansibar.*

385. *Colasposoma compactum*, n. sp. *Breviusculum, aeneum, subnitidum, capite disperse et subtiliter, prothorace fortius punctato, latera versus confertim strigoso, linea media laevi: elytris suturam versus disperse et subtilius, post humeros fortiter rugoso-punctatis. Long. $5\frac{1}{2}$ mill. — Von Mombas.*

386. *Ceralces Natalensis* Baly, Transact. entom. soc. of London 2. ser. IV. p. 88, No. 2.

Var. *Corpore infra, antennarum basi pedibusque ferrugineis. — Von Endara und dem See Jipe.*

387. *Entomoscelis ochroptera* Stål, Vetensk. Akad. Föreländ. XIV. p. 60, No. 1. — Von Mombas und dem See Jipe.

388. *Plagiodera impolita* (Vogel i. lit.). *Bre-*

viter ovata, nigra, supra viridi-cyanca, nitida, antennarum basi, prothoracis lateribus elytrorumque margine externo miniaceis: elytris subtilissime alutaceis, perspicue et disperse punctatis. Long. 6—6½ mill. — Vom See Jipe.

389. *Adorium palliatum*, n. sp. *Acuminato-ovatum, glabrum, atrum, supra opacum, prothorace anoque aurantiacis, illo disperse punctulato, elytris alutaceis, confertim punctatis: abdominis segmento quinto (maris?) apice retuso.* Long. 11 mill. — Von Mombas.

390. *Apophylia nobilitata*, n. sp. *Oblonga, subparallela, opaca, testacea, capitis vitta lata media plagisque duabus postocularibus, prothoracis maculis tribus scutelloque nigris, elytris smaragdinis, confertim granoso-punctatis, margine laterali inaurato.* Long. 7 mill. — Von Endara.

391. *Apophylia murina*, n. sp. *Nigra, confertim punctata, griseo-pubescent, opaca, antennarum basi, coxis femoribusque ferrugineis, capitis linea media laevi.* Long. 5—6 mill. — Von der Insel Sansibar.

392. *Diamphidia pectinicornis* Oliv. Entom. VI. p. 849, No. 13. pl. 1, fig. 10. — Von der Insel Sansibar.

393. *Monolepta flaveola* * Gerst., Monatsber. Akad. d. Wissensch. Berlin 1855. p. 638. — Von Aruscha.

394. *Monolepta rubricosa*, n. sp. *Rubra, subtiliter punctata, nitida, elytrorum fasciis duabus, altera basali, altera antecapicali nigris, antennis, tibiis tarsisque pallide testaceis, illarum articulis duobus ultimis nigro-fuscis.* Long. 4⅔ mill. — Von der Insel Sansibar.

395. *Monolepta didyma*, n. sp. *Pallide testacea, supra glabra, parum nitida, antennis apice fuscis, elytrorum fasciis duabus abbreviatis nigris: elytris subtiliter alutaceis, sat confertim punctatis.* Long. 4 mill. — Von Wanga.

396. *Monolepta vineta*, n. sp. *Testacea, punctata, submitida, capite ventrequae rufo-ferrugineis, antennis apice infuscatis, elytrorum limbo basali et humerali nec non fascia angusta, pone medium sita nigro-piceis.* Long. 4 mill. — Von Mombas.

397. *Monolepta vinosa*, n. sp. *Oblongo-ovata, subtiliter punctata, nitida, testacea, capite, scutello, elytrorum*

sutura et apice pectoreque rufis, vertice elytrorumque fasciis duabus, altera basali, maculari, altera media, vitta marginali inter se conjunctis, nigris. Long. 6 mill. — Vom See Jipe.

398. *Monolepta ephippiata*, n. sp. *Oblonga, punctata, subnitida, testacea, capite, elytrorum fasciis duabus — altera basali, extus abbreviata, altera ante medium sita lata, undulata, ad suturam et marginem lateralem producta — pygidioque nigris, pectore abdomineque rufescentibus.* Long. 4 mill. — Vom See Jipe.

399. *Rhaphidopalpa vinula* * Erichs., Doublett. Verz. 1842, p. 10, No. 188. (= *Galleruca festiva* * Gerst.) — Von Wanga.

400. *Diacantha duplicata*, n. sp. *Flava, supra glabra, nitida, elytris ferrugineis, ad basin late atris.* Long. 8 mill. ♀. — Von Uru.

401. *Xenarthra calcarata*, n. sp. *Tibijs posticis ante apicem longe calcaratis, antennarum articulis 4.—8. arcuatis, tumidulis: lacte rufo-ferruginea, glabra, nitida, elytris subtiliter fusco-punctulatis, ad humeros splendide cyaneis, pedibus antennisque dilute flavis, harum articulis duobus ultimis fuscis.* Long. 6 mill. — Von der Insel Sansibar.

402. *Malacosoma pusilla*, n. sp. *Nigra, glabra, nitida, antennarum basi, fronte, prothorace, elytrorum apice et lateribus nec non pedibus ferrugineis: elytris viridi-micantibus, disperse punctatis et transverse rugulosis.* Long. $3\frac{1}{2}$ mill. — Vom See Jipe und von der Insel Sansibar.

403. *Malacosoma* (?) *litura*, n. sp. *Glabra, nitida, testacea, antennis apicem versus fuscis, prothoracis macula lineari media, elytrorum sutura, pectore abdomineque nigro-piceis: prothorace utrinque oblique punctato-sulcato, elytris subtiliter punctatis, supra marginem lateralem profunde impressis.* Long. $2\frac{1}{2}$ —3 mill. — Von Mombas.

404. *Oedionychis rubiginosa*, n. sp. *Oblongo-ovata, convexa, parum nitida, rufa, antennis basi excepta, mandibularum apice, genibus, tibiis tarsisque nigris: prothorace subtilius et disperse, elytris confertim punctatis.* Long. 9 mill. — Von Moschi.

405. *Oedionychis* (?) *verniciata*, n. sp. *Tarsorum anteriorum articulis duobus primis inflatis: oblongo-*

ovata, parum convexa, glabra, nitidissima, testacea, antennis basi excepta, capite, scutello, elytrorum vitta suturali maculisque duabus discalibus — subhumerali oblonga, posteriore permagna, triquetra — pectore abdominisque basi nigro-piceis: vertice utrinque grosse, prothorace subtiliter punctato. Long. $3\frac{1}{2}$ mill. — Von Uru.

406. *Haltia (Apthona) Cookei*, n. sp. *Ovata, nitida, ferruginea, antennarum apice femoribusque posticis nigro-piceis, femoribus anterioribus fusco-annulatis: elytris subtilissime disperse punctulatis.* Long. 3 mill. — Von Mombas und der Insel Sansibar.

407. *Haltia (Crepidodera) tosta*, n. sp. *Ovata, nitida, rufo-brunnea, prothorace disperse punctulato, ad basin bisulco, elytris acneo-micantibus, striato-punctatis, interstitiis subtilissime punctulatis.* Long. $2\frac{1}{2}$ —3 mill. — Von Mombas.

408. *Argopus pusillus*, n. sp. *Oculis amplis, fronte angusta: breviter ovatus, rufus, disperse punctatus, subnitidus, ore, antennis pedibusque ferrugineis.* Long. $2\frac{1}{2}$ mill. — Vom See Jipe.

409. *Hispa quadrifida*, n. sp. *Antennis prothoracisque disco inermibus, huius lateribus anterieus spinis quatuor longis, basi conjunctis, posterius unica minore obsitis: atra, glabra, elytris leviter acneo-micantibus, irregulariter rude punctatis et multispinosis, spinis dorsalibus pro parte validissimis.* Long. 4— $4\frac{1}{3}$ mill. — Von der Insel Sansibar.

410. *Hispa pachycera*, n. sp. *Antennis crassiusculis, inermibus, prothoracis disco utrinque bi-, lateribus trispinosis: testacea, parce setulosa, elytris profunde seriatim punctatis, interstitiis linearibus, alternis subcostatis, spinis marginalibus gracilioribus, dorsalibus apicem versus conicis, congestis.* Long. $5\frac{1}{3}$ mill. — Von der Insel Sansibar.

411. *Cassida vigintimaculata* Thunb., Nov. Insect. spec. V. p. 86. fig. 95. — Von Endara.

412. *Cassida tigrina* Oliv., Entom. VI. p. 957, No. 56. pl. 5, fig. 78. — Von Endara.

413. *Cassida dorsovittata* * Boh., Monogr. Cassidid. II. p. 395, No. 59. — Von der Insel Sansibar.

414. *Cassida lacrymosa* * Boh., *ibid.* II. p. 397,
No. 61. — Von Mbaramu.

Fam. Erotylidae.

415. *Episcapha scenica*, n. sp. *Nigra, confertim punctata, pubescens, subnitida, prothoracis maculis duabus oblongis, elytrorum annulo humerali fasciisque duabus posterioribus angustis, undulatis sanguineis; antennarum clava angusta, elytrorum striis prope suturam tantum perspicuis.* Long. $12\frac{1}{2}$ mill. — Von Mombas.

Fam. Endomychidae.

416. *Ancylopus unicolor* * Gerst., *Monogr. d. Endomychid.* p. 194, No. 4. — Von Mombas.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Archiv für Naturgeschichte](#)

Jahr/Year: 1871

Band/Volume: [37-1](#)

Autor(en)/Author(s): Gerstäcker [Gerstaecker] Carl Eduard Adolph

Artikel/Article: [Beitrag zur Insektenfauna von Zanzibar. 42-86](#)