**70**(2) 75–83

# A new *Epiophlebia* (Odonata: Epiophlebioidea) from China with a review of epiophlebian taxonomy, life history, and biogeography

FRANK LOUIS CARLE

Rutgers, The State University of New Jersey, Department of Entomology, 96 Lipman Drive, New Brunswick, NJ 08901, USA [Carle@AESOP.Rutgers.edu]

Received 13.ix.2011, accepted 27.viii.2012. Published online at www.arthropod-systematics.de on 28.ix.2012.

### > Abstract

*Epiophlebia diana* **sp.n.** is described from larval specimens collected in the mountains of western Sichuan Province, China. Epiophlebia taxonomy, life history, and biogeography are reviewed, and keys provided for determination of the known adults and larvae of *Epiophlebia* Calvert, 1903. Classification of *Epiophlebia* is revised as follows: *Epiophlebia* s.str. with *E. superstes* (Selys, 1889) [type species] and *E. sinensis* Li & Nel, 2012; and *Rheoepiophlebia* **subgen.n.** with *E. laidlawi* Tillyard, 1921 [type species] and *E. diana* **sp.n.**. Behavioral, ecological and paleontological information is also evaluated and members of *Epiophlebia* acknowledged to have inhabited small high elevation streams of the east Palaearctic for possibly 180 million years. Likely reasons are proposed for the enduring survival of *Epiophlebia*, its lack of a fossil record and the extinction of related groups.

## > Key words

Epiophlebia, Rheoepiophlebia, extinction, biogeography, living fossil.

# 1. Introduction

Traditionally Odonata include Anisoptera (dragonflies), Zygoptera (damselflies), Epiophlebia Calvert, 1903, and a diverse array of fossil forms, among which are both the largest known insects and the oldest known creatures to fly (BRAUCKMANN & ZESSIN 1989; CARPENTER 1992). Fossil dragonflies not only allow the intricate wing venation of Odonata to be correctly homologized with that of other Pterygota (CARLE 1982a; RIEK & KUKALOVÁ-PECK 1984), but offer excellent outgroup taxa for character state polarization in Odonata s.str. (REHN 2003). However, even with an extensive fossil record, the systematic position of the enigmatic Epiophlebia has changed dramatically over the last century. Epiophlebia was considered a zygopteran (Selys 1889; Needham 1903; Tillyard 1914, 1917), an anisopteran (NEEDHAM 1930; LEW 1933), or placed within the artificial group "Anisozygoptera Handlirsch" (HANDLIRSCH 1906-08, 1939; TILLYARD 1925, 1929; TILLYARD & FRASER 1940) along with several Jurassic fossil taxa.

In his monograph on the morphology of *Epiophlebia* ASAHINA (1954) established a separate superfamily for Epiophlebiidae based on its phylogenetic isolation and morphological uniqueness. From related fossil evidence it is apparent that Epiophlebia has been genetically isolated since the Lower Jurassic, and this has resulted in a curious combination of plesiomorphy and autapomorphy. This is illustrated by a few wing features of Epiophlebia: petiole of hind wing well developed, costa widened at nodus, pterostigmata strongly convex posteriorly, and hind wing with one or two cell rows between CuA and wing margin. Epiophlebia also lacks a discal nodus, an often overlooked ventral membranous area along the discal brace, yet some wing flexibility is achieved by a slight flattening of the pleat at the discal brace, and although the costal nodus is hinge-like its flexibility is limited particularly in the forewings by a minimal nodal fissure (CARLE 1982a). Additional autapomorphies include: labrum widened distally, genae with lateral tubercles, antennae five segmented with pedicel elongate and flattened, postfrons with transverse shield-like intraocellar ridge, male occipital region with antitandem lobes, male epiproct with ventral and dorsal rami, male gonocoxae elongate, female abdominal segment 8 with midventral apical spur, and several autapomorphies of the female genitalia listed by KLASS (2008: p. 133). The copulatory apparatus of *Epiophlebia* is bizarre. It includes paired elongate posteriorly directed hamuli with swollen studded hooked apices (penis) and a median process (the homologue of the anisopteran ligula and the zygopteran penis) of abdominal sternum two, and a somewhat bottle shaped sperm vesicle (anisopteran penis) at the anterior end of abdominal sternum three (SCHMIDT 1915; PFAU 1971; CARLE 1982b). Remarkable plesiomorphic character states include: labial end hook occasionally two segmented (though more usually one segmented), well developed glossae and paraglossae, and a trilobate hypopharynx (Asahina 1949). Epiophlebia larvae are unique in that the antennae are five segmented, the head has well developed paralabial ridges, and they can produce sound by rubbing the inner apex of the femora against lateral abdominal files (Figs. 1E, 2E, 3E). Plesiomorphic character states of Epiophlebia larvae include wing pads without a branch of the RA tracheae crossing over the RP trachea, abdomen without transverse muscles (AsaHINA 1954), simple rectal gills, and proventriculus with 16 to 18 well developed denticulate lobes.

Several apomorphies found in extant Odonata support a sister group relationship between Anisoptera and Epiophlebia (CARLE 1982a, 1995; CARLE & KJER 2002; REHN 2003; KLASS 2008), as do diverse molecular markers (CARLE et al. 2008). "Anisozygoptera" was considered polyphyletic by CARLE (1982a), who rearranged the three suborders of Odonata into Zygopteroids and Anisopteroids, the latter of which includes Epiophlebia, based on several characters including tandem hold and associated modifications of the male terminalia. BECHLY (1996) also considered "Anisozygoptera" polyphyletic, but with Tarsophlebioidea Handlirsch, 1908 sister to a group including Zygoptera, Anisoptera and Epiophlebia. In contrast, NEL et al. (1993), BECHLY (1995) and (FLECK et al. 2004) represented "Anisozygoptera" as merely paraphyletic, with Tarsophlebioidea sister to Anisopteroids.

The genus *Epiophlebia* currently includes three described species: *Epiophlebia laidlawi* Tillyard, 1921 from India, Nepal and Bhutan; *Epiophlebia superstes* (Selys, 1889) from Japan; and *Epiophlebia sinensis* Li & Nel, 2012 recently described from northeastern China (LI et al. 2012). Adult *Epiophlebia* are rather rare in collections perhaps due to short flight seasons, isolated habitats, and atypical lifestyles. However, the first larval specimens of *Epiophlebia* from India, China and Japan were already collected before 1930 (TILLYARD 1921; NEEDHAM 1930; CARLE 1983a), and the second known Epiophlebia species, E. laidlawi, was described based on the original larval specimen collected in India. Here, a new species of Epiophle*bia* is described from larval specimens collected by "Dr. David C. Graham in the mountainous regions of western Szechuan" (NEEDHAM 1930). Unfortunately, Needham misidentified larvae of Epiophlebia as Gomphidae (CARLE 1983a), and the significance of these larvae from south-central China was not realized. The new species is compared to larval specimens of both E. superstes from Japan and of E. laidlawi from the type locality near Sonada in India, and determined to be more similar to the Himalayan species. Selected diagnostic features are illustrated for all known Epiophlebia larvae in Figs. 1-3.

# 2. *Epiophlebia diana* sp.n.

Fig. 3A–E

**Material.** Holotype: ultimate female larva in ethanol, China, mountains of western Sichuan, Dr. "D. C. Graham" collector; deposited in Cornell University Insect Collection. Additional material: one penultimate male larva (collected with holotype). Adult unknown.

**Etymology.** Diana – an allusion to the highland huntress and protectress of humankind, who if despoiled, will presage the destruction of mankind by their own handiwork, just as Actaeon was devoured by his own hunting dogs.

**Description (ultimate female larva, holotype).** Compactly built, somewhat limpet-shaped with granulate cuticle and head and thorax rather austropetaliidlike. Total length 26.5 mm, abdomen including terminalia 15.5 mm, hind femur 6.2 mm, prementum 5.7 mm.

Ground color light brown with abdominal pattern in darker brown. Head wider than prothorax and slightly narrower than synthorax, labrum wider than frons, slightly scooplike with shallow median indentation; genae produced forward well beyond anterior margin of compound eyes and ventrally formed into a granule lined ridge along prementum; frontoclypeal region with low transverse ridge; antennae short, ca. 1.8 mm long, and tightly appressed to face, scape and pedicel together as long as first flagellar segment (Fig. 3A); prementum ca.  $2 \times$  as long as wide at base (Fig. 3C), distally triangularly produced to hair fringed edge with deep median tightly appressed cleft, lateral margins strongly sinuate with dorsally directed flange in basal half; labial palps with movable hook slightly shorter than distance from base of movable hook to base of palp, palps rounded-truncate apically with ca. 15 triangular teeth along inner margin; labium in resting position extended into a pocket along anterior edge of mesosternum; occipital lobes well-developed, produced into anterolateral recesses of pronotum.

Anterolateral angle of prothoracic tergum blunt, ventral paracoxal lobes of mesothorax blunt; femora strongly flattened, forefemur ca.  $2.5 \times$  as long as wide (Fig. 3B), foretibia with inner apical brush of hairlike setae; wing-cases parallel, extended posteriorly as far as hind margin of abdominal segment 4; tarsi three jointed.

Abdomen similar to that of Lanthus in outline (Fig. 3D), widest on segment 7, abdominal terga 3 to 10 with low medial ridge; proventriculus 16-lobed: 8 minor folds with 2-3 teeth alternating with 8 major folds with clusters of 4-8 teeth, minor lobes typically alternating with teeth separated and together, major ventral lobes with 5-8 teeth and major dorsal lobes with 4-5teeth; abdominal stridulatory file of segment 3 vestigial and surrounded by smooth area ca. 1/3 length of segment, that of segment 4 ca. 1/2 as high as long, and that of segment 7 ca. 3/4 length of segment (Fig. 3E); lateral abdominal lobes on segment 7 vestigial, moderately developed on 8, and well developed on 9 (Fig. 3D), lateral ridges of segments 7-9 with small irregular spines; paraprocts 1 mm long with hair lined inner surface, epiproct ca.  $0.6 \times$  length of paraprocts, male epiproctal processes slightly divergent, cerci ca.  $1/2 \times \text{length of epiproct.}$ 

**Description (penultimate male larva).** Same as holotype, with the following exceptions: Total length 21.1 mm, abdomen including terminalia 11.9 mm, hind femur 5.4 mm, prementum 4.5 mm including palps. Femora relatively wider than in ultimate instar, wing pads extended to near middle of abdominal segment 2.

**Material used for comparison.** *Epiophlebia superstes*: male and female ultimate and penultimate larvae, Shimizu Japan, 21-XII-1993, H. Karube collector; *Epiophlebia laidlawi*: male and female ultimate and penultimate larvae, Sonada India, 14-IV-1988, G. Patel collector.

# 3. Key to known larvae and adults of Epiophlebiidae

1 Adult: body black with labrum, postclypeus, antefrons, and interocellar ridge largely yellow, with long curved dorsal mesanepisternal stripes, with metepisternal yellow stripes extended below spiracles, with well developed ventral metepimeral stripes, and with well developed ventral, lateral, and middorsal abdomi-



Figs. 1–3. Larvae of *Epiophlebia* species. 1: *Epiophlebia* (*Epiophlebia*) superstes. 2: *Epiophlebia* (*Rheoepiophlebia*) laidlawi.
3: *Epiophlebia* (*Rheoepiophlebia*) diana sp.n. In each figure (A) dorsal view of right antenna, (B) dorsal view of right prothoracic femur, (C) ventral view of prementum, (D) ventral view of abdominal segments 4–10, and (E) lateral view of abdominal segments 3–8 showing stridulatory surface.

nal yellow markings; forewing postnodals typically 12-15, costal side of quadrilateral subequal in length to arcular brace, abdomen 37-42 mm, posterior hamuli with apical lobe shorter than wide, male tergum 10 without middorsal longitudinal carina, male epiproct extended beyond apex of cerci and with ventral rami subacute.

– Larva: cuticle with larger granulations dark brown; antennal pedicel ca.  $2.0 \times$  as long as wide, longer than first flagellar segment (Fig. 1A); prementum ca.  $2.4 \times$ as long as basal width (Fig. 1C); anterolateral angle of prothoracic tergum and ventral paracoxal lobes of mesothorax acute; femora not strongly flattened, forefemur ca.  $3.4 \times$  as long as wide (Fig. 1B), foretibial brush with palpiform setae; abdominal stridulatory file of segment 3 well developed, that of segment 4 ca. as high as long, and that of segment 7 vestigial (Fig. 1E); abdomen with lateral abdominal lobes 7 and 8 well developed (Fig. 1D), male epiproctal processes strongly divergent

..... Epiophlebia (Epiophlebia) – 2

**Type-species:** *Epiophlebia superstes* (Selys), originally named as *Palaeophlebia superstes* Selys (CALVERT 1903).

Larva of E. sinensis Li & Nel not yet known.

1' Adult: body brown with straight dorsal mesanepisternal and short metepisternal yellow stripes, and with vestigial ventral metepimeral stripes and vestigial middorsal abdominal spots, forewing postnodals typically 8-11, costal side of quadrilateral longer than arcular brace, abdomen 45-47 mm, posterior hamuli with apical lobe longer than wide, male tergum 10 with middorsal longitudinal carina, male epiproct not extended beyond apex of cerci and with ventral rami blunt.

– Larva: cuticle with granulations light brown; antennal pedicel ca.  $1.5 \times$  as long as wide, shorter than first flagellar segment (Figs. 2A, 3A); prementum ca.  $2.0 \times$ as long as basal width (Figs. 2C, 3C); anterolateral angle of prothoracic tergum and ventral paracoxal lobes of mesothorax blunt; femora strongly flattened, forefemur  $2.5-3.0 \times$  as long as wide (Figs. 2B, 3B), foretibial brush with hairlike setae; abdominal stridulatory file of segment 3 vestigial and surrounded by triangular smooth area, that of segment 4 ca. 1/2 as high as long, and that of segment 7 well developed (Figs. 2E, 3E); abdomen with lateral abdominal lobes 7 and 8 vestigial (Fig. 2D, 3D), male epiproctal processes slightly divergent

**Type-species:** *Epiophlebia laidlawi* Tillyard. Adult of *Epiophlebia diana* sp.n. not yet known.

2 Adult: abdominal segment 6 laterally beyond antecostal suture and segments 7-10 and terminalia dark brown to black with dorsal yellow spots of segments 8-10 distinct

..... Epiophlebia (Epiophlebia) superstes (Selys)

**Distribution:** Japan: Honshu, Hokkaido, Kyushu, Shikoku.

Flight season: 3-IV-1990, Shikoku (H. Karube collector) to July, Hokkaido (Asahina 1950).

2' Adult: abdominal segment 6 laterally beyond antecostal suture and segments 7-10 and terminalia ferruginous, with dorsal pale spots of segments 8-10 obscure

..... Epiophlebia (Epiophlebia) sinensis Li & Nel

**Distribution:** China: Zheng-nan-gou stream, Heilongjiang province.

**Flight season:** 20–21-VI-2010, Shahezi Zhen (Qiming Li collector).

**3** Larva: labrum with anterior margin entire, scape and pedicel together longer than first flagellar segment (Fig. 2A), prementum with lateral margins slightly sinuous, premental cleft as long as ventral width of palp at midlength (Fig. 2C), forefemur ca.  $3.0 \times$  as long as wide (Fig. 2B), abdominal stridulatory file of segment 7 ca. 1/2 length of segment (Fig. 2E), lateral abdominal lobes slightly protrudent on segment 9 (Fig. 2D)

#### .... Epiophlebia (Rheoepiophlebia) laidlawi Tillyard

**Distribution:** India: Darjeeling; Nepal: Chitre, Shivapuri Hills, Daman, Helambu, Phulping; Bhutan: Thimphu, Taba, Haa, Trongsa (Tillyard 1921; Asahina 1963; Tani & Miyatake 1979; Saville 1990; Sharma & Ofenböck 1996; Nesemann et al. 2008; Brockhaus & Hartmann 2009).

Flight season: 2-V-1979, Shivapuri Hills (O. Tominaga & N. Doi collectors) to 1-VII-1963, Chitre (T. Haruta & A. Hara collectors).

**3'** Larva: labrum with anterior margin widely indented, scape and pedicel together as long as first flagellar segment (Fig. 3A), prementum with lateral margins strongly sinuous, premental cleft longer than ventral width of palp at midlength (Fig. 3C), forefemur ca.  $2.5 \times as$  long as wide (Fig. 3B), abdominal stridulatory file of segment 7 ca. 3/4 length of segment (Fig. 3E), lateral abdominal lobes protrudent on segment 9 (Fig. 3D)

#### ..... Epiophlebia (Rheoepiophlebia) diana sp.n.

**Distribution:** China: mountains of western Sichuan province.

Flight season: unknown.

According to Asahina (1950), upon hatching the shrimp-like prolarva of Epiophlebia superstes is active and able to flip itself a distance of 10 cm. After having reached the water surface the prolarval exuvia is cast and the first instar larva sinks to the stream bed. A total of 14 larval instars were estimated by TOKU-NAGA & ODAGAKI (1939), and it was determined that five to eight years are required for larval development depending on conditions in the larval habitat. During early larval development E. superstes larvae cling to the stones of rushing torrents, with larger instars favoring larger stones in faster current (Asahina 1950). In contrast the larvae of E. laidlawi occur in smaller streams with slower current (TANI & MIYATAKE 1979). However, current velocity for all E. laidlawi sites was rather high according to BROCKHAUS & HARTMANN (2009), and ranged from 15 cm/sec in pools to 200 cm/ sec in riffles. The distinct aquatic habitat requirements of these species from the two subgenera apparently mirror differences in larval morphology. The greater development of the ventral occipital ridges and anteroventral mesothoracic lobes in *E. superstes* apparently is required to more firmly support the labium in faster currents, where the labium is necessarily extended and retracted during feeding. However, the femora of both Rheoepiophlebia species are more strongly flattened than in *E. superstes*, and in the recessed position overlap the trochanter of the following leg to form a sort of suction disk. This morphology may function to prevent these larvae from being washed from their small upstream habitats during periods of high flow. In contrast, the legs of E. superstes are narrower and in the recessed position form a small streamlined disk apparently adapted for moving in faster currents. In addition, FURUKAWA (1934) supposed that the ventral setae of the paraprocts function as adhesive organs, but these setae (found in all known Epiophlebia larvae) point outward about the respiratory opening, and may instead act as a filtering device for water entering the rectal gill chamber.

Epiophlebian larval behavior during the ultimate instar is unique when compared to other Odonata. For *E. superstes* it is known that emergence is preceded by a period of up to five months during which the larvae leave the water and crawl about in the terrestrial environment near the parent stream (TOKUNAGA & ODAGAKI 1939). This unusual shift in larval habitat is also suspected in *E. laidlawi* because TANI & MIYATAKE (1979) were not able to find last instar larvae in the Shivapuri stream. In my experience larval stridulation is yet another unique feature of Epiophlebiidae among Odonata. It is not known whether larvae utilize stridulation to synchronize emergence, optimize habitat occupancy, deter predation, or some other function such as flushing aquatic prey. In any case larvae do stridulate when disturbed, which suggests a protective or warning function. The stridulation sound can be artificially produced by holding the apices of the femora against the lateral abdominal files while bending the abdomen from side to side. The abdominal scraper is apparently formed from cuticle granulations that have been modified into transverse vertical ridges (Figs. 1E, 2E, 3E). The smooth areas about the anterior files apparently serve to enable a stress-free start to the stridulatory movement.

Transformation into the adult generally occurs in the morning on a streamside rock or stem with the larval body in a vertical position, but Asahina (1950) found exuvia on the bark of *Cryptomeria japonica* as far as ten meters from the larval habitat. As is usual for stream inhabiting Odonata, tenerals show the greatest dispersal, with mature individuals eventually aggregating back toward the parent stream or to nearby streams. For example, a teneral female *E. laidlawi* has been collected at the top of the Shivapuri Hills about one km from the parent stream (TANI & MIYATAKE 1979), although mature adults are rarely collected far from suitable larval habitat.

Flight of Epiophlebia is swift and relatively efficient, but maneuverability and acceleration are far less than in Zygoptera or Anisoptera; and after a fifteen minute warm up period, flight can occur at 16°C, or even in the rain at lower temperatures than typical for extant Odonata (RÜPPELL & HILFERT 1993). Feeding flight occurs in the valley of the parent stream near treetop level, where early in the season small insects are fed on throughout the day. As the season progresses, feeding is limited to before 10:00 and to after 15:00 (Asahina 1950). The male search flight for females occurs during mid-day and is typified by a slow upstream flight close to the water surface, with the males weaving forward and back while inspecting suitable oviposition sites for mature females. Males pursue females as they fly between oviposition sites and grasp females with the typical anisopteran tandem hold. Mating is at rest and the huge hooked and studded apex of the posterior hamuli (penis) may not only function in sperm displacement, removal, and transfer, but could also damage the female reproductive tract and thus reduce the efficiency of future mating. Unlike other Odonata the male head of Epiophlebia has posterior anti-tandem lobes (CARLE 1996), which presumably prevent multi-tandem entanglements as sometimes observed in Lestidae.

Flight season is about a month in length (ASAHINA 1950), and varies significantly with latitude and elevation: for *E. superstes* it extends from late March in Kyusyu to July in Hokkaido, and for *E. laidlawi* it extends from early May until early July. *Epiophlebia superstes* and *E. laidlawi* (and thus perhaps the sub-

genera of *Epiophlebia*) apparently also differ in the substrate utilized for oviposition. E. superstes has been reported to oviposit in a wide variety of soft tissued plants, although oviposition is sometimes attempted in hard tissue. Chosen plants typically grow in wet and mostly shaded areas, less than one meter from the stream, and include several genera of spermatophytes: Houttuynia (Saururaceae), Angelica (Apiaceae), Elatostema (Urticaceae), Trautvetteria (Ranunculaceae), Saxifraga (Saxifragaceae), Parasenecio, Petasites, Ligularia (Asteraceae), Lilium, Cardiocrinum (Liliaceae), Arisaema (Araceae), and Wasabia, Cardamine (Brassicaceae); and a few bryophytes: Dumortiera (Dumortieraceae), Conocephalum (Conocephalaceae), and Pellia (Pelliaceae) (Asahina 1934, 1950; Asahina & Eda 1958, 1982; Asahina & Sugimura 1981; Tami-YA & MIYAKAWA 1984; TOKUNAGA & ODAGAKI 1939). However, suitable plants for oviposition are not known at the small streams in the Shivapuri Hills, one of the few localities where both adults and larvae of E. laidlawi have been found (TANI & MIYATAKE 1979). It is possible that suitable oviposition substrates may also include decaying logs, but ASAHINA (1982) has identified Elatostema hookerianum (Urticaceae) as the plant most likely utilized for oviposition in this region. Female Epiophlebia oviposit unaccompanied, and require a small opening of about one cubic meter above the oviposition site for access, possibly because of crude flight capabilities reported by RÜPPELL & HILFERT (1993). Oviposition in plant stems is preceded by trial thrusts, which, if successful, lead to a descent along the stem, after which eggs are laid from the bottom of the stem upward. ASAHINA (1934) reported that females would fly away without ovipositing when plants were found to be too solid. Similar plants are utilized by austropetaliids for oviposition (CARLE 1996), but the more weakly sclerotized epiophlebian abdomen apparently makes the process more time consuming and seemingly laborious. The eggs of each female are arranged in a sinuous row of up to about 1000 eggs maximum, with an egg deposited about every four seconds (Asahina 1950). The eggs are inserted with the anterior coarse end upward, are about 1/4 as wide as long, and average about 1.1 mm in length. Micropyles number 10-14 (reported as 12-14 by ANDO 1962) and are arranged about the anterior end in a subapical ring. Embryonic development is ca. 30 days at 20°C and 20 days at 25.5° C (Asahina 1950).

# 5. Epiophlebian biogeography

Lower Jurassic fossils of the epiophlebian sister group indicate that *Epiophlebia* are the survivors of an evo-

lutionary line that has been genetically isolated for at least 180 million years. Unfortunately, conclusive fossils are not known of Epiophlebia or of its Jurassic stem group. Several synapomorphies, including the widening of the MA-MP interspace, separate both Epiophlebia and its sister Anisopteroids from their Jurassic stem group (CARLE 1982a), which is currently known only from the Palaearctic (HAGEN 1862, 1866; BODE 1905; MARTYNOV 1925; PRITYKINA 1968, 1970, 1977, 1985; Whalley 1985; Fleck & Nel 2002; Nel et al. 2009). In turn, Epiophlebia is distinguished from its sister group Anisopteroids by the lack of several wing features related to increased vagility (CARLE 1982a). It is telling that even the most plesiomorphic of these more vagile Anisopteroids had already spread across Pangaea during the Jurassic, as evidenced by fossils from Antarctica to China (CARPENTER 1969; HANDLIRSCH 1920; PRITYKINA 1980; HONG 1983). Epiophlebia's apparent failure to disperse more widely indicates that Epiophlebia had already specialized to inhabit isolated east Palaearctic highland streams during the early Jurassic. Lack of wing features including a discal nodus, a well developed nodal hinge, or an expanded hind wing anal field (CARLE 1982a) indicate a lower vagility of Epiophlebia when compared to its anisopteran contemporaries. The most plesiotypic extant anisopterans are the Austropetaliidae, which inhabit the mountain streams and spring seeps of the Southern Hemisphere (CARLE & LOUTON 1994; CARLE 1996; CARLE et al. 2008), suggesting that these southern habitats remained unoccupied until the basal radiation and dispersal of Anisoptera. Specilization for highland streams and associated low vagility has also likely prevented dispersal of *Epiophlebia* into the mountains of Europe or its dispersal across the North American/Okhotsk plate (SENO et al. 1996) from Japan to America. In contrast, nearby ecological associates of *Epiophlebia*, the anisopteran genera *Lanthus* Needham, 1897 (Gomphidae) and Tanypteryx Kennedy, 1917 (Petaluridae), occur exclusively in Japan and North America (CARLE 1980, 1986).

The known distribution of the subgenus *Epiophlebia* s.str. includes Japan and northeastern China, and the known distribution of *Rheoepiophlebia* subgen.n. includes the Himalayas and mountains of south-central China. Similar adult morphology of the northern species pair (*E. superstes* and *E. sinensis*) suggests that a vicariance event isolating them on either side of the Sea of Japan may be related to Pleistocene glaciation. In contrast, larvae of the southern species pair are easily distinguished compared to the difficulty often encountered when identifying dragonfly larvae to species. This indicates a much more ancient vicariance event, perhaps the formation of deep river valleys between the Himalayas and the mountains of southcentral China.

Although the utility of epiophlebian autapomorphic features remains largely unknown, it is apparent that the unusual wings of epiophlebians are adapted for flight about cool high elevation habitats (RUPPELL & HILFERT 1993). In addition, a streamlined larval body and a rheophilic lifestyle also enable Epiophlebia to exploit the stable environment and detritus based ecosystems of high elevation spring-fed seeps and streams. Unfortunately, this high energy environment is not conducive to fossil formation, thus explaining the current lack of *Epiophlebia* from the fossil record. As streams descend mountain valleys ambient temperature rises, gradient decreases and sediment accumulates in pool areas (VANNOTE et al. 1980), thus forming the habitat utilized by the borrowing larvae of gomphid, cordulegastrid and chlorogomphid populations (CARLE 1983b). These sentinels patrol along the upper limit of lowland aquatic ecosystems dominated by advanced anisopteran competitors which together help restrict Epiophlebia to its mountain retreats.

Fortunately for Epiophlebia its habitat is an ideal shelter from the perturbations of global extinction events such as meteor impacts and extreme volcanism, which result in extended periods of reduced sunlight (ALVAREZ et al. 1980; KNOLL et al. 2007). The environmental stability afforded small stream communities from emerging ground water and a persistent flow of energy and nutrients from detritus based ecosystems (CARLE 1982c) is strongly contrasted by temporary lentic ecosystems, which although often highly productive are dependent on reliable sunlight and rainfall. For example, extinction of all Stenophlebioidea Pritykina, 1980 and Aeschnidioidea Carle & Wighton, 1990 is associated with large size, a stalking larval feeding mode, and an apparent lentic habitat preference (CARLE & WIGHTON 1990; ZHANG 1999). A synergism favoring an exposed feeding mode and the utilization of relatively predator free temporary aquatic habitats would explain the lack of concealed larvae among fossil Stenophlebioidea and Aeschnidioidea, but would eventually also expose these giant dragonflies to the environmental perturbations associated with major extinction events. In contrast, the diversification of Anisoptera and the prolonged survival of Epiophlebiidae are associated with small size, secretive larval feeding modes, the environmental stability of emerging ground water, and the persistent flow of energy and nutrients through detritus based ecosystems.

### 6. Acknowledgments

I thank the following individuals and institutions for the loan of material: Mrs. Leonora K. Gloyd, E.B. Williamson Collection;

Dr M.J.Westfall, Jr., Florida State Collection of Arthropods and International Odonata Research Institute; Dr O.S.Flint, Jr., United States Museum of Natural History; Dr A. Newton and Dr S. Cover, Museum of Comparative Zoology, Harvard University; Dr. Jon Gelhaus, Academy of Natural Sciences; Dr. D.A. Grimaldi, American Museum of Natural History; W.J. Winstanley, Dr. Q.D. Wheeler, Cornell University Insect Collection; Mr. Brooks and Dr D.T. Goodger, The Natural History Museum; and Mr. Haruki Karube. I am also grateful to E.B. Cosgrove for assistance with larval photography. New Jersey Agricultural Experiment Station publication D-08022-12-97, supported by state funds.

# 7. References

- ALVAREZ L.W., ALVAREZ W., ASARO F., MICHEL H.V. 1980. Extraterrestrial cause for the Cretaceous-Tertiary extinction. – Science 208(4448): 1095–1108.
- ANDO H. 1962. The comparative embryology of Odonata with special reference to a relic dragonfly *Epiophlebia superstes* Selys. – The Japan Society for the Promotion of Science, Tokyo.
- ASAHINA S. 1934. Ecological observations on *Epiophlebia superstes*. Kontyû **8**(2): 103–106.
- ASAHINA S. 1949. On some archaic structures retained in *Epio-phlebia superstes* (Odonata, Anisozygoptera). Mushi 19: 49–51.
- ASAHINA S. 1950. On the life history of *Ephiophlebia superstes* (Odonata, Anisozygoptera). – Proceedings of the VIII International Congress of Entomology, Stockholm, pp. 337– 341.
- Asahina S. 1954. A morphological study of a relic dragonfly *Epiophlebia superstes* Selys (Odonata, Anisozygoptera). The Japan Society for the Promotion of Science, Tokyo, 153 pp.
- ASAHINA S. 1963. Description of the possible adult dragonfly of *Epiophlebia laidlawi* from the Himalayas. Tombo **6**(3/4): 18–20.
- AsaHINA S. 1982. Survey of the relict dragonfly *Epiophlebia laidlawi* Tillyard in Nepal, May 1981. – Reports of the Odonata Specialist Group of the International Union for the Conservation of Nature, No. 1, 10 pp.
- ASAHINA S., EDA S. 1958. A female *Epiophlebia superstes* laying eggs (Odonata, Epiophlebiidae). – Kontyû 26: 1.
- ASAHINA S., EDA S. 1982. Further observations on bryophyte oviposition by *Epiophlebia superstes*. – Tombo 25: 2–6.
- ASAHINA S., SUGIMURA M. 1981. Oviposition of *Epiophlebia superstes* into bryophytes. Tombo **24**(1): 22–23.
- BECHLY G. 1995. Morphologische Untersuchungen am Flügelgeäder der rezenten Libellen und deren Stammgruppenvertreter (Insecta; Odonatoptera), unter besonderer Berücksichtigung der Phylogenetischen Systematik und des Grundplanes der \*Odonata. – Petalura spec. vol. 1: 1–341; Böblingen.

- BECHLY G. 1996. Morphologische Untersuchungen am Flügelgeäder der rezenten Libellen und deren Stammgruppenvertreter (Insecta; Pterygota; Odonata), unter besonderer Berücksichtigung der Phylogenetischen Systematik und des Grundplanes der \*Odonata. – Petalura spec. vol. 2: 1–402; Böblingen.
- BODE A. 1905. Orthoptera und Neuroptera aus dem oberen Lias von Braunschweig. – Jahrbuch der Preussischen Geologischen Landesanstalt 25: 218–245.
- BRAUCKMANN C., ZESSIN W. 1989. Neue Meganeuridae aus dem Namurium von Hagen-Vorhalle (BRD) und die Phylogenie der Meganisoptera (Insecta, Odonata). – Deutsche Entomologische Zeitschrift 36(1–3): 177–215.
- BROCKHAUS T., HARTMANN A. 2009. New records of *Epiophlebia laidlawi* Tillyard in Bhutan, with notes on its biology, ecology, distribution, zoogeography and threat status (Anisozygoptera: Epiophlebiidae). – Odonatologica **38**(3): 203– 215.
- CALVERT P.P. 1903. "A genealogic study of dragonfly wing venation" by J.G. Needham. – Entomological News 14: 208.
- CARLE F.L. 1980. A new *Lanthus* (Odonata: Gomphidae) from eastern North America with adult and nymphal keys to American octogomphines. – Annals of the Entomological Society of America. **73**: 172–179.
- CARLE F.L. 1982a. The wing vein homologies and phylogeny of the Odonata: A continuing debate. – Society of International Odonatologists Rapid Communications 4: 1–66.
- CARLE F.L. 1982b. Evolution of the odonate copulatory process. Odonatologica **11**(4): 271–286.
- CARLE F.L. 1982c. Thoughts on the Origin of Insect Flight. Entomological News **93**(5): 159–172.
- CARLE F.L. 1983a. The first collection of larval *Epiophlebia superstes* (Selys) (Anisozygoptera: Epiophlebiidae). Tombo **26**(1/4): 16.
- CARLE F.L. 1983b. A new Zoraena (Odonata: Cordulegastridae) from eastern North America with a key to adult American Cordulegastridae. – Annals of the Entomological Society of America 76: 61–68.
- CARLE F.L. 1986. The classification, phylogeny, and biogeography of the Gomphidae (Anisoptera), 1. Classification. – Odonatologica 15(3): 275–326.
- CARLE F.L. 1995. Evolution, taxonomy, and biogeography of ancient gondwanian libelluloides, with comments on anisopteroid evolution and phylogenetic systematics (Anisoptera: Libelluloidea). – Odonatologica 24(4): 383–424.
- CARLE F.L. 1996. Revision of Austropetaliidae (Anisoptera: Aeshnoidea). – Odonatologica 25(3): 231–259.
- CARLE F.L., KJER K.M. 2002. Phylogeny of *Libellula* Linnaeus (Odonata: Insecta). – Zootaxa 87: 1–18.
- CARLE F.L., KJER K.M., MAY M.L. 2008. Evolution of Odonata, with special reference to Coenagrionoidea (Zygoptera). – Arthropod Systematics & Phylogeny 66(1): 37–44.
- CARLE F.L., LOUTON J.A. 1994. The larva of *Neopetalia punctata* and establishment of Austropetaliidae fam. nov. (Odonata). – Proceedings of the Entomological Society of Washington **96**(1): 147–155.

- CARLE F.L., WIGHTON D.C. 1990. Odonata. Pp. 51–68 in: GRI-MALDI D.A. (ed.), Insects from the Santana Formation, Lower Cretaceous, of Brazil. – Bulletin of the American Museum of Natural History, Number 195. New York. 191 pp.
- CARPENTER F.M. 1969. Fossil insects from Antarctica. Psyche 76(4): 418–425.
- CARPENTER F.M. 1992. Superclass Hexapoda. In: MOORE R.C., KAESLER R.L. (eds), Treatise on Invertebrate Paleontology, Arthropoda. – Geological Society of America & University of Kansas **4**: 3–4. xxii + 655 pp. Boulder & Lawrence.
- FLECK G., BECHLY G., MARTÍNEZ-DELCLÓS X., JARZEMBOWSKI E. A., NEL A. 2004. A revision of the Upper Jurassic-Lower Cretaceous dragonfly family Tarsophlebiidae, with a discussion on the phylogenetic positions of the Tarsophlebiidae and Sieblosiidae (Insecta, Odonatoptera, Panodonata). – Geodiversitas 26(1): 33–59.
- FLECK G., NEL A. 2002. The first isophlebioid dragonfly (Odonata: Isophlebioptera: Campterophlebiidae) from the Mesozoic of China. – Palaeontology 45(6): 1123–1136.
- FURUKAWA H. 1934. On the nymph of a dragonfly, *Epiophlebia* superstes Selys. – Annotationes Zoologicae Japonenses 14: 451–456.
- HAGEN H.A. 1862. Über die Neuropteren aus dem Lithographischen Schiefer in Bayern. Pars I. – Palaeontographica 10: 96–145.
- HAGEN H.A. 1866. Die Neuropteren des Lithographischen Schiefers in Bayern. Pars I. – Palaeontographica **15**: 57–96.
- HANDLIRSCH A. 1906–1908. Die Fossilen Insekten und die Phylogenie der Rezenten Formen. Ein Handbuch fuer Palaeontologen und Zoologen. – Engelmann, Leipzig, 640 pp.
- HANDLIRSCH A. 1920. Palaeontologie. Systematische Übersicht. Pp. 117–299 in: SCHRÖDER C. (ed.), Handbuch der Entomologie, vol. 3.
- HANDLIRSCH A. 1939. Neue Untersuchungen über die fossilen Insekten, Teil II. – Annalen des Naturhistorischen Museums Wien 49: 1–240.
- HONG Y.C. 1983. Middle Jurassic Fossil Insects in North China. – Geological Publication House, Beijing, 223 pp.
- KLASS K.-D. 2008. The female abdomen of ovipositor-bearing Odonata (Insecta: Pterygota). – Arthropod Systematics & Phylogeny 66: 45–142.
- KNOLL A.H., BAMBACH R.K., PAYNE J.L., PRUSS S., FISCHER W.W. 2007. Paleophysiology and end-Permian mass extinction. – Earth and Planetary Science Letters 256(3–4): 295–313.
- Lew G.T. 1933. Head characters of the Odonata with special reference to the development of the compound eyes. Entomologica Americana 14(2): 41–96.
- LI J., NEL A., ZHANG X., FLECK G., GAO M., LIN L., ZHOU J. 2012. A third species of the relict family Epiophlebiidae discovered in China (Odonata: Epiproctophora). – Systematic Entomology 37(2): 408–412.
- MARTYNOV A.V. 1925. To the knowledge of fossil insects from Jurassic beds in Turkestan. 2. Raphidioptera (continued), Orthoptera (s. l. ), Odonata, Neuroptera. – Izvestiya Akademii Nauk 19(12–15): 569–598.

- NEEDHAM J.G. 1903. A genealogic study of the dragonfly wing venation. Proceedings of the United States National Museum **26**: 703–764.
- NEEDHAM J.G. 1930. A manual of the dragonflies of China A monographic study of Chinese Odonata. – Zoologia Sinica (Series A. Invertebrates of China Volume II, Fascicle 1), Peking, 399 pp.
- NEL A., MARTÍNEZ-DELCLOS X., PLACHER J.C., HENROTAY M. 1993. Les "Anisozygoptera" fossils – Phylogénie et classification (Odonata). – Martinia Numéro Hors-Série 3: 1–311.
- NEL A., BECHLY G., DELCLOS X., HUANG D. 2009. New and poorly known Mesozoic damsel-dragonflies (Odonata: Isophlebioidea: Campterophlebiidae, Isophlebiidae). – Palaeodiversity 2: 209–232.
- NESEMANN H., TACHAMO R.D., SHAH D.N. 2008. Records of Epiophlebiidae (relict Himalayas dragonfly) habitats from central Napal. – Abstracts of Papers of the 18th International Symposium of Odonatology, Nagpur, p. 44.
- PFAU H.K. 1971. Struktur und Funktion des sekundären Kopulationsapparates der Odonaten (Insecta, Palaeoptera), ihre Wandlung in der Stammesgeschichte und Bedeutung für die adaptive Entfaltung der Ordnung. – Zeitschrift für Morphologie und Ökologie der Tiere **70**: 281–371.
- PRITYKINA L.N. 1968. Dragonflies (Odonata) from Karatau. Pp. 26–55 in: ROHDENDORF B.B. (ed.), Yurskie nasekomye Karatau. Nauka, Moscow. 249 pp.
- PRITYKINA L.N. 1970. Triassic and Jurassic Dragonflies of the Liassophlebiidae from Soviet Central Asia. – Paleontological Journal 4(1): 91–102.
- PRITYKINA L.N. 1977. New dragonflies from Lower Cretaceous deposits of Transbaikalia and Mongolia. – Transactions of the Soviet-Mongolian Palaeontological Expedition, Moscow 4: 81–96.
- PRITYKINA L.N. 1980. Odonata. In: ROHDENDORF B.B., Historical Development of the Class Insecta. – Trudy Paleontological Institute, Moscow 175: 128–134.
- PRITYKINA L.N. 1985. Fossil Odonata (Libellulida = Odonata) from Siberia and outer Mongolia. Pp. 5–42 in: RASNITSYN A.P., Fossil Insects from Siberia and Mongolia. – Trudy Paleontological Institute, Moscow **211**: 1–191.
- REHN A.C. 2003. Phylogenetic analysis of higher level relationships of Odonata. – Systematic Entomology 28: 181–239.
- RIEK E.F., KUKALOVÁ-PECK J. 1984. A new interpretation of dragonfly wing venation based upon Early Upper Carboniferous fossils from Argentina (Insecta, Odonatoidea) and basic character states in pterygote wings. – Canadian Journal of Zoology 62: 1150–1166.
- RUPPELL G., HILFERT D. 1993. The flight of the relict dragonfly *Epiophlebia superstes* (Selys) in comparison with that of the modern Odonata (Anisozygoptera: Epiophlebiidae). – Odonatologica 22(3): 295–309.
- SAVILLE N. 1990. Survey of the status of *Epiophlebia laidlawi* (Tillyard) in the Shivapuri Watershed and Wildlife Reserve, 1988. – Report on the Cambridge Entomological Expedition to Nepal, pp. 31–35.

- SCHMIDT E. 1915. Vergleichende Morphologie des 2. und 3. Abdominalsegmentes bei männlichen Libellen. – Zoologische Jahrbücher **39**: 87–196.
- SELVS LONGCHAMPS E. de 1889. Palaeophlebia, nouvelle légion de Caloptérygines, suivi de la description d'une nouvelle Gomphine du Japon: Tachopteryx pryeri. – Comptes Rendus de la Société Entomologique de Belgique 3: 153–154.
- SENO T., SAKURAI T., STEIN S. 1996. Can the Okhotsk plate be discriminated from the North American plate? – Journal of Geophysical Research 101(B5): 11305–11315.
- SHARMA S., OFENBÖCK T. 1996. New discoveries of *Epiophle-bia laidlawi* Tillyard, 1921 in the Nepal Himalaya (Odonata, Anisozygoptera: Epiophlebiidae). – Opuscula Zoologica Fluminensia **150**: 1–11.
- TAMIYA Y., MIYAKAWA K. 1984. On the oviposition habitat of *Epiophlebia superstes* (Selys) (Anisozygoptera: Epiophlebiidae). – Odonatologica 13(2): 301–307.
- TANI K., MIYATAKE Y. 1979. The discovery of *Epiophlebia laidlawi* Tillyard, 1921 in the Kathmandu valley, Nepal (Anisozygoptera: Epiophlebiidae). – Odonatologica 8(4): 329–332.
- TILLYARD R.J. 1914. On some problems concerning the development of the wing venation of the Odonata. – Proceedings of the Linnaean Society New South Wales **39**(1): 163–216.
- TILLYARD R.J. 1917. The Biology of Dragonflies. Cambridge University Press, Cambridge. xii + 396 pp.
- TILLYARD R.J. 1921. On an anisozygopterous larva from the Himalayas (Order Odonata). – Records of the Indian Museum 22: 93–107.
- TILLYARD R.J. 1925. The British Liassic Dragonflies (Odonata). – London, British Museum of Natural History, Fossil Insects 1: 38 pp.
- TILLYARD R.J. 1929. The evolution of the order Odonata. Fourth International Congress of Entomology, Ithaca **1928**: 543–545.
- TILLYARD R.J., FRASER F.C. 1940. A reclassification of the order Odonata, based on some new interpretations of the venation of the dragonfly wing. Parts I, II, III. – Australian Zoologist 9(2): 125–169, 9(3): 195–221, 9(4): 359–396.
- TOKUNAGA M., ODAGAKI S. 1939. Ecological and morphological studies on a dragonfly *Epiophlebia superstes* Selys. – Celebration Bulletin of Professor Sadao Yoshida, pp. 472–496. [in Japanese]
- VANNOTE R.L., MINSHALL G.W., CUMMINS K.W., SEDELL J.R., CUSHING C.E. 1980. The river continuum concept. – Canadian Journal of Fisheries and Aquatic Sciences 37(1): 130–137.
- WHALLEY P.E.S. 1985. The systematics and palaeogeography of the Lower Jurassic insects of Dorset, England. – Bulletin of the British Museum (Natural History), Geology, **39**: 107– 187.
- ZHANG J. 1999. Aeschnidiid nymphs from the Jehol biota (latest Jurassic–Early Cretaceous), China, with a discussion of the family Aeschnidiidae (Insecta, Odonata). – Cretaceous Research 20: 813–627.

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Arthropod Systematics and Phylogeny

Jahr/Year: 2012

Band/Volume: 70

Autor(en)/Author(s): Carle Frank Louis

Artikel/Article: <u>A new Epiophlebia (Odonata: Epiophlebioidea) from China with a</u> review of epiophlebian taxonomy, life history, and biogeography 75-83