PENSOFT

SENCKENBERG world of biodiversity



An integrative revision of the subgenus *Liophloeodes* (Coleoptera: Curculionidae: Entiminae: Polydrusini): taxonomic, systematic, biogeographic and evolutionary insights

Beniamin Wacławik¹, Francesco Nugnes², Umberto Bernardo², Marco Gebiola³, Maja Przybycień¹, Dorota Lachowska-Cierlik¹

1 Institute of Zoology and Biomedical Research, Jagiellonian University, Kraków, Poland.

2 CNR, Institute for Sustainable Plant Protection, SS of Portici, Portici (NA), Italy.

3 Department of Entomology, University of California, Riverside, CA, USA.

http://zoobank.org/D8457AE0-64E6-4346-B9ED-57D1C7B4392F

Corresponding author: Dorota Lachowska-Cierlik (dorota.lachowska-cierlik@uj.edu.pl)

Received	2 March 2021		
Accepted	17 June 2021	Academic Editors	Sergio Pérez-González, Mónica M. Solórzano-
Published	11 August 2021		Kraemer

Citation: Wacławik B, Nugnes F, Bernardo U, Gebiola M, Przybycień M, Lachowska-Cierlik D (2021) An integrative revision of the subgenus *Liophloeodes* (Coleoptera: Curculionidae: Entiminae: Polydrusini): taxonomic, systematic, biogeographic and evolutionary insights. Arthropod Systematics & Phylogeny 79: 419–441. https://doi.org/10.3897/asp.79.e64252

Abstract

The subgenus *Liophloeus* Weise, 1894 of *Liophloeus* Germar, 1817 (Coleoptera: Curculionidae: Entiminae: Polydrusini) consists of five morphologically similar species traditionally diagnosed based on the shape of the aedeagus. However, traits of the genital apparatus exhibit substantial and overlapping inter- and intraspecific variation. All five species have the same ecological requirements and occur in central and eastern Europe, mostly in montane areas. The focus of this work was to verify the taxonomic status and validity of *Liophloeodes* species using a combination of molecular and morphometric techniques. Specimens were collected from the entire distribution range and initially assigned to a species according to the aedeagal shape. Genetic diversity and phylogeny of the subgenus were studied using three molecular markers (two ribosomal, 28S-D2 and ITS2, and one mitochondrial, COI). Moreover, several morphological characters were used for multivariate morphometric analyses. Finally, presence and prevalence of bacterial endosymbionts among species were investigated. Phylogenies based on ribosomal markers suggest that traditional species are correctly delimited, whereas COI phylogeny suggests hybridization and introgression occurring between *Liophloeodes* species. Morphometric analyses confirmed low interspecific diversity. Two major bacterial endosymbionts, *Rickettsia* and *Wolbachia*, were detected in many populations. We argue that *Liophloeodes* consists of young lineages whose evolution and diversification was possibly mediated by cyclic climate change events.

Key words

molecular markers, morphometry, phylogeny, taxonomy, weevils.

Introduction

Integrative taxonomy is a relatively new approach based on the idea that results obtained using different methods should be integrated to increase robustness of taxonomic hypotheses (Dayrat 2005; Schlick-Steiner et al. 2010; Yeates et al. 2011), to which a degree of expected stability may be associated (Padial et al. 2010). Over the last 15 years, this integrative approach has become the most popular (and useful) taxonomic method, giving robust, reliable and often unexpected results for many groups of organisms (Miralles et al. 2011; Schutze et al. 2017; Vitecek et al. 2017, Stec et al. 2020a; Stec et al. 2020b). Integrated data include not only those obtained by "traditional" morphological and "modern" molecular methods, but also rigorous statistical testing of detailed morphometric measurements as well as ecological and biogeographical data, and even infection by microorganisms that may affect the organisms' biology (Gebiola et al. 2012).

Weevils (Coleoptera: Curculionoidea) are one of the most diverse groups of living organisms (McKenna et al. 2009) with more than 60,000 known species (of which over 50,000 belong to the family Curculionidae), and many ecological forms that evolved over millions of years of coevolution with (mostly angiosperm) plants (Oberprieler et al. 2007). Due to this huge morphological diversity, weevils represent a big challenge for taxonomists. Traditional taxonomy has been verified by molecular markers and by phylogenomic data at the level of subfamilies (Marvaldi et al. 2002; Shin et al. 2018). However, phylogenies and taxonomies of many tribes and genera are still poorly known and mostly unresolved. Integrative taxonomy as a tool that allows combining morphological knowledge with molecular data has proven helpful for studies that focus on weevils (Grobler et al. 2006; Toševski et al., 2014; Brown 2017).

The genus Liophloeus Germar, 1817 (Coleoptera: Curculionidae) includes two subgenera: Liophloeus sensu stricto and Liophloeodes Weise, 1894 (Fig. 1). Liophloeus s.s. consists of three species comprising both bisexual and parthenogenetic populations, and its range is much wider than Liophloeodes, as it covers the most of Europe (including Scandinavia and British Isles) and has wider ecological requirements (it can occur both in the cold and wet biotopes and in the warmer habitats in the lowlands). All *Liophloeodes* species are exclusively bisexual, and their geographic range overlaps with only one species of Liophloeus s.s., L. tessulatus. Here we examined the morphological diversity and phylogenetic systematics of the subgenus Liophloeodes. Systematics of Liophloeodes has undergone many changes over time because of the extreme morphological similarity that is noted between its taxa, whereas the distinction between this subgenus and Liophloeus s. s. is straightforward, even in the field. The basis for current taxonomy of *Liophloeodes* is the work by Weise (1894), who established a subgenus that included the following group of species: Liophloeus (Liophloeodes) schmidti Boheman, 1842, Liophloeus

(Liophloeodes) lentus Germar, 1824, Liophloeus (Liophloeodes) chrysopterus Boheman, 1842, Liophloeus (Liophloeodes) gibbus Boheman, 1842 and Liophloeus (Liophloeodes) liptoviensis Weise, 1894, which can be distinguished only by the shape of the aedeagus. Later, this nomenclature has been modified a few times (Apfelbeck 1928; Petri 1912; Reitter 1916). Smreczyński (1958) differentiated two species: Liophloeus (Liophloeodes) lentus and Liophloeus (Liophloeodes) pupillatus Apfelbeck, 1928, and at the same time split the first taxon into several distinct subspecies [Liophloeus (Liophloeodes) lentus lentus, Liophloeus (Liophloeodes) lentus gibbus, Liophloeus (Liophloeodes) lentus liptoviensis, Liophloeus (Liophloeodes) lentus herbstii and Liophloeus (Liophloeodes) lentus ovipennis with uncertain status]. More than 20 years later, Dieckmann (1980) elevated Liophloeus (Liophloeodes) lentus subspecies to the species level [except for uncertain Liophloeus (Liophloeodes) lentus ovipennis]. Currently, the subgenus Liophloeodes comprises five nominal species [it includes four Smreczynski's subspecies - except for Liophloeus (Liophloeodes) lentus ovipennis - and Liophloeus (Liophloeodes) pupillatus] whose taxonomy is rather poorly understood, and identification, based mainly on shape of male genitalia (Fig. 2) is considered extremely challenging. Furthermore, many formerly described taxa, despite being later synonymized, can still be found in faunistic surveys and species checklists which underlines even more the need for a taxonomic revision.

All Liophloeodes species prefer wet and cold biotopes and their host plants are species from the families Apiaceae (Aegopodium spp., Chaerophyllum spp., Heracleum spp.), Asteraceae (Petasites spp., Tussilago spp.) and Urticaceae (Urtica spp.). They can be found near streams and rivers in the mountains or sub-mountainous areas in south-eastern Europe (Fig. 3), across the whole Carpathians, eastern Alps, Dinaric Alps, Balkan Mountains, the Sudetes, and small montane chains in Pannonian Basin. Species belonging to this subgenus are partially sympatric. Their ecological and geographical similarity matches the low level of morphological differentiation among species. A general problem with the morphology of Liophloeodes is the lack of reliable diagnostic characters that would allow for confident species identification. The only diagnostic trait is the shape of the aedeagus; hence, females can only be distinguished by association with males from the same population. However, even identifying males can be problematic, due to the high intraspecific and low interspecific phenotypic plasticity of aedeagus. The diversity of Liophloeodes could be described more as a gradient of differences between species (Smreczyński 1958). Another major problem with Liophloeodes taxonomy is the limited knowledge about populations living south of the Pannonian Basin and, more generally, about their biology. The only exception is a study on Microsporidia infecting some Polish populations of Liophloeus (Liophloeodes) lentus (Ovcharenko et al. 2013).



Figure 1. Habitus of Liophloeus (Liophloeodes) lentus male (a) and female (b) and Liophloeus tessulatus female (c). Scale bar: 5 mm.



Figure 2. Aedeagi of *Liophloeus* (*Liophloeodes*) species in dorsal, ventral, and lateral views. A- *Liophloeus* (*Liophloeodes*) herbstii, B – *Liophloeus* (*Liophloeodes*) gibbus, C – *Liophloeus* (*Liophloeodes*) lentus, D – *Liophloeus* (*Liophloeodes*) liptoviensis, E – *Liophloeus* (*Liophloeodes*) pupillatus.

Taking into consideration all mentioned issues concerning these weevils, their systematics can be considered as highly uncertain and should be verified by integrative taxonomy, using morphological, molecular and morphometric data. Using traditional aedeagus shape-based species identification as a starting hypothesis, the diversity of *Liophloeodes* species from the entire known distribution range was iteratively assessed by a combination of morphological and molecular examination. Three molecular (two ribosomal, one mitochondrial) markers were used as distinct lines of evidence. The morphometric measurements were also taken and analysed as another independent method. Additionally, endosymbionts occurrence and phylogeny has been shown to be another potentially important line of evidence to support differences between species (Gebiola et al. 2012). Endosymbiont research



Figure 3. Geographical distribution of the species belonging to the genus *Liophloeus* (*Liophloeodes*) and contact zone between *Liophloeus* (*Liophloeodes*) *lentus* and *Liophloeus* (*Liophloeodes*) gibbus in Dunajec and Poprad valley.

could also deliver additional information about possible paths of evolution, diversification and speciation within the taxon, as symbionts may have an impact on reproduction isolation by causing reproductive manipulations (Shropshire et al. 2020).

Materials and Methods

Material collection

Specimens were collected in 2009–2010 in Poland and Slovakia, and in 2013–2017 in Poland, Slovakia, the Czech Republic, Austria, Slovenia, Croatia, Bosnia and Herzegovina, Serbia, Hungary, Romania, Bulgaria, and Ukraine (Table S1, Fig. 3). Specimens (481 individuals, now deposited in the Department of Entomology, Jagiellonian University) were collected by hand or using a sweeping net, immediately put in 95% alcohol, transferred to the laboratory and stored at -20°C until use. The collection sites were determined using faunistic literature [mainly Smreczyński (1958), but also local faunistic papers] and knowledge about preferred landscape and environment. Most specimens were collected in the valleys, near streams and rivers in wet and cold biotopes, in plant communities, mainly consisting of Apiaceae, but also *Petasites* spp. and *Urtica* spp.

Liophloeus (Liophloeodes) ovipennis, which was described based on a single specimen collected in the French Alps, is probably a misidentified weevil belonging to *Liophloeus s.s.*, because there are no other data about the occurrence of *Liophloeodes* in this part of Europe. Specimens of *Liophloeus tessulatus* occurring in the sampling areas were also collected to be included in phylogenetic analyses, to help understand interspecific phylogenetic relationship in the genus.

Molecular techniques

DNA was isolated from whole insect bodies. Before the extraction, the abdomen of every specimen was poked laterally with a sterile needle to facilitate DNA extraction. Isolation was made using the NucleoSpin Tissue kit (Macherey-Nagel) following the manufacturer's instructions. Three molecular markers were amplified for *Liophloeodes*: two ribosomal: 28S-D2 (GenBank accession: MN190722-MN191039) and ITS2 (GenBank accession: MN191040-MN191233) and one mitochondrial: the standard COI barcoding region (GenBank accession: MT858362-MT858668), using primers as in Table 1. A nested PCR was used to amplify bacterial DNA, by per-

DNA marker	Primers	References
COI	LCO1490 / HC02198	(Folmer et al. 1994)
28S-D2	D2F / D2R	(Campbell et al. 1994)
ITS2	LC1 / HC2	(Navajas et al. 1992)
16S	27F / 1513R	(Weisburg et al. 1991)
16S Spiroplasma	27F / TKSSsp	(Fukatsu and Nikoh 2000)
16S Cardinium	CLOF / CLOR	(Weeks et al. 2003)
16S Arsenophonus	27F / ARS16SR	(Tsuchida et al.2002)
16S Rickettsia	Rb-F / Rb-R	(Gottlieb et al., 2006)
wsp Wolbachia	wsp_F1 / wsp_R1	(Baldo et al., 2006)
ftsZ Wolbachia	ftsZ_F1 / ftsZ_R1	(Baldo et al., 2006)
16S Microsporidia	V1 / 1492	(Vossbrinck and Friedman 1989; Zhu et al. 1993)
16S Nardonella	16SA1F / Nard733R	(White et al., 2015)

Table 1. Primers used in this study.

forming the first PCR using generic primers targeting the 16S gene (GenBank accession MN621120-MN621139), followed by a second PCR with primers specific to Wolbachia, Arsenophonus, Rickettsia, Spiroplasma, Cardinium, Nardonella and Microsporidia. For Wolbachia, ftsZ (GenBank accession MT500574-MT500577) and wsp (GenBank accession MT611140-MT611153) genes were amplified with primer pairs listed in Table 1. The PCR reaction was conducted using the following mix: 11.5 µl of ddH₂O, 2 µl of 10X DreamTaq buffer, 2 µl of 25 mM MgCl₂, 1 µl of 10 uM for each primer, 0.4 µl of 10 mM dNTPs, 0.1 µl of 5 U/µl Taq polymerase and 2 µl DNA. PCR products were sent for sequencing to the companies HongKe XiLin Biotechnology Co (China) and Macrogen (Netherlands). Obtained sequences were visually analysed and edited using SeqMan (Swindell and Plasterer 1997) or BioEdit (Hall 1999). COI sequences were translated into amino acids using the ExPASy translate tool (Gasteiger et al. 2003). Sequences were aligned by MAFFT (Katoh 2002) using the G-INS-1 algorithm. Due to many insertions and deletions, the ITS2 dataset was aligned using Fastgap (Borchsenius 2009), which allows for coding indels as traits for Bayesian analysis. Pseudomeira obscura Solari & Solari, 1907 (GenBank accessions HE818408 and HE818407) and Eusomus ovulum Germar, 1824 (KU341552 and MH746366), both from the tribe Entiminae were selected as outgroups for phylogenetic analyses based on nuclear markers, because both ITS2 and 28S-D2 were available for these species. For COI, E. ovulum (KU341536) was used along with Graptus triguttatus (Fabricius, 1775) (KY110616) and Prothrombosternus tarsalis Voss (1965) (KU748541), from the tribes Entiminae and Molytinae, respectively. For the phylogenetic reconstruction of Rickettsia symbionts, homologous sequences of several Rickettsia strains available in GenBank used are reported in Table S5. Evolutionary models for each alignment and the best partitioning scheme were chosen using PartitionFinder (Lanfear et al. 2017). Phylogenetic reconstructions were obtained by Bayesian inference using MrBayes v 3.2 with 1 cold and 3 heated Markov chains for 10,000,000 generations, and trees sampled every 1000th generation. (Huelsenbeck and Ronquist 2001) – there have been built tree for every marker and also tree from concatenated nuclear markers. Obtained trees were visualized using Fig-Tree 1.4.3 (Rambaut 2009) and graphically edited using CorelDraw Graphic Suite X7. Genetic distances within and between lineages were calculated using the *p*-distance method in MEGA 6 (Tamura et al. 2013). Phylogeography of *Liophloeodes* populations based on COI was inferred by statistical parsimony using the software TCS (Clement et al. 2000). The network was graphically edited using the software PopArt (Leigh and Bryant 2015) to better visualize taxonomic and geographic signals.

Morphological and morphometric study

After identifying the species based on the male aedeagus morphology (Fig. 2) every female Liophloeodes specimen from the same area was initially assigned to the same species, except for areas where males of two species were collected (those females were classified only as Liophloeodes). All collected specimens of Liophloeus sensu stricto were identified as Liophloeus tessulatus, based on the morphology and the geographical ranges of the species and were female (there are no bisexual populations in the sampling areas, see Smreczynski 1958). The specimens were dissected and their elytra, heads, antennae, legs, pronota, abdomens and spermathecae (females) were glued on cardboard and measured using a Nikon SMZ1500 binocular microscope and the NIS Elements BR 2.30 software (Fig. S1). Due to the destruction of some structures while collecting or mounting, and consequent lack of some measurements, missing data were replaced by the mean of measurements from a particular trait and species (Arbour and Broun 2014). Results of the measurements were presented separately for both sexes using Principal



Figure 4. Bayesian phylogenetic tree based on the ITS2 marker. (GenBank accessions: MN191040–MN191233). The first number in each collapsed clade is the total number of specimens sequenced, the number in parenthesis indicates unidentified specimens (from populations where there were no males). Numbers above branches represent posterior probabilities. *Eusomus ovulum* and *Pseudomeira obscura* were used as outgroups.

Component Analysis (PCA) in the R environment using a modified script (Baur and Leuenberger 2011). Results of PCA were analysed using the Generalized Linear Model (GLM) method with factor scores as dependent variables and species as the predictor (Kuszewska and Woyciechowski 2015). Principal components that explained most of the variance were used in the analysis. When differences were statistically significant, the Tukey post-hoc test was performed to assess differences between species.

Results

Phylogenetic trees and networks

Phylogenies based on 28S-D2 (581 bp) and ITS2 (750 bp – length of full alignment) were consistent with morphological identification based on the aedeagal shape (Fig. 2), with each clade including male specimens assigned to only one particular species (along with the females from the same collection area, where available) (Figs 4–6). However, topologies of trees and networks differed by markers. ITS2 tree (Fig. 4) was divided into two main sister clades: [Liophloeus tessulatus + [Liophloeus (Liophloeodes) lentus +Liophloeus (Liophloeodes) gibbus)]] and [Liophloeus (Liophloeodes) liptoviensis + [Liophloeodes) eus (Liophloeodes) pupillatus + Liophloeus (Liophloeodes) eus (Liophloeodes) pupillatus + Liophloeus (Liophloeodes) eus (Liophloeodes) eus

des) herbstii]]. However, 28S tree topology (Fig. 5) was different with three clades: [Liophloeus (Liophloeodes) lentus], [Liophloeus (Liophloeodes) gibbus], [Liophloeus (Liophloeodes) liptoviensis + Liophloeus (Liophloeodes) pupillatus + Liophloeus tessulatus] and unresolved Liophloeus (Liophloeodes) herbstii. Heterozygosity (double peaks in the chromatograms) was detected at diagnostic polymorphic sites of 28S and ITS2 sequences of specimens occupying contact zones (Fig. S4). Combined tree from both nuclear markers was, similarly to ITS2 tree divided into two main clades: [Liophloeus tessulatus + [Liophloeus (Liophloeodes) lentus + Liophloeus (Liophloeodes) gibbus]] and [Liophloeus (Liophloeodes) liptoviensis + [Liophloeus (Liophloeodes) pupillatus + Liophloeus (Liophloeodes) herbstii]] (Fig.6).

Phylogeny based on the 650-bp COI alignment was incongruent with traditional taxonomy and nuclear phylogenies (Fig. S2, Figs 7–11). *Liophloeus tessulatus* was the only monophyletic clade (Fig. S2, Fig. 7, clade A). Three clades corresponding to three morphological species, [*Liophloeus* (*Liophloeodes*) liptoviensis, *Liophloeus* (*Liophloeodes*) lentus and *Liophloeus* (*Liophloeodes*) gibbus)] could be distinguished (Fig. S2), yet each of those clades also included specimens from the other species (at least specimens from two species in one clade). The *Liophloeus* (*Liophloeodes*) liptoviensis clade includes specimens from this species along with specimens from *Liophloeus* (*Liophloeodes*) gibbus, *Liophloeus* (*Liophloeodes*) herbstii and *Liophloeus* (*Liophloeodes*) pupillatus









Figure 6. Bayesian phylogenetic tree based on the combined markers 28S-D2 and ITS (GenBank accessions: MN190722–MN191039 and MN191040–MN191233). The first number is the number of specimens in the clade, the number in parentheses is the number of unidentified specimens (from populations where there were no males). Numbers above branches represent posterior probabilities. *Eusomus ovulum* and *Pseudomeira obscura* were used as outgroups.



Figure 7. Bayesian phylogenetic tree of the subgenus *Liophloeodes* based on COI sequences (GenBank accession: MT858362-MT858668): *Liophloeus tessulatus* and *Liophloeus (Liophloeodes) liptoviensis*, clades A and B.

from neighbour populations (Fig. 7, clade B). Also, in the *Liophloeus* (*Liophloeodes*) *lentus* clade, we can find specimens from neighbouring populations of *Liophloeus* (*Liophloeodes*) *liptoviensis* (Fig. 8, clade C). *Liophloeus* (*Liophloeodes*) *gibbus* clade includes few subclades (Figs 9–11, clades D–H), one of them (Fig. 9, clade D) most-



Figure 8. Bayesian phylogenetic tree of the subgenus *Liophloeodes* based on COI sequences: *Liophloeus* (*Liophloeodes*) *lentus*, clade C.

ly contains *Liophloeus* (*Liophloeodes*) *lentus* specimens from Poland, localized western from the contact zone between *Liophloeus* (*Liophloeodes*) *lentus* and *Liophloeus* (Liophloeodes) gibbus (Dunajec and Poprad valley, Fig.3). Liophloeus (Liophloeodes) herbstii and Liophloeus (Liophloeodes) liptoviensis can also be found in the Lio-



Figure 9. Bayesian phylogenetic tree of the subgenus *Liophloeodes* based on COI sequences: *Liophloeus* (*Liophloeodes*) *lentus* and *Liophloeus* (*Liophloeodes*) gibbus, clade D.

phloeus (*Liophloeodes*) *gibbus* clade. In some clades, a geographical structure could be seen. For example, there were subclades of *Liophloeus* (*Liophloeodes*) *liptoviensis*

clade that included *Liophloeus* (*Liophloeodes*) *liptoviensis* specimens from Romania, Ukraine, or Polish Western Carpathians, along with specimens from different



Figure 10. Bayesian phylogenetic tree of the subgenus *Liophloeodes* based on COI sequences: *Liophloeus* (*Liophloeodes*) gibbus, clade E.

species from the same areas (Fig. 7, clade B). Similarly, all *Liophloeus* (*Liophloeodes*) *lentus* from Slovakia formed one big subclade, which also included specimens of *Liophloeus* (*Liophloeodes*) *liptoviensis* from Slovakia (Fig. 8, clade C). All Polish populations of *Liophloeus* (*Liophloeodes*) *lentus* clustered in one of the *Liophloeus*



Figure 11. Bayesian phylogenetic tree of the subgenus *Liophloeodes* based on COI sequences *Liophloeus* (*Liophloeodes*) gibbus and *Liophloeus* (*Liophloeodes*) lentus, clades F&G&H.

(*Liophloeodes*) gibbus subclades, along with some Polish *Liophloeus* (*Liophloeodes*) gibbus specimens (Fig. 9, clade D). All and *Liophloeus* (*Liophloeodes*) pupillatus specimens gathered in the *Liophloeus* (*Liophloeodes*) *liptoviensis* clade (Fig. 7, clade B). The COI statistical parsimony networks (Figs. 12a, 12b) also confirmed the presence of a strong geographic signal, with different species connected by reticulation events.

Genetic distances

For all three markers and all *Liophloeodes* species, the highest proportions of differing nucleotides were found

when they were paired with *Liophloeus tessulatus*. The most differing species among *Liophloeodes* species was *Liophloeus (Liophloeodes) liptoviensis*, however the differences within this subgenus were minor. For COI: minimum *Liophloeus (Liophloeodes). liptoviensis-Liophloeus (Liophloeodes) pupillatus 0.9%*, maximum *Liophloeus (Liophloeodes) herbstii-Liophloeus (Liophloeodes) lentus 8.6%*; for 28S-D2: minimum: *Liophloeus (Liophloeodes) gibbus 0.4%*, maximum *Liophloeus (Liophloeodes) gibbus 0.4%*, maximum *Liophloeus (Liophloeodes) lentus 2.1%*, *Liophloeus (Liophloeodes) lentus 2.1%*, *for ITS minimum: Liophloeus (Liophloeodes) lentus 3.6%*, maximum: *Liophloeus (Liophloeodes) lentus 2.6%*, maximum: *Liophloeus (Liophloeodes) lentus 2.1%*, *for ITS minimum: Liophloeus (Liophloeodes) lentus 3.6%*, maximum: *Liophloeus Liophloeus (Liophloeodes) lentus 3.6%*, maximum: *Liophloeus Liophloeus Liophloeus (Liophloeus 3.6%*, maximum: *Liophloeus Liophloeus Liophloeus (Liophloeus 3.6%*, maximum: *Liophloeus Liophloeus (Liophloeus 3.6%*, maximum: *Liophloeus Liophloeus 1.6%*, maximum: *Liophloeus (Liophloeus 3.6%*, maximum: *Liophloeus 1.6%*, maximum: *Lioph*



Figure 12a. TCS network inferred from mitochondrial sequences, with groups corresponding to species. The relative size of circles is proportional to the number of sequences of the same haplotype.

phloeus (Liophloeodes) gibbus-Liophloeus (Liophloeodes) pupillatus 19.7%). The highest distances were detected for the ITS2 and the lowest for 28S-D2 (Tables S2–S4).

Endosymbiont survey

Of the seven groups of symbionts searched for in Liophloeodes, only two were found, Wolbachia and Rickettsia. (Table S6). Wolbachia and Rickettsia were recorded in 43% and 76% of tested specimens respectively. All obtained Wolbachia wsp sequences were identical and they belonged to the strain that can be also found in other beetles (Otiorhynchus singularis GU111688, Byturus ochraceus AJ585380). Wolbachia ftsZ sequences were identical, and this strain has also been found in many other arthropod groups (spiders-MN594716, wasps-MH742743, flies-CP042904, butterflies-KC959172). Differently, Rickettsia 16S rDNA sequences were more diverse than ftsZ and wsp, but they were all closely related to the ones that have been previously found in other weevil species (Fig. S3).





Figure 13b. Principal component analysis (PCA) of morphometric measurements of males. Scatterplot shows first against second shape PC. The variance explained by each principal component is given in parentheses.

Morphometrics

Females of all *Liophloeodes* species grouped together in the PCA scatter plot, however, in the plot of the first against the second shape PC, females of *Liophloeus tessulatus* formed a distinct group (Fig. 13a). In the PCA scatter plot of first against second shape PC for males, all species were overlapping (Fig. 13b). PCA for females showed a positive correlation of the elytra length (0.29) and width (0.24), abdomen length (0.25) and width (0.28), distance between eyes (0.25), femur (0.26), tibia (0.28), pronotum length (0.27) and width (0.26), scape length (0.24), rostrum width (0.26) and head width (0.29) with the first component, which integrated information about the body shape. The first principal component was used in the GLM as it explained 60% of the variance (with 10% of variance explained by the second principal component). Results of GLM showed that there is a statistically significant difference (p<0.001) in factor scores from the PCA between species. *Post-hoc* Tukey test for females confirmed the distinctiveness of *Liophloeus tessulatus* [statistically significant differences in all pairings,

except for pairing with Liophloeus (Liophloeodes) liptoviensis with Liophloeus tessulatus being significantly bigger] and showed a statistically significant difference between Liophloeus (Liophloeodes) liptoviensis, and Liophloeus (Liophloeodes) gibbus, with Liophloeus (Liophloeodes) liptoviensis being significantly bigger. PCA for males showed strong negative correlation of the elytra length (-0.29) and width (-0.27), abdomen length (-0.28)and width (-0.3), distance between eyes (-0.25), femur (-0.28), tibia (-0.29), pronotum length (-0.29) and width (-0.29) scape length (-0.28), rostrum width (-0.26) and head width (-0.3) with the first component, which integrated information about the body shape. The first principal component was used in the GLM, as it explained 72% of the variance (with 7% of the variance explained by the second principal component). Results of GLM showed that there is a statistically significant difference (p < 0.001) in factor scores from the PCA between species. Post-hoc Tukey test for males showed significant difference between Liophloeus (Liophloeodes) liptoviensis and Liophloeus (Liophloeodes) lentus, with Liophloeus (Liophloeodes) lentus being significantly bigger.

Discussion

Diversity of Liophloeodes

Results of traditional species identification were usually consistent with data from faunistic papers, with some differences. In few regions where the subgenus was previously recorded no Liophloeodes were found (northern Slovenia, Croatia, a big part of south-western Romania). In northern Slovenia (Triglav) specimens of Liophloeus (Liophloeodes) liptoviensis were collected instead of Liophloeus (Liophloeodes) herbstii, which was the species known from this region. This last issue may be explained by the previous misidentification of these two species (the aedeagi of both species are often similar). Some new areas of distribution for Liophloeodes were also found (e.g., Balkan Mountains in Bulgaria). Nuclear markers showed full congruence with morphological identification of species (Figs 4-6), whereas the mitochondrial marker displayed a strong geographic pattern that suggests widespread hybridization and introgression events (Fig. S2, Figs 7-12a). This may be at least partly explained by the presence of two bacterial endosymbionts, Wolbachia and Rickettsia, which infect the most of species, with no pattern of particular symbiont infecting a particular species or population occupying a particular area. Detected Wolbachia and Rickettsia strains were previously found in other weevils (Malloch and Fenton 2005; Merville et al. 2013) and other beetles (Roehrdanz et al. 2019), respectively.

The lack of Microsporidia might be surprising due to its detection in the earlier study (Ovcharenko et al. 2013). However, the detection was based on one population from the contact zone between *Liophloeus* (*Liophloeodes*) *lentus* and *Liophloeus* (*Liophloeodes*) *gibbus* in the Dunajec and Poprad Valley, so we can assume that Microsporidia infection is occasional in this subgenus.

The hypothesis of past hybridization and subsequent introgression seems to be supported by the detection of evidence of heterozygosity: double peaks in chromatograms, mostly at polymorphic and diagnostic sites of 28S-D2 and ITS2 sequences of specimens occupying contact zones (Fig. S4). The lack of ecological differences (here confirmed during collecting) along with similar endosymbiont infections and low genetic interspecific distances also suggest that Liophloeodes species may consist of lineages that have not fully sorted yet. Further support to this hypothesis is provided by the morphometric study, which did not show clear differences between Liophloeodes species, suggesting that they have not differentiated morphologically yet. Great caution should be taken when concluding about distinction between Liophloeus (Liophloeodes) herbstii and Liophloeus (Liophloeodes) pupillatus based on morphometric data, as only a few specimens were examined in our study.

As for the status of *Liophloeus tessulatus*, this species strongly differs from all *Liophloeodes* species morphologically, ecologically (it occurs both in wet/cold as in dry/warm biotopes, whereas *Liophloeodes* is restricted only to the former habitat type) and sexually (so far only bisexual populations of *Liophloeodes* have been detected). However, phylogenetic analyses based on nuclear markers consistently placed it within *Liophloeodes* clades (Figs 4–6) and heterozygosity was also detected in this species. To sum up, traditional systematics of *Liophloeodes* by ribosomal phylogenies, but the integration of different types of data along with the status of *Liophloeus tessulatus* that is not congruent with traditional systematics lead to some questions that we attempt to address.

Incongruence of mtDNA and nuclear DNA phylogeny

Lack of congruence between mtDNA and nuclear DNAs is often found in phylogenetic studies on many groups of organisms (Roca et al 2005; Larmuseau et al. 2010; Lumme et al. 2017; Thielsch et al. 2017; Wallis et al. 2017; Weigand et al. 2017) including insects (Linnen and Farrell 2007; Gompert et al. 2008; Hinojosa et al. 2019). Two main causes of this problem are: a) incomplete lineage sorting and b) introgression following hybridization. When the incongruence between two markers has a geographical structure (for example, when one mtDNA haplotype is shared by two species or by geographically close populations of those species), the incongruence is suspected to be the result of hybridization. When populations under speciation were divided by some geographic barrier for a long time and then came into secondary contact, accumulated mutations may not allow successful mating and genetic mixing of the two species. However, in some cases, a stable hybridization zone might emerge and introgression of mtDNA from one species to another might occur on a large scale (Toews and Brelsford 2012). This can even lead to the replacement of the "native" haplotypes by the introgressed ones (Babik et al. 2005). Disagreement between the nuclear and mitochondrial DNA phylogenies may be strongly influenced by natural selection (Boratyński et al. 2011; Toews et al. 2014), infection by bacterial endosymbionts (Hurst and Jiggins 2005; Whitworth et al. 2007; Gompert et al. 2008), sexual selection, unequal survival of hybrids, differences in survival and dispersion between sexes (Bonnet et al. 2017). Moreover, invasions of populations to new areas are also often followed by stronger introgression from the local population to the invading one (Currat et al. 2008; Phuong et al. 2017).

While the Liophloeus tessulatus clade in the COI tree consists only of specimens from this species (and there are no specimens from this species in other clades), the other three Liophloeodes clades include specimens from more than one species, with one dominating in each. All clades have a strong geographical structure (Fig. S2, Fig. 12b). Probably contact between those two species was followed by hybridization and introgression, which might have led to the removal of Liophloeus (Liophloeodes) lentus COI haplotypes from Polish populations. Furthermore, some specimens belonging to this clade live in the Sudetes, which is a separated mountain range in the west, almost 300 km from the contact zone. Thus, it can be speculated that the present contact zone is just one of the areas where hybridization and introgression occurred and/or that Liophloeus (Liophloeodes) lentus populations dispersed west. The other area where hybridization and introgression could have occurred or is currently occurring is the eastern part of Polish Carpathians, where Liophloeus (Liophloeodes) liptoviensis and Liophloeus (Liophloeodes) gibbus are often sympatric. Heterozygosity detected in specimens from this area may support the hypothesis that this is an ongoing process. The heterozygosity found in nuclear markers, and a COI phylogeny with clades of intermixed species suggests also other places where hybridization could occur, such as Tatra in Slovakia [Liophloeus (Liophloeodes) lentus and Liophloeus (Liophloeodes) liptoviensis], Romania [Liophloeus (Liophloeodes) herbstii and Liophloeus (Liophloeodes) liptoviensis], and Pannonian Basin in Hungary and Serbia [different populations of Liophloeus (Liophloeodes) lentus]. One of the factors that could increase introgression is Wolbachia spread between the species and populations (hitchhiking effect), and this endosymbiont was detected mostly in areas of possible hybridization (Miyata et al. 2020). Incongruence between gene trees often occurs in species that underwent many geographical range shifts in the past and now occur sympatrically in several areas, often in montane environments (Rodríguez et al. 2010; Haines et al. 2017; Ortego et al. 2017; Tóth et al. 2017), a scenario that suits Liophloeodes well.

Status of Liophloeus tessulatus

Liophloeus tessulatus includes both parthenogenetic and bisexual populations, yet in areas where it is sympatric

with Liophloeodes no bisexual populations were found. Based on this information and the unexpected position of Liophloeus tessulatus in the phylogenetic trees (Figs 4-6), we propose two possible scenarios for its evolutionary history. According to the first one, lineages of Liophloeodes and Liophloeus sensu stricto (including Liophloeus tessulatus) diverged long time ago and then both evolved and adapted to different environmental conditions. A consequence would be that the position of Liophloeus tessulatus in the phylogenetic trees is misleading, and its taxonomic position should be established based on genetic distances of all three markers (Tables S2–S4), morphology, and ecology. In this scenario, Liophloeodes populations, restricted to wet and cold biotopes, would shrink their ranges to mountain refugia during times of climate warming, which would lead to their diversification and subsequent speciation. The more cosmopolitan Liophloeus sensu stricto would be more resistant to climate change events and this would, along with its partially parthenogenetic mode of reproduction, lead to its present wide range of distribution. Because in populations sympatric with Liophloeodes those weevils are parthenogenetic, it is possible that divergence between the two lineages occurred somewhere inside present Liophloeodes range and the parthenogenetic type of reproduction evolved only once. The second hypothesis derives from the position of *Liophloeus tessulatus* in the phylogenetic trees, and according to it, Liophloeus tessulatus is one of Liophloeodes species that diverged during subgenus evolution along with other species. Thus, different morphology and ecological requirements are the results of different adaptations during climate change events and range shifts. However, the ancestor of Liophloeus tessulatus changed its ecological niche, expanded its range and a new type of reproduction evolved. Niche shifts were one of the key factors in weevil evolution and diversification (Marvaldi et al. 2002) and parthenogenetic forms often invaded new areas, also in Europe in glacial and interglacial periods (Kajtoch et al. 2009; Kajtoch et al. 2012) However, the parthenogenetic reproduction may be quite a new adaptation in this lineage. Heterozygosity found in Liophloeus tessulatus may be a result of recent hybridizations between bisexual populations. It is also worth mentioning that parthenogenetic populations of Liophloeus tessulatus are triploid (Suomalainen 1955), which, along with the 28S-D2 heterozygosity might indicate a hybrid origin of this species - hybridization is in fact likely the main cause of emerging parthenogenetic forms among weevils (Stenberg and Lundmark 2004; Neiman et al. 2009). The sequence of these events (emergence of parthenogenesis, niche shift, geographic expansion, possible hybridization) remains unsolved. This hypothesis suggests that bisexual, montane populations of Liophloeus tessulatus from Western Europe are a distinct species that diverged before the niche shift and evolution of parthenogenesis. However, without examining those populations, no definitive conclusions about the status of Liophloeus tessulatus can be derived. If the second hypothesis was true, Liophloeus tessulatus should be included in the Liophloeodes subgenus.

Low morphometric diversity

Results of the morphometric analysis suggest small shape differences in selected body parts between species, which seem to correspond with Smreczyński's suggestion (1958) that the aedeagal shape is the only trait that can be treated as diagnostic, and the other traits display a gradient of variation among species. To support (or reject) this conclusion, traits mentioned by Smreczyński such as the shape of the rostrum (how it expands to the end), tooth on the femur, the structure of the abdomen's margin were visually checked, and the validity of his opinion was confirmed. Given the ecological requirements of Liophloeodes species, these results are not surprising. All species prefer the same biotope and this ecological conservatism probably led to speciation - changing conditions caused the division of ranges and withdrawal to glacial refugia, which was likely followed by isolation and speciation. This association of diverged species to the same biotope, along with repeated hybridizations that may have occurred, might result in small morphometric diversity.

There is another interesting aspect of this problem: small interspecific diversity can be a result of high intraspecific diversity that probably evolved before speciation, so it can be considered a kind of ancestral polymorphism (Williams et al. 2015). However, it is important to remember that males of two species [*Liophloeus* (*Liophloeodes*) lentus and *Liophloeus* (*Liophloeodes*) liptoviensis)] display significant diversity, which could be the result of an ongoing differentiation. The presence of hybrid specimens with intermediate features might have heavily affected the morphometric results, erasing the diversity between species groups, which may cause problems with species identification (Nugnes et al. 2017).

Inconsistency between phylogenies of nuclear markers

Lack of congruence between phylogenies derived from different nuclear markers happens more rarely than between nuclear and mitochondrial DNA, but their causes are similar: introgression and incomplete lineage sorting. The random distribution of specimens in the phylogenetic tree usually suggests the latter cause (Page and Charleston 1997; Buckley et al. 2006; Vaezi and Brouillet 2009). This is not the case of *Liophloeodes* because all species are clearly distinguished in all nuclear marker trees (Figs 4–6), and there are no signs of interspecific admixture (as it is in COI – Fig. S2). The main difference is the distribution of species in the trees and their histories.

Two things should be considered here. The first is the conservatism of the 28S-D2 rDNA marker, which can affect the phylogeny of young, recently derived species (Jordal and Kambestad 2014). ITS2 marker is far more sensitive to recent divergence processes (Jousselin et al. 2006). The second argument is that ITS2 phylogeny is more robust, based on the statistical support of nodes. The entire ITS2 tree is indeed fully resolved and strong-

ly supported, whereas the placement of *Liophloeus* (*Liophloeodes*) herbstii, *Liophloeus* (*Liophloeodes*) lentus and *Liophloeus* (*Liophloeodes*) gibbus in the 28S rDNA tree is unsupported. Morevoer the topology of ITS2 tree is consistent with the tree from combined nuclear marker.

Evolutionary history of Liophloeodes

Based on the ITS2 phylogeny, there are two main phylogeographic groups of Liophloeodes. The first, consisting of Liophloeus (Liophloeodes) lentus and Liophloeus (Liophloeodes) gibbus, is the most frequent in the north of the Pannonian Basin, and surrounds it nearly completely through (in clockwise geographical order): West Carpathians, Podolian Upland, Apuseni, Milevska Planina/ Vitosha, Dinaric Alps, the Sudetes, and Western Carpathians. On the east, there are more than 400 km of break between Apuseni and Vitosha. Among these two species, Liophloeus (Liophloeodes) lentus is the westernmost one (it occurs in the Alps, the Sudetes, Pannonian Basin, and Western Carpathians). It is interesting that in this tree, this clade is sister to generally more western Liophloeus tessulatus, which may support the hypothesis of divergence of this species by niche shifts.

The second ITS2 clade [(Liophloeus (Liophloeodes) liptoviensis + [(Liophloeus (Liophloeodes) pupillatus + (Liophloeus (Liophloeodes) herbstii]] has more eastern distribution, with one exception [Liophloeus (Liophloeodes) liptoviensis from Slovenia]. It can be hypothesised that the first strong divergence in the history of Liophloeodes was between populations occupying central Carpathians and populations from more western and southern mountain ranges.

Probably recurrent glacial and interglacial events caused recurrent changes in the ranges of distribution with expansions and reductions. Due to *Liophloeodes* ecological requirements, these weevils probably expanded their ranges during glaciations and withdrew to refugia during interglacial periods (which we can observe now, as they occur mostly in mountain valleys). Maybe climate changes caused the division of distinct species lineages and populations of diverged lineages came into secondary contact when their ranges expanded. Some of them might have also dispersed through mountain ranges, even during interglaciations because the climate was stable there. For example, sympatry of Liophloeus (Liophloeodes) gibbus and Liophloeus (Liophloeodes) liptoviensis in the eastern part of Polish Carpathians can be the result of the dispersion of both species through Carpathians (the first one to the west, the latter to the east).

Detailed *Liophloeodes* phylogeography is difficult to resolve. The topology of all phylogenetic trees suggests strong intraspecific gene flow between populations – we can observe small clades consisting of populations from wider areas such as Poland and Romania. Alternatively, many small clades consist of specimens restricted to very limited ranges. A possible explanation for this is that the evolution of this subgenus was influenced by repeating isolations of populations in refugia, which could lead

Table 2. Synthetic summary of the integrative approach with all independent lines of evidence listed. Y=starting hypothesis supported; N=starting hypothesis not supported; P=starting hypothesis partially supported; IPC=integration by partial congruence; species hypothesis stability: S=stable, U=unstable.

Liophloeodes species	Morphology (H0)	28S-D2	ITS2	COI	Morphometry	Symbionts	IPC
Liophloeus (Liophloeodes) lentus	Y	Y	Y	Р	N	Ν	S
Liophloeus (Liophloeodes) gibbus	Y	Y	Y	Р	N	Ν	S
Liophloeus (Liophloeodes) liptoviensis	Y	Y	Y	Р	N	Ν	S
Liophloeus (Liophloeodes) herbstii	Y	Y	Y	N	N	Ν	U
Liophloeus (Liophloeodes) pupillatus	Y	Y	Y	N	N	Ν	U

to speciation, and by subsequent contacts of already diverged populations that resulted in hybridization.

Conclusions

Six independent lines of evidence were used: traditional morphology (as the starting hypothesis), phylogenies based on three molecular markers, morphometry and endosymbiont occurrence/phylogeny (Table 2). The latter two methods were inconclusive in supporting species differentiation. Phylogeny based on both ribosomal markers supported traditional morphology, whereas the different topology of the COI phylogeny can be explained by hybridization and introgression events. Examining all the lines of evidence considered we conclude that:

1. Although not supported by all lines of evidence, we argue that traditional systematics of *Liophloeodes* based on the shape of genitalia, an important trait for reproductive isolation (Langerhans 2016), should not be changed, lacking conclusive evidence to challenge taxonomic stability.

2. Species of *Liophloeodes* probably represent young lineages with evidence of hybridization and/or introgression detected in the COI phylogeny and the heterozygosity found in nuclear markers of specimens from contact zones.

3. The status of *Liophloeus tessulatus* and the *Liophloeus sensu stricto* subgenus remains uncertain. Morphology, morphometrics, ecology, reproduction, and genetic data suggest its distinctiveness from *Liophloeodes*, however phylogenetic analyses indicate that it may be another *Liophloeodes* species. For more sound conclusions on the taxonomic status of this species, and of the *Liophloeus* s.s subgenus, more populations (especially bisexual) and the remaining species of the subgenus, respectively, must be examined.

Acknowledgments

We would like to thank Stanisław Knutelski, Łukasz Kajtoch and Miłosz Mazur for help in the material collection. We are grateful to Anna Giulia Nappo and Liberata Gualtieri for help in the laboratory work. We express our special thanks for professor Bogusław Petryszak who is no longer with us for his help and inspiration for years. The research has been funded by the Polish National Science Center (Preludium grant 2015/17/N/NZ8/01571). We would also like to thank Rob Barber (UK) for proofreading.

References

- Apfelbeck V (1928) Fauna insectorum balcanica. IX. Ad cognitionem curculionidarum (Col.). Pars II. Beiträge zur Kenntnis der Gattungen Otiorhynchus Germ., Tropiphorus Schonh., Liophloeus Germ., Liparus Oliv., Plinthomeleus Rttr. Und Scleropterus Schonh. Mit Beschreibung Neuer Arten. Glasnik Zemaljskog Muzeja Bosni i Hercehovini. Sarajevo, 40: 75–86.
- Arbour J, Brown C (2014) Incomplete specimens in geometric morphometric analyses. Methods in Ecology and Evolution 5(1). https://doi. org/10.1111/2041-210X.12128
- Babik W, Branicki W, Crnobrnja-Isailović J, Cogălniceanu D, Sas I, Olgun K, Poyarkov NA, Garcia-París M, Arntzen JW (2005) Phylogeography of two European newt species-discordance between mtD-NA and morphology. Molecular Ecology 14(8): 2475–91. https:// doi.org/10.1111/j.1365-294X.2005.02605.x
- Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MCJ, Tettelin H, Werren JH (2006) Multilocus sequence typing system for the endosymbiont *Wolbachia pipientis*. Applied and Environmental Microbiology 72: 7098–7110. https://www.doi.org/10.1128/AEM.00731-06
- Baur H, Leuenberger C (2011) Analysis of ratios in multivariate morphometry. Systematic Biology 60(6): 813–825. https://doi. org/10.1093/sysbio/syr061
- Bonnet T, Leblois R, Rousset F, Crochet PA (2017) A reassessment of explanations for discordant introgressions of mitochondrial and nuclear genomes. Evolution 71(9): 2140–2158. https://doi.org/ 10.1111/evo.13296
- Boratyński Z, Alves P, Berto S, Koskela E, Mappes T, Melo-Ferreira J (2011) Introgression of mitochondrial DNA among *Myodes* voles: Consequences for energetics? BMC Evolutionary Biology 11(1). https://doi.org/10.1186/1471-2148-11-355
- Borchsenius F (2009). FastGap 1.2 (p. Department of Biosciences, Aarhus University, Denm). https://www.aubot.dk/FastGap_home.htm
- Brown SDJ (2017) A revision of the New Zealand weevil genus Irenimus Pascoe (1876) (Coleoptera: Curculionidae: Entiminae). Zootaxa 4263(1): 1–42. https://doi.org/10.11646/zootaxa.4263.1.1
- Buckley TR, Cordeiro M, Marshall DC, Simon C (2006) Differentiating between hypotheses of lineage sorting and introgression in New Zealand Alpine Cicadas (Maoricicada Dugdale). 55(3): 411–425.
- Campbell BC, Steffen-Campbell JD, Werren JH (1994) Phylogeny of the *Nasonia* species complex (Hymenoptera: Pteromalidae) inferred from an internal transcribed spacer (ITS2) and 28S rDNA

sequences. Insect Molecular Biology 2(4): 225–237. https://doi.org/ 10.1111/j.1365-2583.1994.tb00142.x

- Clement M, Posada D, Crandall KA (2000) TCS: A computer program to estimate gene genealogies. Molecular Ecology 9: 1657–1659. https://doi.org/10.1046/j.1365-294X.2000.01020.x
- Currat M, Ruedi M, Petit RJ, Excoffier L (2008) The hidden side of invasions: Massive introgression by local genes. Evolution 62(8): 1908–1920. https://doi.org/10.1111/j.1558-5646.2008.00413.x
- Dayrat B (2005) Towards integrative taxonomy. Biological Journal of the Linnean Society 85: 407–415. https://doi.org/10.1111/j.1095-8312.2005.00503.x
- Dieckmann L (1980) Beiträge zur Insektenfauna der DDR: Coleoptera
 Curculionidae (Brachycerinae, Otiorhynchinae, Brachyderinae).
 Beiträge Zur Entomologie. Berlin, 30(1): 145–310.
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology 3(5): 294–299. https://doi.org/10.1371/journal. pone.0013102
- Fukatsu T, Nikoh N (2000) Endosymbiotic microbiota of the bamboo pseudococcid Antonina crawii (Insecta, Homoptera). Applied and Environmental Microbiology 66(2): 643–650. https://doi. org/10.1128/AEM.66.2.643-650.2000
- Gasteiger E, Gattiker A, Hoogland C, Ivanyi I, Appel RD, Bairoch A (2003) ExPASy: The proteomics server for in-depth protein knowledge and analysis. Nucleic Acids Research. https://doi.org/10.1093/ nar/gkg563
- Gebiola M, Gómez-Zurita J, Monti MM, Navone P, Bernardo U (2012) Integration of molecular, ecological, morphological and endosymbiont data for species delimitation within the *Pnigalio soemius* complex (Hymenoptera: Eulophidae). Molecular Ecology 21(5): 1190– 1208. https://doi.org/10.1111/j.1365-294X.2011.05428.x
- Gompert Z, Forister ML, Fordyce JA, Nice CC (2008) Widespread mito-nuclear discordance with evidence for introgressive hybridization and selective sweeps in *Lycaeides*. Molecular Ecology 17(24): 5231–5244. https://doi.org/10.1111/j.1365-294X.2008.03988.x
- Gottlieb Y, Ghanim M, Chiel E, Gerling D, Portnoy V, Steinberg S, Tzuri G, Rami A, Belausov E, Mozes-daube N, Gershon M, Gal S, Katzir N, Zchori-fein E, Horowitz AR, Kontsedalov S (2006). Identification and localization of a *Rickettsia* sp. in *Bemisia tabaci* (Homoptera: Aleyrodidae). Applied and Environmental Microbiology 72(5): 3646–3652. https://doi.org/10.1128/AEM.72.5.3646
- Grobler GC, Janse van Rensburg L, Bastos ADS, Chimimba CT, Chown SL (2006) Molecular and morphometric assessment of the taxonomic status of *Ectemnorhinus* weevil species (Coleoptera: Curculionidae, Entiminae) from the sub-Antarctic Prince Edward Islands. Journal of Zoological Systematics and Evolutionary Research 44(3): 200–211. https://doi.org/10.1111/j.1439-0469.2006.00358.x
- Haines ML, Stuart-Fox D, Sumner J, Clemann N, Chapple DG, Melville J (2017) A complex history of introgression and vicariance in a threatened montane skink (*Pseudemoia cryodroma*) across an Australian sky island system. Conservation Genetics 18(4): 939–950. https://doi.org/10.1007/s10592-017-0945-7
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium 41: 95–98. https://doi.org/10.14601/phytopathol_mediterr-14998u1.29
- Hinojosa J, Koubínová D, Szenteczki M, Pitteloud C, Dinca V, Alvarez N, Vila R (2019) A mirage of cryptic species: Genomics uncover striking mitonuclear discordance in the butterfly *Thymelicus sylves*-

tris. Molecular ecology 28(17): 3857–3868. https://doi.org/10.1111/ mec.15153

- Hurst GDD, Jiggins FM (2005) Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: the effects of inherited symbionts. Proceedings. Biological Sciences/ The Royal Society 272(1572): 1525–1534. https://doi.org/10.1098/ rspb.2005.3056
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogeny. Bioinformatics 17: 754–755
- Jordal BH, Kambestad M (2014) DNA barcoding of bark and ambrosia beetles reveals excessive NUMTs and consistent east-west divergence across Palearctic forests. Molecular Ecology Resources 14(1): 7–17. doi:10.1111/1755-0998.12150
- Jousselin E, Van Noort S, Rasplus JY, Greeff JM (2006) Patterns of diversification of Afrotropical Otiteselline fig wasps: Phylogenetic study reveals a double radiation across host figs and conservatism of host association. Journal of Evolutionary Biology 19(1): 253–266. https://doi.org/10.1111/j.1420-9101.2005.00968.x
- Kajtoch Ł, Korotyaev B, Lachowska-Cierlik D (2012) Genetic distinctness of parthenogenetic forms of European *Polydrusus* weevils of the subgenus *Scythodrusus*. Insect Science 19(2):183–194. https:// doi.org/10.1111/j.1744-7917.2011.01448.x
- Kajtoch Ł, Lachowska-Cierlik D, Mazur M (2009) Genetic diversity of the xerothermic weevils *Polydrusus inustus* and *Centricnemus leucogrammus* (Coleoptera: Curculionidae) in Central Europe. European Journal of Entomology 106(3): 325–334. https://doi.org/ 10.14411/eje.2009.040
- Katoh K (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research 30(14): 3059–3066. doi:10.1093/nar/gkf436
- Kuszewska K, Woyciechowski M (2015) Age at which larvae are orphaned determines their development into typical or rebel workers in the honeybee (*Apis mellifera L.*). PLoS ONE 10(4). https://doi. org/10.1371/journal.pone.0123404
- Lanfear R, Frandsen PB, Wright AM, Senfeld T, Calcott B (2017) Partitionfinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. Molecular Biology and Evolution 34(3): 772–773. https://doi.org/10.1093/ molbev/msw260
- Langerhans RB, Anderson CM, Heinen-Kay JL (2016) Causes and Consequences of Genital Evolution. Integrative and Comparative Biology 56(4): 741–751. https://doi.org/10.1093/icb/icw101
- Larmuseau MHD, Raeymaekers JAM, Hellemans B, Houdt JKJ Van, Volckaert FAM (2010) Mito-nuclear discordance in the degree of population differentiation in a marine goby. Heredity 105(6): 532– 542. https://doi.org/10.1038/hdy.2010.9
- Leigh JW, Bryant D (2015) POPART: Full-feature software for haplotype network construction. Methods in Ecology and Evolution 6: 1110–1116. https://doi.org/10.1111/2041-210X.12410
- Linnen C, Farrell B (2007) Mitonuclear discordance is caused by rampant mitochondrial introgression in *Neodiprion* (Hymenoptera: Diprionidae) sawflies. Evolution; International Journal of Organic Evolution 61: 1417–1438. https://doi.org/10.1111/j.1558-5646.20-07.00114.x
- Lumme J, Ziętara MS, Lebedeva D (2017) Ancient and modern genome shuffling: Reticulate mito-nuclear phylogeny of four related allopatric species of *Gyrodactylus* von Nordmann, 1832 (Monogenea: Gyrodactylidae), ectoparasites on the Eurasian minnow *Phoxinus phoxinus* (L.) (Cyprinidae). Systematic Parasitology 94(2): 183– 200. https://doi.org/10.1007/s11230-016-9696-y

Marvaldi AE, Sequeira AS, O'Brien CW, Farrel BD (2002) Molecular and morphological phylogenetics of weevils (Coleoptera, Curculionoidea): Do niche shifts accompany diversification? Systematic Biology 51(5): 761–785. https://doi.org/10.1080/10635150290102465

- McKenna DD, Sequeira AS, Marvaldi AE, Farrell BD (2009) Temporal lags and overlap in the diversification of weevils and flowering plants. Proceedings of the National Academy of Sciences 106(17): 7083–7088. https://doi.org/10.1073/pnas.0810618106
- Merville A, Venner S, Henri H, Vallier A, Menu F, Vavre F, Heddi F, Bel-Venner M (2013) Endosymbiont diversity among sibling weevil species competing for the same resource. BMC Evolutionary Biology 13: 28. https://doi.org/10.1186/1471-2148-13-28
- Miralles A, Vasconcelos R, Perera A, Harris DJ, Carranza S (2011) An integrative taxonomic revision of the Cape Verdean skinks (Squamata, Scincidae). Zoologica Scripta 40(1): 16–44. https://doi.org/10.-1111/j.1463-6409.2010.00453.x
- Miyata MN, Nomura M, Kageyama D (2020) Wolbachia have made it twice: Hybrid introgression between two sister species of Eurema butterflies. Ecol Evol. 10: 8323–8330. https://doi.org/10.1002/ ece3.6539
- Navajas M, Cotton D, Kreiter S, Gutierrez J (1992) Molecular approach in spider mites (Acari: Tetranychidae): preliminary data on ribosomal DNA sequences. Experimental & Applied Acarology 15(4): 211–218. https://doi.org/10.1007/BF01246563
- Neiman M, Meirmans S, Meirmans PG (2009) What can asexual lineage age tell us about the maintenance of sex? Annals of the New York Academy of Sciences 1168: 185–200. https://doi.org/10.1111/ j.1749-6632.2009.04572.x
- Nugnes F, Bernardo U, Viggiani G (2017) An integrative approach to species discrimination in the *Anagrus atomus* group sensu stricto (Hymenoptera: Mymaridae), with a description of a new species. Systematics and Biodiversity 15: 582–599. https://doi.org/10.1080 /14772000.2017.1299811
- Oberprieler R, Marvaldi A, Anderson R (2007) Weevils, weevils, weevils everywhere. Zootaxa. 520: 491–520. https://doi.org/10.11646/ zootaxa.1668.1.24
- Ortego J, Noguerales V, Cordero PJ (2017) Geographical and ecological drivers of mitonuclear genetic divergence in a Mediterranean grass-hopper. Evolutionary Biology 44: 505–521. https://doi.org/10.1007/s11692-017-9423-x
- Ovcharenko M, Świątek P, Ironside J, Skalski T (2013) Orthosomella lipae sp. n. (Microsporidia) a parasite of the weevil, Liophloeus lentus Germar, 1824 (Coleoptera: Curculionidae). Journal of Invertebrate Pathology 112(1): 33–40. https://doi.org/10.1016/j.jip.2012.10.007
- Padial JM, Miralles A, De la Riva I, Vences M (2010) The integrative future of taxonomy. Frontiers in Zoology 7:16. doi:10.1186/1742-9994-7-16
- Page RDM, Charleston MA (1997) From Gene to Organismal Phylogeny: reconciled trees and the gene tree/species tree problem. Molecular Phylogenetics and Evolution 7(2): 231–240. https://doi. org/10.1006/mpev.1996.0390
- Petri K (1912) Siebenbiirgens Kaferfauna auf Grund ihrer Erforschung bis zum Jahre 1911. In Sieberbiirgischen Vereins fur Naturwissenschaften zu Hermannstadt.
- Phuong MA, Bi K, Moritz C (2017) Range instability leads to cytonuclear discordance in a morphologically cryptic ground squirrel spe-

cies complex. Molecular Ecology 26(18) 4743- 4755. https://doi. org/10.1111/mec.14238

- Rambaut A (2009) FigTree, a graphical viewer of phylogenetic trees. Institute of Evolutionary Biology University of Edinburgh. http:// tree.bio.ed.ac.uk/software/figtree
- Reitter E (1916) Fauna Germanica. Die Käfer des Deutschen Reiches. 5 Bände, K. G. Lutz, Stuttgart.
- Roca AL, Georgiadis N, Brien SJO (2005) Cytonuclear genomic dissociation in African elephant species. Nat Genet. 37(1): 96–100. https://doi.org/10.1038/ng1485
- Rodríguez F, Pérez T, Hammer SE, Albornoz J, Domínguez A (2010) Integrating phylogeographic patterns of microsatellite and mtD-NA divergence to infer the evolutionary history of chamois (genus *Rupicapra*). BMC Evolutionary Biology 10(1). https://doi.org/ 10.1186/1471-2148-10-222
- Roehrdanz ARL, Wichmann SS (2019) Wolbachia Multilocus sequence typing of singly infected and multiply infected populations of northern Corn Rootworm (Coleoptera: Chrysomelidae) Annals of the Entomological Society of America 107: 832–841. https://doi. org/10.1603/AN14006
- Schlick-Steiner BC, Steiner FM, Seifert B, Stauffer C, Christian E, Crozier RH (2010) Integrative taxonomy: a multisource approach to exploring biodiversity. Annual Review of Entomology 55: 421–438. https://doi.org/10.1146/annurev-ento-112408-085432
- Schutze MKMK, Virgilio M, Norrbom A, Clarke ARAR (2017) Tephritid integrative taxonomy: where we are now, with a focus on the resolution of three tropical fruit fly species complexes. Annual Review of Entomology 62(1): 147–164. https://doi.org/10.1146/annurev- ento-031616-035518
- Shin S, Clarke DJ, Lemmon AR, Moriarty Lemmon E, Aitken AL, Haddad S, Farrell BD, Marvaldi AE, Oberprieler RG, Mckenna DD (2018) Phylogenomic data yield new and robust insights into the phylogeny and evolution of weevils. Molecular Biology and Evolution 35(4): 823–836. https://doi.org/10.1093/molbev/msx324
- Shropshire JD, Leigh B, Bordenstein SR (2020) Symbiont-mediated cytoplasmic incompatibility: What have we learned in 50 years? Elife (9). doi: 10.7554/eLife.61989Smreczyński S (1958) Vorstudien zu einer Monographie des Subgenus *Liophloeodes*. Acta Zoologica Cracoviencia 3(3): 67–120.
- Stec D, Krzywański Ł, Zawierucha K, Michalczyk Ł (2020a) Untangling systematics of the *Paramacrobiotus areolatus* species complex by an integrative redescription of the nominal species for the group, with multilocus phylogeny and species delineation within the genus *Paramacrobiotus*. Zoological Journal of the Linnean Society 188(3): 694–716. https://doi.org/10.1093/zoolinnean/zlz163
- Stee D, Tumanov DT, Kristensen RM (2020b) Integrative taxonomy identifies two new tardigrade species (Eutardigrada: Macrobiotidae) from Greenland. European Journal of Taxonomy, 614, 1–40. https:// doi.org/10.5852/ejt.2020.614
- Stenberg P, Lundmark M (2004) Distribution, mechanisms and evolutionary significance of clonality and polyploidy in weevils. Agricultural and Forest Entomology 6(4): 259–266. https://doi.org/10.1111/ j.1461-9555.2004.00231.x
- Suomalainen E (1955) A further instance of geographical parthenogenesis and Polyploidy in the weevils. Archivum Societatis Botanicae Zoologicae Fennicae "Vanamo" 9: 350–354.
- Swindell SR, Plasterer TN (1997) SEQMAN. Contig assembly. Methods in Molecular Biology (Clifton, N.J.) 70: 75–89. https://doi. org/10.1385/0-89603-358-9:75

- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. Molecular Biology and Evolution 30(12): 2725–2729. https://doi.org/10.1093/ molbev/mst197
- Thielsch A, Knell A, Mohammadyari A, Petrusek A, Schwenk K (2017) Divergent clades or cryptic species? Mito-nuclear discordance in a Daphnia species complex. BMC Evolutionary Biology 17(1): 1–9. https://doi.org/10.1186/s12862-017-1070-4
- Toews DPL, Brelsford A (2012) The biogeography of mitochondrial and nuclear discordance in animals. Molecular Ecology 21(16): 3907–3930. https://doi.org/10.1111/j.1365-294X.2012.05664.x
- Toews DPL, Mandic M, Richards JG, Irwin DE (2014) Migration, mitochondria, and the yellow-rumped warbler. Evolution 68(1): 241– 255. https://doi.org/10.1111/evo.12260
- Toševski I, Caldara R, Jović J, Baviera C, Hernández-Vera G, Gassmann A, Emerson BC (2014) Revision of *Mecinus heydenii* species complex (Curculionidae): integrative taxonomy reveals multiple species exhibiting host specialization. Zoologica Scripta, 43(1): 34–51. https://doi.org/10.1111/zsc.12037
- Tóth JP, Varga ZS, Verovnik R, Wahlberg N, Váradi A, Bereczki J (2017) Mito-nuclear discordance helps to reveal the phylogeographic patterns of *Melitaea ornate* (Lepidoptera: Nymphalidae). Biological Journal of the Linnean Society 20: 1–15. https://doi.org/10.1093/ biolinnean/blw037
- Tsuchida T, Koga R, Shibao H, Matsumoto T, Fukatsu T (2002) Diversity and geographic distribution of secondary endosymbiotic bacteria in natural populations of the pea aphid, *Acyrthosiphon pisum*. Molecular Ecology 11(10): 2123–2135. https://doi.org/10.1046/j.1365-294x.2002.01606.x
- Vaezi J, Brouillet L (2009) Phylogenetic relationships among diploid species of *Symphyotrichum* (Asteraceae: Astereae) based on two nuclear markers, ITS and GAPDH. Molecular Phylogenetics and Evolution 51(3): 540–553. doi:10.1016/j.ympev.2009.03.003
- Vitecek S, Kučinić M, Previšić A, Živić I, Stojanović K, Keresztes L, Bálint M, Hoppeler F, Waringer J, Graf W, Pauls SU (2017) Integrative taxonomy by molecular species delimitation: multi-locus data corroborate a new species of Balkan Drusinae micro-endemics. BMC Evolutionary Biology 17(1): 1–18. https://doi.org/10.1186/ s12862-017-0972-5
- Vossbrinck CR, Friedman S (1989) A 28s ribosomal RNA phylogeny of certain cyclorrhaphous Diptera based upon a hypervariable region. Systematic Entomology 14(4): 417–431. https://doi. org/10.1111/j.1365-3113.1989.tb00296.x

- Wallis GP, Cameron-Christie SR, Kennedy HL, Palmer G, Sanders TR, Winter DJ (2017) Interspecific hybridization causes long-term phylogenetic discordance between nuclear and mitochondrial genomes in freshwater fishes. Molecular Ecology 26(12): 3116–3127. https:// doi.org/10.1111/mec.14096
- Weeks AR, Velten R, Stouthamer R (2003) Incidence of a new sex-ratio-distorting endosymbiotic bacterium among arthropods. Proceedings of the Royal Society B: Biological Sciences 270(1526): 1857–1865. https://doi.org/10.1098/rspb.2003.2425
- Weigand H, Weiss M, Cai H, Li Y, Yu L, Zhang C, Leese F (2017) Deciphering the origin of mito-nuclear discordance in two sibling caddisfly species. Molecular Ecology 26(20): 5705–5715. https:// doi.org/10.1111/mec.14292
- Weisburg WG, Barns SM, Pelletier DA, Lane DJ (1991) 16S ribosomal DNA amplification for phylogenetic study J Bacteriol. 173(2): 697– 703. https://doi.org/10.1128/jb.173.2.697–703.1991
- Weise J (1894) Zur Gattung Liophloeus Germ. Deutsche Entomologische Zeitschrift, Berlin, 1894(2): 257–266. [VIII–1894]
- White JA, Richards NK, Laugraud A, Saeed A, Curry MM, McNeill MR (2015) Endosymbiotic candidates for parasitoid defense in exotic and native New Zealand Weevils. Microbial Ecology 70(1): 274–286. https://doi.org/10.1007/s00248-014-0561-8
- Whitworth TL, Dawson RD, Magalon H, Baudry E (2007). DNA barcoding cannot reliably identify species of the blowfly genus *Protocalliphora* (Diptera: Calliphoridae). Proceedings. Biological Sciences / The Royal Society 274: 1731–1739. https://doi.org/10.1098/ rspb.2007.0062
- Williams PH, Byvaltsev AM, Cederberg B, Berezin MV, Ødegaard F, Rasmussen C, Richardson LL, Huang J, Sheffield CS, Williams ST (2015) Genes suggest ancestral colour polymorphisms are shared across morphologically cryptic species in arctic bumblebees. PLoS ONE: 10(12). https://doi.org/10.1371/journal.pone.0144544
- Yeates DK, Seago A, Nelson L, Cameron SL, Joseph L, Trueman JWH (2011) Integrative taxonomy, or iterative taxonomy? Systematic Entomology 36(2): 209–217. https://doi.org/10.1111/j.1365-3113.-2010.00558.x
- Zhu X, Wittner M, Tanowitz HB, Kotler D, Cali A, Weiss LM (1993) Small subunit rRNA sequence of *Enterocytozoon bieneusi* and its potential diagnostic role with use of the polymerase chain reaction. Journal of Infectious Diseases 168(6): 1570–1575. https://doi. org/10.1093/infdis/168.6.1570

Supplementary material 1

File 1

Authors: Wacławik et al. (2021) Data type: .pdf

Explanation note: Figure S1. Morphometric measurements.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/asp.79.e64252.suppl1

Supplementary material 2

File 2

Authors: Wacławik et al. (2021) Data type: .pdf

- Explanation note: Figure S2. A Bayesian phylogenetic tree of the subgenus Liophloeodes based on COI sequences (GenBank accession: MT858362–MT85-8668), with Eusomus ovulum as an outgroup. Bayesian posterior probability values are given above branches. Scale bar represents 0.03 substitutions per site.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.
- Link: https://doi.org/10.3897/asp.79.e64252.suppl2

Supplementary material 3

File 3

Authors: Wacławik et al. (2021) Data type: .pdf

Explanation note: Figure S3. A Bayesian phylogenetic tree of Rickettsia sequences based on 16S rDNA sequences (GenBank accession MN621120–MN621139) with Orientia tsutsugamushi, Wolbachia-E. formosa, Wolbachia-N. vitripennis as outgroups. Bayesian posterior probability values are given above branches. Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited. Link: https://doi.org/10.3897/asp.79.e64252.suppl3

Supplementary material 4

File 4

Authors: Wacławik et al. (2021)

Data type: .pdf

- **Explanation note: Figure S4.** Double peak indicating heterozygosity.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/asp.79.e64252.suppl4

Supplementary material 5

File 5

Authors: Wacławik et al. (2021)

Data type: .xlsx

Explanation note: Table S1. List of collected specimens. Species in bold (column D) have been recognized based on the morphological study of particular specimens, species with normal font have been recognized based on study of different males found in the same place.

Copyright notice: This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/asp.79.e64252.suppl5

Supplementary material 6

File 6

Authors: Wacławik et al. (2021) Data type: .xlsx

- **Explanation note: Table S2.** Uncorrected p-distance values. Interspecific COI percentage genetic distances are below diagonal, with standard deviations above diagonal.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/asp.79.e64252.suppl6

Supplementary material 7

File 7

Authors: Wacławik et al. (2021) Data type: .xlsx

- **Explanation note: Table S3.** Uncorrected p-distance values. Interspecific 28S-D2 percentage genetic distances are below diagonal, with standard deviations above diagonal.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.
- Link: https://doi.org/10.3897/asp.79.e64252.suppl7

Supplementary material 8

File 8

Authors: Wacławik et al. (2021) Data type: .xlsx

Explanation note: Table S4. Uncorrected p-distance values. Interspecific ITS percentage genetic distances are below diagonal, with standard deviations above diagonal.

- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.
- Link: https://doi.org/10.3897/asp.79.e64252.suppl8

Supplementary material 9

File 9

Authors: Wacławik et al. (2021)

Data type: .pdf

- **Explanation note: Table S5.** Accession numbers of Rickettsia strains available from Genbank used in phylogenetic analysis.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.
- Link: https://doi.org/10.3897/asp.79.e64252.suppl9

Supplementary material 10

File 10

Authors: Wacławik et al. (2021)

Data type: .xlsx

- **Explanation note: Table S6.** Endosymbionts detected in *Liophloeus*.
- **Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons. org/licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/10.3897/asp.79.e64252.suppl10

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Arthropod Systematics and Phylogeny

Jahr/Year: 2021

Band/Volume: 79

Autor(en)/Author(s): Waclawik Benjamin, Nugnes Francesco, Bernardo Umberto, Przybycien Maja, Lachowska-Cierlik Dorota

Artikel/Article: An integrative revision of the subgenus Liophloeodes (Coleoptera: <u>Curculionidae: Entiminae: Polydrusini): taxonomic, systematic, biogeographic and</u> <u>evolutionary insights 419-441</u>