<u>PENSOFT</u>

SENCKENBERG world of biodiversity



More than Olceclostera bifenestrata: New species and morphology of immature stages of Olceclostera Butler, 1879 (Lepidoptera: Bombycoidea, Apatelodidae)

Elton Orlandin¹, Mônica Piovesan¹, Daniel Herbin², Eduardo Carneiro¹

1 Laboratório de Estudos de Lepidoptera Neotropical, Departamento de Zoologia, Universidade Federal do Paraná. P.O. Box 19020, 81531-980 Curitiba, Paraná, Brazil

2 Le Clos de Lutché, F-31380 Garidech, France

https://zoobank.org/F59B7D6A-008D-4D8E-8727-D6E7E818D1E7

Corresponding author: Elton Orlandin (orlandinelton@gmail.com)

Received9 June 2023Accepted15 October 2023Published20 December 2023

Academic Editors Andreas Zwick, Martin Wiemers

Citation: Orlandin E, Piovesan M, Herbin D, Carneiro E (2023) More than *Olceclostera bifenestrata*: New species and morphology of immature stages of *Olceclostera* Butler, 1879 (Lepidoptera: Bombycoidea, Apatelodidae). Arthropod Systematics & Phylogeny 81: 1063–1088. https://doi. org/10.3897/asp.81.e107507

Abstract

Apatelodidae is a family of Neotropical bombycoids that still needs to be studied in several aspects, as many groups of species have not yet undergone a careful systematic revision. On the other hand, recent studies showed that some species known to be widely distributed in fact form species complexes. Until now, *Olceclostera bifenestrata* Schaus, 1912, described from Costa Rica, supposedly has a wide distribution, reaching the south of Brazil. We reviewed specimens from South America identified as *O. bifenestrata*. Using morphological data and DNA barcodes, we discovered that South American specimens belong to four new species: *Olceclostera jairana* **sp. nov.**, *Olceclostera quilombola* **sp. nov.**, *Olceclostera xeta* **sp. nov.** and *Olceclostera wayana* **sp. nov.** The four species can be distinguished mainly by characters of genitalia, mostly in aedeagus structures. However, wing characters (number of hyaline spots) and abdomen characters (number of patches of larger scales) may be useful for classifying species groups in *Olceclostera*. Additionally, we illustrate and describe for the first time all the life stages of an *Olceclostera* species, including its chaetotaxy and life history, and provide the first SEM images of the egg of an Apatelodidae species. We present a discussion about the morphological characters of adults and immatures and their relevance to the systematics of Apatelodidae.

Keywords

American silkworm moth, *Olceclostera jairana* **sp. nov.**, *Olceclostera quilombola* **sp. nov.**, *Olceclostera wayana* **sp. nov.**, *Olceclostera xeta* **sp. nov.**, systematics, taxonomy

1. Introduction

Apatelodidae is a family of Neotropical bombycoids (Lepidoptera: Bombycoidea) that lacks conspicuous colors or economic or medical importance. As a consequence, the group has been largely neglected in biodiversity studies, especially if compared to other large-sized Bombycoidea (e. g. Saturniidae and Sphingidae) (Kitching et al. 2018; Ballesteros Mejia et al. 2020). More recently, some taxonomic efforts carried out mainly by Herbin (Herbin 2015,

Copyright Elton Orlandin et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

2017, 2018, 2021a, 2021b; Herbin and Monzon Sierra 2015; Herbin and Beccacece 2018; Herbin and Mielke 2018; Herbin et al. 2021) have shown that the family is more diverse than previously thought. The study of the male and female genitalia structures and the use of molecular tools has shown that many species comprise cryptic species complexes (Herbin 2017; Herbin et al. 2021; Orlandin and Carneiro 2021). Currently, the family comprises 200 species and ten subspecies, distributed in 13 genera (Kitching et al. 2018; Hamilton et al. 2019; Herbin 2021a, 2021b; Herbin et al. 2021; Orlandin and Carneiro 2021) but studies testing the monophyly of the genera are lacking.

The genus Olceclostera Butler, 1879, was described in Notodontidae, based on Olceclostera irrorata Butler, 1879 and subsequently transferred to Lasiocampidae as a synonym of Parathyris Hübner 1819 (Kirby 1892). However, the synonymy was ignored by Schaus, who first described species of Olceclostera in Bombycidae (Schaus 1892) but later in Eupterotidae (Schaus 1894, 1900, 1905, 1910, 1912, 1920). Subsequently, Draudt (1929) transferred Olceclostera to Zanolinae (Bombycidae), currently a synonym of Apatelodidae (Zwick 2011), where the genus stands. Olceclostera harbors 31 species (Kitching et al. 2018; Herbin, 2021a, 2021b), 27 described until 1929 (Draudt 1929), based only on a few external characters (e.g. color, wing shape, and the number of hyaline spots). Based on Draudt (1929), some species are still believed to have a wide distribution. An example of this is Olceclostera bifenestrata Schaus, 1912. Described from Costa Rica, O. bifenestrata is cited as a widely distributed species (Draudt 1929; Biezanko 1986), and specimens from South America are identified as this species in collections.

The immature stages of Olceclostera species are also poorly known. Only the larvae of two North American species have been described (Lintner 1874; Packard 1895; Franclemont 1973; Stehr 1987; Peigler 1994; Wagner 2005), and additional images of some species from Central America can be found in online databases (Janzen and Hallwachs 2009). Also, host plant records are only available for a few species (Franclemont 1973; Biezanko et al. 1974; Wagner 2005; Janzen and Hallwachs 2009; Robinson et al. 2010; Diniz et al. 2013). However, none of the studies provides a detailed description of all immature stages, including the larval chaetotaxy, an important set of characters that contribute to the systematics of many groups of Lepidoptera (Peterson 1962; Scoble 1992) and which have shown substantial differences when comparing the immatures of some genera of Apatelodidae (Orlandin et al. 2021, 2022a, 2022b).

Aiming to test if the specimens from South America recognized as *O. bifenestrata* belong to this species, the morphology of males and females, including genitalia, was studied. To confirm if this morphological similarity reflects a close relationship, we perform a molecular analysis based on COX1 (barcode region, 658pb). Additionally, we describe in detail all immature stages from one of these new species adding a discussion of its relevance to the systematics of Apatelodidae.

2. Material and methods

2.1. Specimens examined

About 200 specimens, including all *Olceclostera* types, were examined directly or indirectly from the following collections (abbreviations used throughout the text):

CDH – Personal collection of Daniel Herbin; Garidech, France. CUIC - Cornell University Insect Collection; Ithaca, New York, United States of America. DD - Personal collection of Diego Rodrigo Dolibaina; Vitória, Espírito Santo, Brazil. DZUP - Coleção Entomológica Padre Jesus Santiago Moure, Departamento de Zoologia, Setor de Ciências Biológicas, Universidade Federal do Paraná; Curitiba, Paraná, Brazil. MACN - Museo Argentino de Ciencias Naturales "Bernardino Rivadavia"; Buenos Aires, Argentina. MfN - Museum für Naturkunde; Berlin, Germany. MHNG – Muséum d'histoire naturelle de Genève; Geneva, Switzerland. MNRJ - Museu Nacional da Universidade Federal do Rio de Janeiro; Rio de Janeiro, Brazil. NHMUK - The Natural History Museum; London, United Kingdom. USNM - National Museum of Natural History, Smithsonian Institution; Washington, D.C., United States of America. VOB - Personal collection of Vitor Osmar Becker, Reserva Serra Bonita; Camacan, Bahia, Brazil. ZSM - Zoologische Staatssammlung München; Munich, Germany.

In addition, we accessed all original descriptions of *Olceclostera* species (Walker 1855, 1865; Grote 1864; Packard 1864; Butler 1879; Edwards 1886; Druce 1890; Schaus 1892, 1894, 1900, 1905, 1910, 1912, 1920, 1927; Dyar 1906; Jones 1908; Draudt 1929; Herbin 2018, 2021; Herbin and Mielke 2018). In the section "Material examined" where the information on the types is provided, the labels are separated from one another by transverse bars.

2.2. Morphology

Immature. The first instar was illustrated with and without the long setae that ornamented the body. These setae were carefully removed using tweezers to allow the visualization of its insertions and other structures present along the body. Images of the eggs were taken using scanning electron microscopy (SEM-JEOL JSM 6360-LV). Images of the egg 1st, and 2nd instars, and pupa were taken using an optical stereomicroscope adapted with focus stacking (Leica Application Suite Version 4.12.0 [Build 86]), and subsequent instars were recorded with a Sony DSC-HX100V digital camera. We used ImageJ Software (Schneider et al. 2012) to measure the eggs, cephalic capsules, larvae, and pupae. The terminology follows Peterson (1962) and Stehr et al. (1987).

Adults. For the description of adults, the head, palpi, legs, and abdomen were detached, soaked in hot 10% potassium hydroxide solution (KOH) for approximately 3 min, washed in water, examined, illustrated, and then stored in glass vials containing glycerin. Images were taken using an optical stereomicroscope adapted with focus stacking (Leica Application Suite Version 4.12.0 [Build 86]) and a Sony DSCHX100V digital camera. The terminology follows Kristensen (2003), but the term "fultura inferior" (= juxta) is applied *sensu* Petersen (1904); the term aedeagus is applied *sensu* Snodgrass (1935); and followed Zwick (2009) for the terminology of the semi–oval sclerotized plates located ventrally of the subscaphium [gnathos] (Fig. 1).

2.3. Immatures rearing

A female was collected using a reflective sheet placed adjacent to light during sampling at Nova Petrópolis, Joaçaba, Santa Catarina, Southern Brazil (27°06'02"S, 51°36'32"W) on 21–31 December 2019. The female was placed in a plastic container for oviposition. After hatching from the eggs, a set of hostplant species, including

Allophylus sp. (Sapindaceae), Duranta erecta L. (Verbenaceae), Handroanthus sp. (Bignoniaceae), Ilex paraguariensis A. St.-Hil. (Aquifoliaceae), Persea americana Mill. (Lauraceae), Psidium cattleyanum Sabine (Myrtaceae) and Schinus terebinthifolia Raddi (Anacardiaceae) were offered to the larvae. These hostplant species were chosen based on previous records of Apatelodidae host plants (Biezanko et al. 1974; Janzen and Hallwachs 2009; Robinson et al. 2010; Orlandin et al. 2021, 2022a, 2022b). The larvae were reared with the chosen hostplant species until pupae. Whenever necessary, the branches were replaced to maintain fresh, abundant, and high-quality leaves for the immatures. Immediately after each moult, the dry head capsules were retained and stored in glass vials. From the fifth instar on, the larvae were transferred to similar containers with the hostplant branches and a layer of dark soil beneath. A few specimens from each immature stage were killed through immersion in boiling



Figure 1. Terminology adopted for *Olceclostera* genitalia: A male, ventral view; B male, lateral view; C male, phallus, lateral view; D female, lateral view.

water, fixed in Kahle-Dietrich 10% (Borror and deLong 1971) for 72 h, subsequently preserved in ethanol 70%, and then deposited in the Immature Collection of Lepidoptera from the Entomological Collection "Pe. Jesus Santiago Moure, Universidade Federal do Paraná" under voucher number DZUPIL 0170.

2.4. Genetic divergence and phylogenetic inference

A series of 30 Apatelodidae specimens were selected for DNA sequencing. A leg was removed and sent for sequencing of the COX1 (barcode region, 658pb) at the Canadian Center for DNA Barcoding (CCDB) (Hebert et al. 2003). DNA extraction, PCR amplification, and sequencing of the COX1 (barcode region, 658pb) followed standard protocols (CCDB 2013). Posteriorly, we accessed the Barcode of Life Data Systems (BOLD) (Ratnasingham and Hebert 2007) and mined the FASTA files of the COX1 (barcode region, 658pb) of the Apatelodidae, mainly *Olceclostera* species. Sequence data of specimens analyzed here are available on GenBank and/ or BOLD Systems (for GenBank and BOLD Systems accession numbers, see Table S1).

Alignment was then performed with MAFFT (Katoh and Standley 2013). The most optimal model of nucleotide evolution was selected by ModelFinder (Kalyaanamoorthy et al. 2017) using the AICc criterion, and the best fit model according to AICc was GTR+F+I+G4. The tree was rooted with *Drepatelodes friburgensis* (Schaus, 1924) following Hamilton et al. (2019) that recovered *Drepatelodes* as the sister group of other Apatelodidae genera. Maximum likelihood phylogenies were inferred using IQ-TREE v. 1.6.8 (Nguyen et al. 2015), with branch supports reported as 5,000 standard bootstraps. Alignment, model selection, and phylogenetic analyses were performed in the multifunctional platform PhyloSuite (Zhang et al. 2020).

Genetic distances of the "O. bifenestrata group" were calculated by the Tamura-Nei distance model (Tamura and Nei 1993) in MEGA11 (Tamura et al. 2021).

3. Results

3.1. "Olceclostera bifenestrata group"

Comments. The "Olceclostera bifenestrata group" comprises eight species, namely: Olceclostera angelica (Grote, 1864), Olceclostera bifenestrata, Olceclostera indistincta (Edwards, 1886), Olceclostera jairana **sp. nov.**, Olceclostera quilombola **sp. nov.**, Olceclostera seraphica Dyar, 1906, Olceclostera wayana **sp. nov.** and Olceclostera xeta **sp. nov.**

Diagnosis. This group of species is easily distinguished from other species of *Olceclostera* by the following com-

bination of characters: forewings grayish, with sinuous brownish lines, two subapical hyaline spots, and one pair of large patches of broader scales laterally on the abdomen. The South American species are reliably separated from the North American species *O. angelica*, *O. indistincta* and *O. seraphica* by their larger size, less grayish coloring (the North American species are light grey) (see Franclemont 1973: plate 3, figs 14–16), molecular distance (Table 1), and geographic distribution (Schaus 1912; Franclemont 1973). They can be distinguished from *O. bifenestrata* (Fig. 2A, B) by the sclerotized subscaphium region with a narrow median ventral hook (Fig. 3).

3.1.1. Olceclostera jairana Orlandin and Piovesan sp. nov.

https://zoobank.org/45D8B4CB-9987-48F0-ADF7-791CF-1D05EE5

Figures 2C, D, 4A, B, 5, 6A, 7-16

Type locality. Parque Estadual do Guartelá, Tibagi Municipality, Paraná State, Brazil.

Diagnosis. Olceclostera jairana sp. nov. differs from O. quilombola sp. nov. by external and genitalia characters: on the forewing of O. jairana sp. nov., the two subapical hyaline spots between M_1 and M_3 are rounded, with the one between M2-M3 smaller and almost unnoticeable in some cases. Otherwise, these spots are rectangular and always visible in O. quilombola sp. nov. In the dorsal forewing, posterior to the hyaline spots, O. jairana sp. **nov.** has a sinuous band of light gray scales, absent in O. quilombola sp. nov. (Fig. 2C-F). Olceclostera jairana sp. nov. has all tibias with a band of long scales with light brown apex, while O. quilombola sp. nov. presents these scales with dark brown apex (Fig. 6A, B). In the genitalia, O. jairana sp. nov. presents, in lateral view, a more rounded tegumen, narrower socii, and a more elongated uncus, and, in ventral view, the anterior projection of the saccus is thinner than in O. quilombola sp. nov. The aedeagus of O. jairana sp. nov. presents apex with a lateral projection bearing one to three spines, while in O. quilombola sp. **nov.** this projection has no spines (Figs 10, 17).

Externally, O. jairana sp. nov. is very similar to O. xeta sp. nov. and O. wayana sp. nov., but O. jairana sp. nov. has all tibias with a band of long scales with light brown apex while O. xeta sp. nov. and O. wayana sp. nov. present these scales with grayish brown apex (Fig. 6A, C). Dorsally, the abdomen of O. jairana sp. nov. has a copper color, while O. xeta sp. nov. and O. wayana sp. nov. have a grayish color (Fig. 2C, D, G–J). The genitalia of O. jairana sp. nov. present, in lateral view, a narrower socci than O. xeta sp. nov. and O. wayana sp. nov., and the process of sacculus is about twice as big as in O. xeta sp. nov. and O. wayana sp. nov. has an apex with spines and vesica without cornuti, while in O. xeta sp. nov. and O. wayana sp. nov. the aedeagus does not have spines and the bas-



Figure 2. Adult male of *Olceclostera* species: *Olceclostera bifenestrata* from Costa Rica. A dorsal view; B ventral view. *Olceclostera jairana* sp. nov. Holotype male from Parque Estadual do Guartelá, Tibagi, Paraná, Brazil. C dorsal view; D ventral view. *Olceclostera quilombola* sp. nov. Holotype male from Parque Estadual das Lauráceas, Adrianópolis, Paraná, Brazil. E dorsal view; F ventral view. *Olceclostera xeta* sp. nov. Holotype male from RPPN Fazenda da Mata, Querência do Norte, Paraná, Brazil. G dorsal view; H ventral view. *Olceclostera wayana* sp. nov. Holotype male from French Guiana. I dorsal view; J ventral view.



Figure 3. Male genitalia of Olceclostera bifenestrata. A ventral view; B lateral view; C aedeagus, lateral view.



Figure 4. Adult female of *Olceclostera* species: *Olceclostera jairana* **sp. nov.**, from Nova Petrópolis, Santa Catarina, Brazil. **A** dorsal view; **B** ventral view. *Olceclostera quilombola* **sp. nov.**, from São Luís do Paraitinga, São Paulo, Brazil. **C** dorsal view; **D** ventral view. *Olceclostera xeta* **sp. nov.**, from Estancia Dimas, Alto Paraná, Paraguay. **E** dorsal view; **F** ventral view.

6	
S	
l_{o}	
2	
Ŭ,	
2	
\sim	
đ	
ŝ	
Ö	
·2	
ě	
st	
>	
8	
ă	
e o	
Ē.	
<u>_</u>	
5	
aı	
d	
Ę	
2	
άd	
а	
π	
2	
S1	
ıе	
G,	
÷	
q	
n.	
10	
stε	
S	
10	
ĕ	
2	
0	
- cu	
ō	
Я	
a	
S	
8	
ă	
Ð	
p	
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	
•	
Se	
o) se	
pb) se	
8pb) se	
558pb) se	
, 658pb) se	
n, 658pb) se	
ion, 658pb) se	
gion, 658pb) se	
region, 658pb) se	
e region, 658pb) se	
de region, 658pb) se	
ode region, 658pb) se	
rcode region, 658pb) se	
varcode region, 658pb) se	
(barcode region, 658pb) se	
e (barcode region, 658pb) se	
me (barcode region, 658pb) se	
gene (barcode region, 658pb) se	
gene (barcode region, 658pb) se	
K1 gene (barcode region, 658pb) se	
)X1 gene (barcode region, 658pb) se	
OX1 gene (barcode region, 658pb) se	
COX1 gene (barcode region, 658pb) se	
or COX1 gene (barcode region, 658pb) se	
for COX1 gene (barcode region, 658pb) se	
) for COX1 gene (barcode region, 658pb) se	
%) for COX1 gene (barcode region, 658pb) se	
(%) for COX1 gene (barcode region, 658pb) se	
ss (%) for COX1 gene (barcode region, 658pb) se	
ces (%) for COX1 gene (barcode region, 658pb) se	
nces (%) for COX1 gene (barcode region, 658pb) se	
tances (%) for COX1 gene (barcode region, 658pb) se	
istances (%) for COX1 gene (barcode region, 658pb) se	
distances (%) for COX1 gene (barcode region, 658pb) se	
c distances (%) for COX1 gene (barcode region, 658pb) se	
tic distances (%) for COX1 gene (barcode region, 658pb) se	
netic distances (%) for COX1 gene (barcode region, 658pb) se	
enetic distances (%) for COX1 gene (barcode region, 658pb) se	
genetic distances (%) for COX1 gene (barcode region, 658pb) se	
ie genetic distances (%) for COX1 gene (barcode region, 658pb) se	
ise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
wise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
irwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
vairwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
Pairwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
1. Pairwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
e 1. Pairwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	
ole 1. Pairwise genetic distances (%) for COX1 gene (barcode region, 658pb) se	

Moderline for the formation of the fo	1		,					,				4						,				
BO34719 <i>- 0 biferestria</i> D 3.110 <i>- 0 piferestria</i> D 3.110 <i>- 0 piferestria</i> D 3.110 <i>- 0 piferestria</i> D 3.110 <i>- 0 piferestria</i> D 3.111 <i>B</i> -1-1 <i>0 angelica</i> D 3.111 <i>B</i> -1-1 <i>n</i> D 3.111 <i>B</i> -1 <i>n</i> D 3.111 <i>B</i> -1 <i>n</i> D 3.111 <i>B</i> -1 <i>n</i> D 3.111 <i>B</i> -1 <i>n</i>			1	2	3	4	S	9	-	æ	6	10	11	12	13	14	15	16	17	18	19	20
PD35410-0bijmetrata000IIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIIII <th< td=""><td>JL/</td><td>ABO847-19 – O. bifenestrata</td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td><td></td></th<>	JL/	ABO847-19 – O. bifenestrata																				
$ (2.1B1187-14-C) \ argetica a brow and a b$	MI	APD354-10 – O. bifenestrata	0,000																			
$ (TB1186-1-0 \ argetica a brain br$	N	CLB1187-14 – <i>O. angelica</i>	0,043	0,043																		
ONS703-08-0000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000000	S	CLB1186-14 - <i>O</i> . angelica	0,041	0,041	0,002																	
ONS704-06-0indisincta00450045000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400040004000400	١Ħ	ONS703-08 – O. indistincta	0,045	0,045	0,002	0,003																
<b>CONB247-09-</b> C. seraphica $0.046$ $0.048$ $0.048$ $0.048$ $0.048$ $0.048$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.047$ $0.040$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$ $0.000$	Ħ	CONS704-08 – O. indistincta	0,045	0,045	0,002	0,003	0,000															
<b>CONB248:09-</b> C. seraphica <b>(0.044(0.045(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.047(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043(0.043</b>	Ť.	CONB247-09 – O. seraphica	0,046	0,048	0,046	0,048	0,048	0,048														
(094868 - C) jairana sp. nov. $(100$ $(009)$ $(008)$ $(003)$ $(009)$ $(010)$ $(008)$ $(003)$ $(003)$ $(010)$ $(000)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$ $(010)$	١Ħ	CONB248-09 – O. seraphica	0,044	0,046	0,045	0,047	0,047	0,047	0,005													
(0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0)         (0) <td>١X</td> <td>.094868 – <i>O. jairana</i> <b>sp. nov.</b></td> <td>0,100</td> <td>0,099</td> <td>0,091</td> <td>0,088</td> <td>0,093</td> <td>0,093</td> <td>0,098</td> <td>0,100</td> <td></td>	١X	.094868 – <i>O. jairana</i> <b>sp. nov.</b>	0,100	0,099	0,091	0,088	0,093	0,093	0,098	0,100												
094854 - 0 jairana <b>sp. nov.</b> 0,1000,0920,0910,0820,0920,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,09	R	.094867 – <i>O. jairana</i> <b>sp. nov.</b>	0,100	0,099	0,091	0,088	0,093	0,093	0,098	0,100	0,000											
094855 - <i>O</i> jairana <b>sp. nov.</b> 0,1020,1010,0950,0920,0970,0930,0930,0960,0960,0960,0000,010mmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmmm<	В.	094854 – <i>O. jairana</i> <b>sp. nov.</b>	0,100	0,099	0,091	0,088	0,093	0,093	0,098	0,100	0,000	0,000										
094853 - O. jairana <b>sp. nov.</b> 0,0920,0910,0980,0930,0930,0930,0930,0930,0930,0930,0930,0930,0940,0940,0160,0160,0160,0160,0160,0160,0160,0170,0170,0170,0170,0170,0170,0170,0170,0170,0170,0170,0170,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0140,0	SR	094855 – <i>O. jairana</i> <b>sp. nov.</b>	0,102	0,101	0,095	0,092	0,097	0,097	0,098	0,100	0,000	0,000	0,000									
094852 - O. jairana sp. nov.0,0960,0950,0980,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0910,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0110,0130,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,093	ЭR	094853 – <i>O. jairana</i> <b>sp. nov.</b>	0,092	0,091	0,090	0,088	0,093	0,093	0,090	0,096	0,009	0,009	0,009	0,010								
094851 - O. jairana <b>sp. nov.</b> 0,0950,0950,0910,0880,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0830,0830,0930,0930,0830,0830,0930,0930,0930,0830,0830,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0930,0	ЭR	094852 – O. jairana sp. nov.	0,096	0,095	0,090	0,088	0,093	0,093	0,095	0,100	0,016	0,016	0,016	0,016	0,009							
094850 - <i>O quitombola</i> <b>sp. nov.</b> 0,090 0,083 0,081 0,085 0,085 0,086 0,096 0,063 0,063 0,062 0,058 0,068 0,060 ··· (10) (10) (10) (10) (10) (10) (10) (10)	OR	094851 – <i>O. jairana</i> <b>sp. nov.</b>	0,095	0,095	0,091	0,088	0,093	0,093	0,092	0,097	0,017	0,017	0,017	0,018	0,011	0,017						
094899- <i>O. quilombola</i> <b>sp. nov.</b> 0,090 0,089 0,087 0,085 0,089 0,089 0,090 0,094 0,067 0,067 0,066 0,062 0,088 0,063 0,003 0,003 0,071 0,01 0,01 0,018 0,012 0,012 0,012 0,022 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,092 0,091 0,074 0,078 0,078 0,009 0,092 0,091 0,074 0,078 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,091 0,074 0,078 0,009 0,009 0,009 0,091 0,074 0,078 0,003 0,003 0,034 0,034 0,044 0,074 0,078 0,009 0,009 0,091 0,074 0,078 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,009 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,000 0,00	OR	094850 – <i>O. quilombola</i> <b>sp. nov.</b>	0,090	0,090	0,083	0,081	0,085	0,085	0,086	0,090	0,063	0,063	0,063	0,062	0,058	0,058	0,060					
094856 <i>-O. xeta</i> <b>sp. nov.</b> 0,105 0,104 0,111 0,108 0,114 0,114 0,109 0,103 0,092 0,092 0,092 0,092 0,092 0,081 0,088 0,077 0,068 0,071 0,1 0 0,0 0 0,009 0,090 0,092 0,092 0,092 0,092 0,092 0,092 0,091 0,074 0,078 0,009 0,099 0,092 0,092 0,092 0,092 0,092 0,092 0,091 0,074 0,078 0,009 0,099 0,093 0,093 0,092 0,093 0,092 0,091 0,074 0,078 0,009 0,009 0,009 0,003 0,003 0,035 0,035 0,034 0,044 0,078 0,003 0,003 0,003 0,004 0,012 0,004 0,012 0,002 0,092 0,092 0,092 0,092 0,092 0,092 0,091 0,074 0,078 0,009 0,003 0,003 0,003 0,034 0,034 0,044 0,078 0,003 0,003 0,003 0,034 0,044 0,078 0,003 0,004 0,091 0,074 0,078 0,003 0,003 0,034 0,044 0,078 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,004 0,	ЭR	094849 – <i>O. quilombola</i> <b>sp. nov.</b>	0,090	0,089	0,087	0,085	0,089	0,089	0,090	0,094	0,067	0,067	0,067	0,066	0,062	0,058	0,063	0,003				
094897 - <i>O. xeta</i> <b>sp. nov.</b> 0,106 0,105 0,098 0,101 0,101 0,101 0,097 0,101 0,097 0,095 0,095 0,095 0,096 0,091 0,074 0,078 0,096 0,091 0,074 0,078 0,095 0,095 0,095 0,095 0,095 0,096 0,091 0,074 0,078 0,005 0,003 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035 0,035	OR	094856 – <i>O. xeta</i> <b>sp. nov.</b>	0,105	0,104	0,111	0,108	0,114	0,114	0,099	0,103	0,092	0,092	0,092	0,092	0,081	0,088	0,087	0,068	0,071			
04898 <i>– O. xeta</i> <b>sp. nov.</b> 0,106 0,105 0,103 0,105 0,105 0,105 0,097 0,101 0,095 0,095 0,095 0,096 0,091 0,074 0,078 0,005 0,003 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,034 0,	R	094897 – <i>O. xeta</i> <b>sp. nov.</b>	0,106	0,105	0,098	0,101	0,101	0,101	0,097	0,101	0,095	0,095	0,095	0,095	0,089	0,096	0,091	0,074	0,078	0,009		
094899 - O. wayaaa sp. nov. $0,116$ $0,115$ $0,115$ $0,118$ $0,118$ $0,118$ $0,118$ $0,118$ $0,118$ $0,118$ $0,118$ $0,107$ $0,100$ $0,102$ $0,102$ $0,102$ $0,006$ $0,009$ $0,098$ $0,098$ $0,079$ $0,080$ $0,038$ $0,038$ $0,034$ $0,034$	OR	094898 – O. xeta sp. nov.	0,106	0,105	0,103	0,105	0,105	0,105	0,097	0,101	0,095	0,095	0,095	0,095	0,089	0,096	0,091	0,074	0,078	0,005	0,003	
	ЭR	094899 – O. wayana <b>sp. nov.</b>	0,116	0,115	0,115	0,118	0,118	0,118	0,107	0,110	0,102	0,102	0,102	0,102	0,096	660'0	0,098	0,079	0,080	0,035	0,038	0,034

al portion of the vesica is covered by small cornuti (Figs 10, 19, 21).

Description of adults (Figs 2C, D, 4A, B, 5, 6A, 7-11). Head. MALE. Vertex and frons pinkish beige; compound eyes naked; antenna approximately 1/3 the length of the costal margin of the anterior wing, bipectinate from the base to apex, ventrally naked, dorsally covered by pearl scales; labial palpus dorsally covered by brown scales, ventrally covered by a compact block of pinkish beige scales (Fig. 5A); first and second segment thick and slightly curved, the second about twice the length of the first, third segment reduced and oval (Fig. 5B); proboscis absent. FEMALE. Similar to the male, slightly smaller than the male's head; antenna, pecten shorter than the male's pecten; palpus thinner and shorter than the male's palpus (Fig. 4A, B). - Thorax. MALE. Dorsally and ventrally gray; dorsally with a longitudinal medial brownish line. - Legs. All legs smooth, femur and tibia gray; tibias with a band of long scales with light brown apex projecting towards the posterior region (Fig. 6A); tarsomeres brownish. Foreleg: coxa thick, developed, approximately 3/4 the length of the femur; femur about the same length as the tibia; tibia anteriorly flattened, epiphysis approximately half-length of the tibia (Fig. 7A). Midleg: femur slightly smaller than the tibia; distally tibia bears a pair of spurs, the inner one longer than the outer one (Fig. 7B). Hindleg: femur smaller than the tibia; distal half of tibia bears two pair of spurs, the inner longer than the outer ones (Fig. 7C). All tarsomeres have a similar length, posteriorly with scattered small spines; tarsal claws simple. FEMALE. Similar to the male, epiphysis approximately 1/3 the length of the tibia, narrower than the male; ventral surface distitarsus of fore, mid, and hindlegs with a patch of small spines and an inner row with four conspicuous spines (Fig. 7D). - Wings shape and venation. Venation (Fig. 8), same as that described for Olceclostera sp. by Costa-Lima (1950). MALE. Forewing length: 14–19.5 mm (n = 10); elongated triangular shape. Upperside: Ground color gravish with scattered black scales; basal region limited by a fine irregular black line; discal region marked between two sinuous brownish lines; on dcm, a patch of black scales surrounded by a patch of light gray scales; post-discal region with a thin irregular black line; two rounded subapical hyaline spots, between M₁ and  $M_3$ ; posterior to the hyaline spots a sinuous band of light gray scales; apical region light gray; between M₂ and CuA₂ marginal region with black scales and outer margin slightly crenulated and marked with black scales. Underside: Ground color similar to the upperside but more brownish; post-discal region with a sinuous brownish line followed by a thin, sinuous black line. Hindwing. Frenulum long and simple (Fig. 8). Upperside: Ground color grayish brown; on dcm, a faint black dot; postdiscal region with two faint sinuous brownish lines; outer margin, between M₃ and CuA₂, slightly crenulated and marked with black scales. Underside: Similar to the upperside, but with scattered black scales; dots and lines more marked; inner margin lighter. FEMALE. Forewing length: 19-22 mm (n = 06); general color similar to the male, the apex is more acute than in males; on dcm, the patch of black scales is smaller than in the male and absent on the underside; hyaline spot between M2-M3 reduced. Hindwing. Frenulum grayish brown, as multiple bristles, similar to other female Apatelodidae species (Orlandin et al. 2021, 2022a). Similar to the male, but on dcm the dot of black scales is smaller than in the male. — Abdomen. MALE. Dorsally coppery and gray, ventrally gray with scattered black scales; pleural region of segments A1 to A3 with patches of broader scales, creamy at the base and apically coppery (Fig. 9A); into this patch, on A2 and A3, a row of white scales (Fig. 9B). FEMALE. Dorsally grayer than the male; at the posterior region of each segment, a band of light gray scales; ventral and pleural region similar to the male. — Genitalia. MALE (Fig. 10). Tegumen shield-like, sclerotized in all its extension; anterior projection of saccus tapered, dorsally curved; ventrally, the posterior margin of saccus has a semicircular projection projected posteriorly; uncus dorsoventrally flattened, ventrally projected, 3/4 bifid V-shaped, with a pair of laterally flattened socii at the base; subscaphium sclerotized, with a median ventral hook, proximal region, ventrally with two semi-oval sclerotized plates [gnathos] with numerous tiny spines; valva less sclerotized than uncus and tegumen, rectangular, tongue-like, laterally covering the uncus apex; costa (= processus superior) forming a slightly sclerotized process bearing some setae; sacculus (= processus inferior) slightly more sclerotized than valva; process of sacculus finger-like, projected at 90° from the valva; fultura inferior sclerotized and rectangular; aedeagus cylindrical with truncated caecum, ventrally projected forming a protuberance; in lateral view, apex with a projection bearing one to three spines (more frequently two); these structures are best seen when the vesica is everted. FEMALE (Fig. 11). Eighth tergum formed by a trapezoidal plate bearing wide lateral arms ventrally reaching the lamella antevaginalis; anterior apophysis thin and cylindrical, slightly spatulate at the tip; lamella antevaginalis flat, almost as long as the lamella postvaginalis; lamella postvaginalis sclerotized, rectangular; ductus bursae about two times the length of corpus bursae, approximately the same diameter for the entire length; corpus bursae ovoid, membranous with a signum heart-shaped, bearing inward-pointing spines and with the central portion less sclerotized with fewer spines; papilla analis dorsoventrally elongated, sclerotized and covered with setae.



Figure 5. Male labial palpus of *Olceclostera jairana* sp. nov. A with sclaes; B without scales.



Figure 6. Lateral view of the scales that cover the tibia of *Olceclostera*. A *Olceclostera jairana* sp. nov.; B *Olceclostera quilombola* sp. nov.; C *Olceclostera xeta* sp. nov. The arrow points to the apex of the scales, which shows a different brown hue in each species.



Figure 7. Legs of *Olceclostera jairana* sp. nov. A male foreleg, lateral view; B male midleg, lateral view; C male hindleg, lateral view; D female, foreleg tarsomeres.

**Description of immature stages (Figs 12–16).** *Eggs* (Fig. 12). Average duration (from oviposition to hatching): approximately 15 days (n = 110); pie-shaped 1.5–



Figure 8. Olceclostera jairana sp. nov. male wing shape and venation.



Figure 9. *Olceclostera jairana* sp. nov. Large patch of broader scales on the abdomen. A lateral view; B detail.



Figure 10. Male genitalia of *Olceclostera jairana* sp. nov. A ventral view; B lateral view; C aedeagus, lateral view, apex with two spines; D aedeagus, apex with one spine; E aedeagus, apex with three spines.

1.7 mm in diameter and 0.4 mm in height (n = 10); chorion transparent without visible ornamentation; yolk greenish during the first days of development, becoming pale greenish-yellow before eclosion (Fig. 12A–C). The micropylar area bears seven petal-shaped rosette cells of

distinct sizes; around the micropylar rosette, there are ten secondary cells (Fig. 12D); these micropylar cells surround the six openings (Fig. 12E). Aeropyles rounded, organized forming geometric shapes, in this region the egg is quite rough (Fig. 12F, G).



Figure 11. Female genitalia of Olceclostera jairana sp. nov. A ventral view; B dorsal view; C lateral view; D signum detail.

First instar (Fig. 13A, B). Average duration: approximately five days (n = 94); body length 4.8-5.1 mm (n = 3); head capsule width 0.54-0.57 mm (n = 4), hypognathous, slightly flattened in front, yellowish with brownish spots and stripes, stemma 3 larger than the other stemmata, ventral margin of labrum moderately notched; head bears only primary setae (see chaetotaxy below). Body yellow, cylindrical, dorsally with faint black stripes, laterally, with black stripes on subdorsal, lateral, and subventral regions; long setae cover the whole body, except ventrally; dorsally and subdorsally covered with black and yellow setae, laterally and subventrally only with yellow setae, most of them grouped over verrucae; spiracles elliptical, T1 and A8 ones larger than the others; abdominal prolegs fully developed on A3-A6 and A10, with two sclerotized plates, one anterior and one posterior; crochets in biordinal mesoseries. - Chaetotaxy. Head. (Fig. 14) One pair of frontal setae (F1), between them one pair of frontal pores (Fa); two pairs of adfrontal setae (AF1, AF2), between them one pair of adfrontal pores (AFa); three anterior pairs of setae (A1, A2, A3) and one pair of pores (Aa) above A2; two pairs of posterior setae (P1, P2) and two pairs of pores (Pa, Pb); three

pairs of microdorsal setae (MD1, MD2, MD3), and a pair of pores (MDa) between MD2 and MD3; on clypeus, two pairs of setae (C1, C2); on labrum three central (Lrm1, Lrm2, Lrm3) and three lateral pairs of setae (Lrl1, Lrl2, Lrl3). Laterally, one pair of lateral setae (L1) and one pair of pores (La); three stemmatal pairs of setae (S1, S2, S3), and two pairs of stemmatal pore (Sa, Sb); three substemmatal pairs of setae (SS1, SS2, and SS3), and a pair of microgenal pores (MGa). - Body (Fig. 15). Thorax. T1. pronotal plate brownish, weakly sclerotized; XD setae located in the anterior area of the pronotal plate close to four secondary setae; D1 and D2 setae solitary, located in the posterior region of the pronotal plate; SD1 and SD2 setae solitary, SD1 seta at the margin of the pronotal plate; L setae grouped with numerous secondary setae on a conspicuous spherical gravish yellow vertuca; SV setae grouped with numerous secondary setae on a large yellowish verruca; V setae grouped with some secondary setae on a sclerotized gravish plate. T2. D1 seta solitary on a reduced gravish yellow plate, D2 seta grouped with a secondary seta on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow verruca; L setae grouped with numerous secondary



**Figure 12.** Egg of *Olceclostera jairana* **sp. nov.** A three days after oviposition, dorsal view; **B** close to eclosion, lateral view; under scanning electronic microscopy (SEM) **C** dorsal view; **D** rosette cells; **E** micropylar openings; **F** aeropyles; **G** magnification of the aeropyles.

setae on a grayish verruca; SV setae grouped with numerous secondary setae on a grayish verruca; V setae grouped with some secondary setae on a sclerotized grayish plate. T3. D1 seta grouped with a secondary seta on a grayish yellow plate, D2 as D1; SD setae grouped with numerous secondary setae on a grayish verruca; L, SV, and V setae



**Figure 13.** Immature stages of *Olceclostera jairana* **sp. nov. A** first instar, dorsal view; **B** first instar, lateral view; **C** second instar, dorsal view; **D** second instar, lateral view; **E** third instar, dorsal view; **F** third instar, lateral view; **G** fourth instar, dorsal view; **H** fourth instar, lateral view; **I** fifth instar, dorsal view; **J** fifth instar, lateral view.



Figure 14. Head chaetotaxy of first instar of Olceclostera jairana sp. nov. A frontal view; B lateral view.



Figure 15. Body chaetotaxy of first instar of Olceclostera jairana sp. nov. A dorsal view; B lateral view.

as in T2. — *Abdomen*. A1, A2 and A7. D1 seta grouped with two secondary setae on a dark gray plate, D2 seta grouped with six secondary setae on a dark gray verruca; SD setae grouped with eight secondary setae on a dark gray verruca; L setae grouped with seven secondary setae on a dark gray verruca; SV setae grouped with seven secondary setae on dark gray verruca; V seta grouped with a secondary seta on a grayish plate. A3–A6. D1, D2, and L setae as in segments A1, A2, and A7; SD setae grouped

with five secondary setae on a dark gray verruca; SV and V setae on the prolegs on two dark grey plates with some secondary setae. A8. D1 seta grouped with a secondary seta on a grayish yellow plate, D2 seta solitary; SD setae grouped with numerous secondary setae on a grayish yellow verruca; L setae as in the other abdominal segments; SV setae not visible; V seta isolated. A9. D1 and D2 setae grouped with a secondary seta on a grayish yellow plate; SD setae grouped with numerous secondary seta on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a grayish yellow plate; SD setae grouped with numerous secondary setae on a gray setae seta



Figure 16. Pupa of *Olceclostera jairana* sp. nov. A dorsal view;B lateral view; C ventral view;D details of male genital scars;E details of female genital scars.

grayish yellow verruca; SV and V setae not visible. A10. D1, D2, and SD setae arranged on a large grayish-yellow anal plate; L and SV setae on the proleg on two grayish plates with some secondary setae.

**Second instar** (Fig. 13C, D). Average duration: 6–8 days (n = 68); head capsule width 1.1–1.2 mm (n = 5); body length 14.2–14.7 mm (n = 2); head brownish with yellowish spots and stripes; with numerous secondary setae. Body gray with yellow stripes; setae grayish shorter than the first instar; T1–T3 and A8–A10 dorsally darker; between T1 and T2, dorsally with a sparse tuft of reddish setae; A1–A7, D1, and D2 verrucae reddish; legs and abdominal prolegs reddish.

*Third instar* (Fig. 13E, F). Average duration: 6-8 days (n = 30); head capsule width 1.63–1.75 mm (n = 7); body length 21 mm (n = 1); head color as in second instar. Body gray with yellowish stripes, dorsally with darker spots; setae grayish shorter than the second instar; T1 and T2

with setae anteriorly projecting; between T2 and T3, dorsally with a tuft of red setae; A3–A5, and A8 dorsally with a perceptible tuft of dark setae; A9–A10 with longer setae posteriorly projecting; legs and abdominal prolegs red.

*Fourth instar* (Fig. 13G, H). Average duration: 9–11 days (n = 18); head capsule width 2.24–2.41 mm (n = 10); body length 29 mm (n = 1); head color as in third instar. Body gray sprinkled with dark and yellowish spots; T1–T3 bears long gray setae anteriorly projecting, covering the head; between T2 and T3, dorsally with a tuft of bright red setae; A1–A8, dorsally with a tuft of dark setae, longer on A3–A5, and A8; A9 and A10 with longer setae posteriorly projecting; legs and abdominal prolegs dark red.

*Fifth instar* (Fig. 13I, J). Average duration: 9-11 days (n = 15); head capsule width 3.6–3.8 mm (n = 2); body length approximately 45.4–45.9 mm (n = 2); head color as in fourth instar. Body as in the fourth instar, but with smaller setae. Prepupa, average duration: 2-4 days.

**Pupae (Fig. 16).** Average duration: 58–60 days (n = 10); length 21–22 mm (n = 2), obtect, stout, heavily sclerotized, all appendages concealed to the body; antennae broad at the base, tapering rapidly to the apex, reaching two thirds the length of the wings; eyes small; proboscis not visible, wings short with truncated apex; margin of segments A1–A6 notched; spiracles narrowly elliptical; without cremaster or anal hooks. The first imagos emerged 60 days after pupation.

Life history. After being collected, the female was placed in a plastic container where it oviposited 110 eggs on the container wall. After about 15 days, the larvae hatched from the egg by cutting a small semicircular opening on the dorsal surface of the chorion. The larvae preferred to feed on *Handroanthus* sp. (Bignoniaceae). During the first and second instars, larvae fed by scraping both the adaxial and abaxial surfaces of the leaf; from the third on, larvae ate the entire leaf.

During the first and second instars, the larvae were very active all day and night, and when threatened, they released themselves from the host plant to the ground by a silk strand. From the third instar on, the larvae were less active during the day, spending this period resting on the branches of the host plant. In the last instar, when threatened, the larvae bristled the dorsal setae of the thoracic segments, making visible the arrangement of the red setae located in this region.

When larvae reached the prepupa, they left the host plant and walked on the soil of the container, burying in it after a few hours to pupate. The underground chamber was not found. After pupating, the earlier imagos emerged in about 60 days, while the latter in about 120 days. There was no difference between the emergence time of females and males. When both emerged on the same night, they immediately mated. An unmated female oviposited about 130 eggs, while a female that had copulated oviposited about 270 eggs.

**Distribution.** Based on museum records, the species is present in the Argentina state of Entre Ríos and the Brazilian states of Espírito Santo, Minas Gerais, Paraná, Rio de Janeiro, Santa Catarina, and São Paulo (Fig. 22).

**Etymology.** This species is named in honor of the parents of the first author, who always encouraged the first and second authors to persist in their studies; the specific epithet is a combination of their first names, Jair and Ana.

Material examined. *Holotype*. (Fig. 2C, D) BRAZIL. 1 ♂; Parque Estadual do Guartelá, 980 m, Paraná, deposited at the DZUP, with the following labels: / Holotypus/ Parque Estadual do Guartelá, Tibagi, Paraná, Brasil 11–12.X.2018, 980 m, 24°33'59.85"S 50°15'25"W, Carneiro, Venâncio, Moreira and Orlandin leg/ Holotypus *Olceclostera jairana* Orlandin and Piovesan det. 2023/ DZ 43.101/ BC-DZ/ Gen. Prep. E. Orlandin 2021/ DZ 43.101 *Olceclostera jairana* Orlandin and Piovesan det. 2023/.— *Paratypes*. ARGENTINA. *Entre Ríos*: Parque Nacional El Palmar, Intendencia, trampa de luz 26 m, 31°52.126'S 58°12.502'W, E. Núñez Bustos, C. Kopuchian and N. García leg. 2

ර්ථ, MACN-Bar-Lep 01621 (Sample ID)/ MACN-En 24657 (Voucher Museum) and MACN-Bar-Lep 01742/ MACN-En 25200 (MACN). BRAZIL. Espírito Santo: Santa Teresa, Curva do Violão, 690-890 m, 19°59'16.01"S 40°33'3.97"W 1 👌 29-30.VI.2019, D. Dolibaina leg. (DD). Minas Gerais: Aiuruoca 1300 m, 22°00'S 44°38'W 1 👌 10-12.X.2018, Col. Becker 157220 (VOB). Espera Feliz, Sítio Sereno, 1200 m, 20°21'27.78"S 41°51'3.93"W, 3 ♂♂ 11-14.I.2021, D. Dolibaina leg. (DD). Paraná: Adrianópolis, Parque Estadual das Lauráceas 24°48'45.90"S, 48°41'9.54"W 1 👌 18-20.IV.2018, O. Mielke, Casagrande, Carneiro, Piovesan, Dantas and Santos leg./ DZ 45.719/ Gen. Prep. E. Orlandin 2021/ DZ 45.719 Olceclostera jairana Orlandin and Piovesan det. 2023; 1 🖑 18-20.IV.2018, O. Mielke, Casagrande, Carneiro, Piovesan, Dantas and Santos leg./ DZ 45.813/ Gen. Prep. E. Orlandin 2021/ DZ 45.813 Olceclostera jairana Orlandin and Piovesan det. 2023; 1 👌 26-29.X.2019, Orlandin E. and Cafisso C. leg./ DZ 52.855/ Gen. Prep. E. Orlandin 2021/ DZ 52.855 Olceclostera jairana Orlandin and Piovesan det. 2023; 1 3 26-29.X.2019, Orlandin E. and Cafisso C. leg./ DZ 52.856 (DZUP). Campina Grande do Sul, 897 m, 25°15'11.1"S 48°52'59.14"W 2 🖑 🖓 10.XI.2017 (CDH). Guaratuba, 622 m, 25°49'55.30"S 48°55'48.51"W 1 🖑 20.X.2011/ genitalia prep. D. Herbin, ref H. 966; 840 m, 25°51'47.55"S 48°55'01.87"W 1 👌 12.XI.2017. Ponta Grossa, Parque Estadual de Vila Velha, 1.060 m, 25°14'34"S, 50°0'14"W 1 👌 23-25.I.2020, E. Orlandin and C. Cafisso leg./ BC-DZ/ DZ 40.950; 1 👌 23-25.I.2020, E. Orlandin and C. Cafisso leg./ DZ 52.863; 1  $\stackrel{\scriptstyle <}{\scriptstyle \sim}$  14–20.XII.2020, E. Orlandin, M. Piovesan and C. Cafisso leg./ DZ 52.869/ Gen. Prep. E. Orlandin 2021/ DZ 52.869 Olceclostera jairana Orlandin and Piovesan det. 2023/ (DZUP). São José dos Pinhais, Chácara Corta-Vento, 970 m, 25°42'09.3"S, 49°04'19.3"W 1 👌 24-25.II.2020, E. Orlandin and C. Cafisso leg./ BC-DZ/ DZ 52.186; 1 🖑 16.XI.2020, E. Orlandin, C. Cafisso and M. Piovesan leg./ DZ 53.021/ Gen. Prep. E. Orlandin 2021/ DZ 53.021 J Olceclostera jairana Orlandin and Piovesan det. 2023; 1 🖑 17-18.IX.2020, E. Orlandin, C. Cafisso and M. Piovesan leg./ DZ 52.317 (DZUP). Tibagi, Parque Estadual do Guartelá, 975 m, 24°33'59"S, 50°15'26"W 1 ් 18.I.2012/ genitalia prep. D. Herbin, ref H. 1652; 4 ඊට් 18.I.2012 (CDH); 950-1000 m, 24°33'44"S, 50°15'33"W 1 🖑 25-27.VII.2017, Dantas, Dias, Dolibaina and Queiroz-S Leg./ DZ 45.810/ Gen. Prep. E. Orlandin 2021/ DZ 45.810 Olceclostera jairana Orlandin and Piovesan det. 2023; 1 ♀ 25-27.VII.2017, Dantas, Dias, Dolibaina and Queiroz-S Leg./ DZ 45.742 (DZUP). Rio de Janeiro: Cachoeiras de Macacu, Valério 1 👌 10.XII.2011, N. Tangerini col./ Ex. Coleção Nirton Tangerini/ DZ 45.752/ Gen. Prep. E. Orlandin 2021/ DZ 45.752 Olceclostera jairana Orlandin and Piovesan det. 2023 (DZUP). Itatiaia, Parque Nacional do Itatiaia, 1000 m, 22°26'55.9"S 44°36'39.5"W 1 🖒 10-20.I.2021, A.H.B Rosa leg./ DZ 52.287/ Gen. Prep. E. Orlandin 2021/ DZ 52.287 Olceclostera jairana Orlandin and Piovesan det. 2023 (DZUP); 910 m 1 3 4-8.XII.2019, A. Soares, G. Marconato, M. A. Costa and N. Tangerini col., COL MNRJ H/N° 0601108, MN-LEP 0001686, 807 m, 22°27'16"S 44°36'29"W; 2 ♂♂ 7-8.IV.2021, T. Zacca leg./ MN-LEP 0002166 and MN-LEP 0002264/ T. Zacca prep. 2021, DNA-LAPEL 0000000243 and DNA-LAPEL 0000000280 (MNRJ). Petrópolis, 1 🖑 6.VI.1959, Gagarin leg./ Ex. Col. Gagarin/ DZ 45.722/ Gen. Prep. E. Orlandin 2018/ DZ 45.722 Olceclostera jairana Orlandin and Piovesan det. 2023 (DZUP). Santa Catarina: Joaçaba, Nova Petrópolis, 811 m, 27°6'20.09"S, 51°37'14.83"W 1 ♂ 02.IV.2020, E. Orlandin leg., Ex. Larvae, DZ 52.191/ BC-DZ; 1 Q (Fig. 4A, B) 02.VI.2020, E. Orlandin leg., Ex. Larvae, DZ 52.194/ BC-DZ/ Gen. Prep. E. Orlandin 2021/ DZ 52.194♀ 1 ♀ 20.VII.2020, E. Orlandin leg., Ex. Larvae, DZ 52.182; 800 m, 27°06'20.6"S 51°37'15.3"W 1 🖒 02.IV.2020, A.H. Schneeberger, V.O. Souza and E. Orlandin leg./ DZ

53.111/ Gen. Prep. E. Orlandin 2021/ DZ 53.111 Olceclostera jairana Orlandin and Piovesan det. 2023 (DZUP). Immature stages: 001.1-141 eggs 26 first instar; 001.2 - 3 second instar; 001.3 - 1 third instar; 001.4 -1 fourth instar; 001.5 - 2 fifth instar 2 pupae; 001.6 - 4 first instar head capsule; 001.7 - 6 second instar head capsule; 001.8 - 11 third instar head capsule; 001.9 - 11 fourth instar head capsule; 001.10 - 2 fifth instar head capsule; 001.11 - 2 pupa exúvia. PARATYPUS, Olceclostera jairana, XII.2019-IV.2020, Nova Petrópolis, Joaçaba, Santa Catarina, Brasil, 811 m, 27°6'20"S 51°37'14"W, E. Orlandin leg. / PARATI-PO / DZUPIL 0170 (DZUPIL). São Bento do Sul, Rio Vermelho, 3 ්ථ, .X.2004; 503 m, 26°20'00.77"S, 49°18'28.25"W 1 ් 23.X.2011/ genitalia prep. D. Herbin, ref H. 970; 9 33 23.X.2011 (CDH). São Paulo: Caconde, 21°28'45"S, 46°35'34"W 1 👌 10-14.I.2018, L. Venancio leg./ DZ 52.943/ Gen. Prep. E. Orlandin 2021/ DZ 52.943 3/Olceclostera jairana Orlandin and Piovesan det. 2023 (DZUP). Jundiaí, Japi, 23°13'S 46°56'W, 900 m 1 👌 13-16.II.2001, V.O. Becker col., Col. Becker 131737 (VOB). Ribeirão Grande, Intervales, 24°16'12"S 48°24'36"W, 1000 m 1 3, 06-07.XI.2021, V.O. Becker col., Col. Becker 165410 (VOB).

**Remarks.** We found at the same locality specimens with one, two, or three spines at the apex of the aedeagus. However, specimens with a different number of spines present a molecular difference of less than 2% (Table 1).

#### 3.1.2. Olceclostera quilombola Orlandin, Piovesan and Carneiro sp. nov.

https://doi.org/zoobank.org/BAA7EC5F-9C6C-4BEB-B466-135E7316F719

Figures 2E, F, 4C, D, 17, 18

**Type locality.** Parque Estadual das Lauráceas, Adrianópolis Municipality, Paraná State, Brazil.

Diagnosis. Olceclostera quilombola sp. nov. differs from O. jairana sp. nov. by the following characters: on the forewing of O. quilombola sp. nov. the two subapical hyaline spots between M₁ and M₃ are rectangular and always visible, while these spots are rounded, with the one between M₂-M₃ almost unnoticeable in some cases in O. jairana sp. nov. (Figs 2C, D, 4A, B). Olceclostera qui*lombola* **sp. nov.** has all tibias with a band of long scales with dark brown apex, while O. jairana sp. nov. presents these scales with light brown apex (Fig. 6A, B). In the genitalia, O. quilombola sp. nov. presents, in lateral view, a more flat tegumen, wider socii, and a less elongated uncus, and, in ventral view, the anterior projection of the saccus is wider than O. jairana sp. nov. The aedeagus of O. quilombola sp. nov. is thinner than that of O. jairana sp. nov. and does not have spines at the apex (Figs 10, 17).

Olceclostera quilombola **sp. nov.** differs from O. xeta **sp. nov.** and O. wayana **sp. nov.** by external and genitalia characters: on the forewing of O. quilombola **sp. nov.**, the two subapical hyaline spots, between  $M_1$  and  $M_3$ , are rectangular and always visible, while in O. xeta **sp. nov.** and O. wayana **sp. nov.**, these spots are rounded and shorter. Olceclostera quilombola sp. nov. has all tibias with a band of long scales with dark brown apex, while O. xeta sp. nov. and O. wayana sp. nov. present these scales with grayish brown apex (Fig. 6B, C). Dorsally, the abdomen of O. quilombola sp. nov. has a copper color, while O. xeta sp. nov. and O. wayana sp. nov. have a grayish color (Fig. 2E-J). The genitalia of O. quilombola sp. nov. presents, in lateral view, the socci apex narrower and triangular, while O. xeta sp. nov. and O. wayana sp. nov. have the socci apex rounded. The process of sacculus of O. quilombola sp. nov. is about twice as big as that of O. xeta sp. nov. and O. wayana sp. nov. The aedeagus of O. quilombola sp. nov. has vesica without cornuti, while in O. xeta sp. nov. and O. wayana sp. nov. the basal portion of the vesica is covered by small cornuti (Figs 17, 19, 21).

Description (Figs 2E, F, 4C, D, 6B, 17, 18). Head. MALE. As in O. jairana sp. nov. Female. Similar to the male, slightly smaller than the male's head; antenna, pecten shorter than the male's pecten; palpus, thinner and shorter than the male's palpus. — Thorax. MALE and FEMALE. As in O. jairana sp. nov. — Legs. As in O. jairana sp. nov., but all tibias with a band of long scales with dark brown apex projecting towards the posterior region (Fig. 6B). — Wings shape and venation. As in O. jairana sp. nov. (Figs 2E, F, 4C, D). MALE. Forewing length: 18-20.5 mm (n = 10). Upperside: As in O. jairana sp. nov., but ground color gray; subapical region with two rectangular hyaline spots between  $M_1$  and  $M_3$ ; outer margin crenulated. Underside: Similar to O. jairana sp. nov. Hindwing. Upperside and underside: Similar to O. jairana sp. nov., but the outer margin crenulated between M₃ and CuA₂. **FEMALE.** Forewing length: 25.8 mm (n = 01); similar to O. *jairana* sp. nov.; general color similar to the male, the apex is more acute than in males; on dcm, the patch of black scales and the hyaline spots are broader than in the male. Hindwing. Similar to the male, but on dcm, the dot of black scales is broader than in the male. — Abdomen. MALE and FEMALE. As in O. jairana sp. nov. — Genitalia. MALE (Fig. 17). Similar to O. jairana sp. nov., but the aedeagus is thinner and does not have spines at the apex. FEMALE (Fig. 18). As in O. *jairana* sp. nov., but the eighth tergum and the ostium bursae are narrower.

**Distribution.** Based on museum records, the species is present in the Brazilian states of Brasília, Espírito Santo, Minas Gerais, Paraná, Rio de Janeiro, and São Paulo (Fig. 22).

**Etymology.** This species is named in honor of the quilombola communities that inhabit the region of the "Parque Estadual das Lauráceas" the type locality of this species.

Material examined. *Holotype*. (Fig. 2E, F) BRAZIL. 1 ♂; Parque Estadual das Lauráceas, Adrianópolis, 900 m, Paraná, Brazil, deposited at the DZUP, with the following labels: / Holotypus/ P[ar]q[ue] Est[adual] das Lauráceas, Adrianópolis, Paraná, Brasil 26–29.X.2019,



Figure 17. Male genitalia of Olceclostera quilombola sp. nov. A ventral view; B lateral view; C aedeagus lateral view.



Figure 18. Female genitalia of Olceclostera quilombola sp. nov. A ventral view; B dorsal view; C lateral view; D signum detail.

900m 24°48'45.90"S, 48°41'9.54"W, Orlandin E. and Cafisso C. leg./ Holotypus Olceclostera quilombola Orlandin, Piovesan and Carneiro det. 2023/ DZ 51.895/ BC-DZ/ Gen. Prep. E. Orlandin 2021/ DZ 51.895 Olceclostera quilombola Orlandin, Piovesan and Carneiro det. 2023/.- Paratypes. BRAZIL. Brasília: Planaltina, 1000 m, 15°35'S 47°42'W 1 👌 15.V.1982, V.O. Becker col., Col. Becker 40190 (VOB). Espírito Santo: 1 👌 2741, CUIC 000060413 (CUIC). Santa Teresa, Reserva Biológica de Augusto Ruschi, 839 m, 19°54'19"S 40°34'07"W 1 👌 13–16.XI.2012; 841 m, 19°54'50"S 40°31'58"W 1 👌 15.XI.2012/ genitalia prep D. Herbin, ref H. 1651; 19°53'21"S 40°32'41"W 1 🖒 16.XI.2012 (CDH). Minas Gerais: Alto Caparaó, Parque Nacional do Caparaó, 1309 m, 20°25'09"S 41°51'04"W 1 👌 11.XI.2012/ Gen. Prep. D. Herbin H1658 (CDH). Caraca, 1300 m 1 & 07-10.V.1996, Becker col., Col. Becker 112800 1 👌 19.I.1998, V.O. Becker col., Col. Becker 112710 (VOB). Espera Feliz, Sítio Sereno, 1200 m, 20°21'27.78"S 41°51'3.93"W 2 ♂♂ 11–14.I.2021, D. Dolibaina leg. (DD). Itamonte, Pousada Fazenda Serra Bonita, 1400 m, 22°14'34.7"S 44°43'11.9"W 1 👌 08-11.I.2021, A. H. B. Rosa leg./ DZ 52.292/ Gen. Prep. E. Orlandin 2021/ DZ 52.292 Olceclostera quilombola Orlandin, Piovesan and Carneiro det. 2023 (DZUP). Paraná: Adrianópolis, 900 m, 24°48'45.90"S, 48°41'9.54"W 1 🖑 26–29.X.2019, Orlandin E. and Cafisso C. leg./ DZ 52.854 (DZUP). Curitiba, 920 m, 1 3 20.XII.1974, V.O. Becker col., Col. Becker N°563 1 & 13.IX.1974, V.O. Becker col., Col. Becker N°564 (VOB). Guaratuba, 600 m, 1 &, 05. VIII. 1975, V.O. Becker col., Col. Becker N°529 (VOB). São José dos Pinhais, Chácara Corta-Vento, 970 m, 25°42'09.3"S, 49°04'19.3"W 1 ♂ 29.I.2020, Orlandin E., Cafisso C. and Carneiro E. leg./ DZ 47.920/ BC-DZ; 1 3 29.I.2020, Orlandin E., Cafisso C. and Carneiro E. leg./ DZ 52.857; 1 3 25-26.II.2020, E. Orlandin and C. Cafisso leg./ DZ 53.030/ Gen. Prep. E. Orlandin 2021/ DZ 53.030 Olceclostera quilombola Orlandin, Piovesan and Carneiro det. 2023; 2 3 25-26.II.2020, E. Orlandin and C. Cafisso leg., DZ 52.942 and DZ 53.029; 1 3 17-18.IX.2020, E. Orlandin C. Cafisso and M. Piovesan leg./ DZ 52.320; 1 3 16.XI.2020, E. Orlandin C. Cafisso and M. Piovesan leg./ DZ 53.020 (DZUP). Rio de Janeiro: Itatiaia, Parque Nacional do Itatiaia, Cascata de Maromba, 1177 m, 22°25'45"S 44°37'08"W 1 👌 6.XI.2012/ genitalia prep D. Herbin, ref H. 1659 (CDH). Nova Friburgo, 1100 m 1 🗇 21.I.1998, V.O. Becker col., Col. Becker 10796 1 3 10.X.1998, V.O. Becker col., Col. Becker 117489 (VOB). Petrópolis, 1 👌 16.X.1955, D'Almeida col./ DZ 45.718/ Gen. Prep. E. Orlandin 2018/ DZ 45.718 Olceclostera quilombola Orlandin, Piovesan and Carneiro det. 2023; 1 👌 30.X.1953, D'Almeida and Berla col./ DZ 45.748 (DZUP). Teresópolis, 1000 m 1 👌 15.I.1985, V.O. Becker col., Col. Becker 54762, Genitália 2644 (VOB). São Paulo: Apiaí, 750 m 17 ♂♂ 15.VII.2005; 4 ♂♂ 12.X.2006; 4 3 2.XI.2007, Coll. D. Herbin (CDH). Campos do Jordão, 1600m, 22°46'S 45°31'W 1 👌 23-27.I.2001, V.O. Becker col., Col. Becker 130775 (VOB). Capão Bonito, Intervales, 900 m, 24°16'S 48°24'W 1 3, 05-07.XII.2001, V.O. Becker col., Col. Becker 134144 (VOB). Guapiara, Paivinha, 800 m 2 3 3 6-8.III.2005; 2 3 3 2-5.IV.2005; 5 ් ටී 25–27.VII.2005; 3 ් ටී 16–19.IX.2005; 1 ් 23–25.X.2005; 3 ් ් 14-16.II.2006, Coll. D. Herbin (CDH). Guaratinguetá, S[anta] Maria da Serra, 1900 m, 22°25'S 45°25'W, 3 3 3 24-26.III.2022, V.O. Becker col., Col. Becker 165753 (VOB). Salesópolis, [Estação Biológica de] Boracéia 1 👌 05.IX.1942, D'Almeida col./ DZ 45.789; 1 👌 27.IX.1948, D'Almeida col./ DZ 45.830 (DZUP). São José do Barreiro, 1640 m, 22°43'26"S 44°36'57"W 2 🖑 .X.2021, V.O. Becker col., Col. Becker 164501 (VOB). São Luís do Paraitinga, 900 m, 23°20'S 45°06'W 1 3 12-17.XI.2001, V.O. Becker col., Col. Becker 133834, 700 m, 23°20'S 39°03'W 1 ♀ (Fig. 4C, D) 26.X.2019, V.O. Becker col., Col. Becker 160738 (VOB).

# 3.1.3. Olceclostera xeta Orlandin, Herbin and Carneiro sp. nov.

https://zoobank.org:/D73ED286-6BC6-4D7E-B9B4-C2141A08DF50

Figures 2G, H, 4E, F, 19, 20

**Type locality.** RPPN Fazenda da Mata, Querência do Norte Municipality, Paraná State, Brazil.

**Diagnosis.** Externally, *O. xeta* **sp. nov.** can be distinguished from *O. jairana* **sp. nov.** by all tibias with a band of long scales with grayish brown apex, while *O. jairana* **sp. nov.** presents these scales with light brown apex (Fig. 6A, C). Dorsally, the abdomen of *O. xeta* **sp. nov.** has a grayish color, while *O. jairana* **sp. nov.** has a copper color (Fig. 2C, D, G, H). The genitalia of *O. xeta* **sp. nov.** presents, in lateral view, a wider socci than *O. jairana* **sp. nov.**, and the process of sacculus is half the length of that of *O. jairana* **sp. nov.** The aedeagus of *O. xeta* **sp. nov.**, does not have spines, and the basal portion of the vesica is covered by small cornuti (Figs 10, 19).

Olceclostera xeta sp. nov. differs from O. quilombola sp. nov. by external and genitalia characters: on the forewing of O. xeta sp. nov. the two subapical hyaline spots, between  $M_1$  and  $M_3$ , are rounded and smaller. Otherwise, these spots are rectangular in O. quilombola sp. nov. Dorsally, the abdomen of O. xeta sp. nov. has a grayish color, while O. quilombola sp. nov. has a copper color (Fig. 2E-H). Olceclostera xeta sp. nov. has all tibias with a band of long scales with gravish brown apex, while O. quilombola sp. nov. presents these scales with dark brown apex (Fig. 6B, C). The genitalia of O. xeta sp. nov. presents the process of sacculus with half of the length of O. quilombola sp. nov., and the basal portion of the vesica is covered by small cornuti, while the aedeagus of O. quilombola sp. nov. has vesica without cornuti (Figs 17, 19).

Olceclostera xeta **sp. nov.** is similar to O. wayana **sp. nov.**, but slightly darker gray and smaller in size (17 mm forewing length). The subapical hyaline spots are slightly larger. In the genitalia, O. xeta **sp. nov.** differs from O. wayana **sp. nov.** by the presence of a protrusion on the apex of the aedeagus.

**Description (Figs 2G, H, 4E, F, 6C, 19, 20).** *Head.* **MALE.** As in *O. jairana* **sp. nov. FEMALE.** Antenna, pecten shorter than the male's pecten. — *Thorax.* **MALE and FEMALE.** As in *O. jairana* **sp. nov.** — *Legs.* All tibias with a band of long scales with grayish-brown apex projecting towards the posterior region (Fig. 6C). — *Wings shape and venation.* **MALE and FEMALE.** As in *O. jairana* **sp. nov.** (Figs 2G, H, 4E, F). **MALE.** Forewing length: 16.5–17 mm (n = 03). Upperside and underside: Similar to *O. jairana* **sp. nov.** Hindwing. Upperside and underside: Similar to *O. jairana* **sp. nov. FEMALE.** Forewing length: 17–21 mm (n = 03); general color similar to the male, the apex is more acute than in males; lighter grey than the female of *O. jairana* 



Figure 19. Male genitalia of Olceclostera xeta sp. nov. A ventral view; B lateral view; C aedeagus lateral view; D cornuti detail.



Figure 20. Female genitalia of Olceclostera xeta sp. nov. A latero-ventral view; B dorsal view; C signum detail.

**sp. nov.** and *O. quilombola* **sp. nov.** The hyaline spot between  $M_1-M_2$  is small than in males, and the hyaline spot between  $M_2-M_3$  is reduced or absent. Hindwing. As in male. *Abdomen.* **MALE and FEMALE.** Similar to *O. jairana* **sp. nov.**, but dorsally gray. — *Genitalia.* **MALE.** (Fig. 19). Similar to *O. jairana* **sp. nov.**, but aedeagus without spines; vesica with the basal portion covered by small cornuti. **FEMALE.** (Fig. 20). Similar to *O. jairana* **sp. nov.**, but the ostium bursae are broader and the signum leaf-shaped.

**Distribution.** Based on museum records, the species is present in the Brazilian states of Paraná and the Para-

guayan provinces of Alto Paraná, Canindeyú, Concepción, Cordillera, Hayes, and Paraguarí (Fig. 22).

**Etymology.** This species is named after the Xetá indigenous people. The Xetá were original inhabitants of the northwestern Paraná and the last ethnic group in the state to come into contact with white colonizers (in 1954) who invaded their territory and decimated the Xetá population.

Material examined. Holotype. (Fig. 2G, H) BRAZIL. 1 ♂; RPPN Fazenda da Mata, Querência do Norte 280 m, Paraná, Brazil, deposited at the DZUP, with the following labels: / Holotypus/ Brasil, Paraná, Querência do Norte, RPPN Faz[enda] da Mata 280 m, 13.IX.2020 22°56'15"S 53°32'05"W, Casagrande, Mielke, Carneiro and Orlandin leg./ Holotypus Olceclostera xeta Orlandin, Herbin and Carneiro det. 2023/ DZ 52.940/ BC-DZ./.- Paratypes. BRAZIL. Paraná: Querência do Norte, RPPN Fazenda da Mata 280 m, 22°56'15"S 53°32'05"W 2 3 3 13.IX.2020, Casagrande, Mielke, Carneiro and Orlandin leg./ DZ 43.365 and DZ 52.158/ Gen. Prep. E. Orlandin 2018 and Gen. Prep. E. Orlandin 2021/ DZ 43.365 and DZ 52.158 Olceclostera xeta Orlandin, Herbin and Carneiro det. 2023 (DZUP). PARAGUAY. Cordillera: Takuara Renda, 25°39'S 56°55'W 14 ♂♂ 1 ♀ 05–08.VIII.2011, 3 3 3 10-14.XI.2011, Leg. U. Drechsel, Coll. D. Herbin (CDH). Pirareta, 25°29'S 56°56'W 1 👌 28.VII.2010; 1 👌 14–18.VII.2011; 3 👌 24-28.VII.2011; 3 3 3 31.VIII.2011, Leg. U. Drechsel, Coll. D. Herbin (CDH). Puerto Naranjahay, 62 m, 24°58'S 57°12'W 1 👌 1.VI.2015, Leg. U. Drechsel, Coll. D. Herbin (CDH). Alto Paraná: Estancia Dimas, 25°33'S 55°13'W 2 3 3 19–22.VII.2012; 2 3 3 6–8.VII.2013; 1 ∂ 10–12.III.2011, Leg. U. Drechsel, Coll. D. Herbin 1♀ (Fig. 4E, F), Leg. U. Drechsel/ Gen. Prep. D. Herbin H1660/ (CDH). Limoy, 245 m, 24°45'S 54°27'W 1 👌 6-8.III.2014, Leg. U. Drechsel, Coll. D. Herbin (CDH). Hayes: Chaco Lodge, 115 m, 22°30'S 59°18'W 1 ♂ 1 ♀ 4–6. XI.2015, Leg. U. Drechsel, Coll. D. Herbin (CDH). Concepción: San Luis, 170 m, 22°24'S 57°28'W 2 33 12-14.V.2016, Leg. U. Drechsel, Coll. D. Herbin (CDH). Canindeyú: Mbaracayú, 194 m, 24°08'S 55°31'W, 3 38 27-29.XI.2013, Leg. U. Drechsel, Coll. D. Herbin (CDH). Carapá, 240 m, 24°22'S 54°23'W 2 ♂♂ 1-4.IV.2009, Leg. U. Drechsel (CDH). **Paraguarí: Mbatoví**, 383 m, 25°35'S 57°05'W 2 33 17–18.IV.2014, Leg. U. Drechsel, Coll. D. Herbin/ BC-Her 4924 and BC-Her4695 (CDH).

# 3.1.4. Olceclostera wayana Herbin, Orlandin and Carneiro sp. nov.

https://zoobank.org/6B8921EC-5000-42F4-B471-61EB664978E4

Figures 2I, J, 21

Type locality. Piste Patagai-Counamama, French Guiana.

**Diagnosis.** Externally, *O. wayana* **sp. nov.** is similar to *O. xeta* **sp. nov.**, slightly lighter grey and larger (21 mm forewing length). The two subapical hyaline spots between  $M_1$  and  $M_3$  are tiny. In the genitalia, *O. wayana* **sp. nov.** differs from *O. xeta* **sp. nov.** by the absence of a protrusion on the apex of the aedeagus and vesica with a plate with minute teeth.

**Description (Figs 21, J, 21).** *Head.* MALE. As in *O. xeta* **sp. nov. FEMALE.** Unknown. — *Thorax.*—As in *O. xeta* **sp. nov.** — *Legs.* As in *O. xeta* **sp. nov.** — *Wings shape and venation.* As in *O. jairana* **sp. nov.** (Fig. 21, J). Forewing length: 21 mm (n = 04). Upperside and underside: Similar to *O. xeta* **sp. nov.**, slightly lighter grey. The subapical hyaline spots between  $M_1$  and  $M_3$  are tiny. Hindwing. Upperside and underside: Similar to *O. xeta* **sp. nov.** — *Genitalia.* (Fig. 21). Similar to *O. xeta* **sp. nov.**, but bigger, without a protrusion on the apex of the aedeagus and vesica with a plate with minute teeth.

#### Distribution. French Guiana.



Figure 21. Male genitalia of *Olceclostera wayana* sp. nov. A ventral view; B lateral view; C aedeagus lateral view; D detail of vesica with a plate with minute teeth.



Figure 22. Geographical distribution of new species of Olceclostera.

**Etymology.** This species is named after the Wayana indigenous people, inhabiting along the Maroni River, at the border between French Guiana and Suriname.

Material examined. *Holotype*. (Fig. 2I, J) FRENCH GUIANA. 1 ♂; French Guiana, deposited in MNHN, with the following labels: / Holotype, *Olceclostera wayana*, des. Herbin, Orlandin and Carneiro 2023 / Guyane Française, Piste Patagai-Counamama, 05°22'59.51"N 53°12'27.25"W, Alt. 49 m 23.XI.2013, leg. D. Herbin / genitalia prep. D. Herbin ref. H.1166 / BC-Her4834 / CDH 3.042.—*Paratypes*. FRENCH GUIANA. **Piste Paul Isnard**, PK 34 1 ♂ 28.VII.2001, leg. M. Laguerre, Coll. D. Herbin, genitalia prep. D. Herbin ref. H.1702 (CDH). Citron 1 ♂ 11–17.III.1986, leg. G. Tavakilian, Coll. D. Herbin (CDH). **Piste Patagai-Counamama**, 05°20'34.23"N 53°12'47.86"O, Alt. 58m 1 ♂ 02.XI.2013, leg. O. Felis, Coll. D. Herbin, genitalia prep. D. Herbin ref. H.1703 (CDH).

#### 3.2. Maximum Likelihood Tree

Based on sequences of the mitochondrial COX1 gene (barcode region, 658pb), our reconstructed tree recovers *O. jairana* **sp. nov.** and *O. quilombola* **sp. nov.** as a sister group to *O. xeta* **sp. nov.** and *O. wayana* **sp. nov.**, although this relationship is not well supported. This clade is sister to a cluster that includes the Central American *O. bifenestrata* and the North American *O. angelica*, *O. indistincta*, and *O. seraphica* (Fig. 23). This relationship,

however, shows low support. The topology obtained between the *Olceclostera* new species supports their distinctiveness, whose genetic distances ranged from 3.4% to 10% (Table 1).

#### 4. Discussion

Olceclostera jairana sp. nov., O. quilombola sp. nov., O. *xeta* **sp. nov.** and *O. wayana* **sp.nov.** are morphologically similar to the North American species O. angelica, O. indistincta, and O. seraphica (see Franclemont 1973). The four new species are cryptic with the Central American species O. bifenestrata and are identified as this species in the collections and in the few studies in which it was cited (Draudt 1929; Biezanko 1986). In fact, the new South American species cluster together with O. bifenestrata, O. angelica, O. indistincta, and O. seraphica, thus forming a monophyletic group well differentiated from the other 16 species of the genus analyzed and the outgroup. Although our COX1 tree showed a low support value for this clade, there are morphological characters that corroborated this group. These eight species share traits such as the presence of two hyaline spots on the forewings and only one pair of large patches of broader scales laterally on the abdomen. Hyaline spots on the



Figure 23. Maximum likelihood phylogenetic tree, inferred with IQ-TREE, based on cytochrome c oxidase subunit I (COX1, barcode region, 658pb) showing the relationship among new species of *Olceclostera*, twenty other species of *Olceclostera*, and representatives of ten genera of Apatelodidae as outgroups. Numbers indicate  $ML \ge 70$ , and * indicates maximum support of 100. Low support values were not shown. The scale bar represents the expected number of nucleotide substitutions per site.

forewings and patches of broader scales on the abdomen are also present in the other *Olceclostera* species but with different configurations. These characters, together with characters of the genitalia (uncus bifid V-shaped, gnathos converted to sclerotized plates, and subscaphium sclerotized), are proposed as diagnostic of the genus.

The configuration of the hyaline spots and the number of patches of broader scales laterally on the abdomen can be used to distinguish groups of species of *Olceclostera*. For example, some species have only a yellowish hyaline spot on their forewings and one pair of large patches of broader scales laterally on each abdominal segment (e. g. *Olceclostera guanduna* Drautd, 1929, *Olceclostera in*- dentata Schaus, 1910, O. irrorata, Olceclostera leticiensis Herbin, 2021, Olceclostera microps (Walker, 1855) and Olceclostera reperta (Walker, 1865)), while others have only a whitish hyaline spot on their forewings and two pairs of large patches of broader scales laterally on the proximal abdominal segments (e. g. Olceclostera amoria Druce, 1890, Olceclostera azteca Schaus, 1894, Olceclostera magniplaga Schaus, 1910 and Olceclostera porioni Herbin and Mielke, 2018). Although the patches of larger scales on the abdomen call us attention, these patches are mentioned in the descriptions of only seven species (O. angelica, O. irrorata, O. azteca, O. magniplaga, O. reperta, O. indentata, O. microps). However, these spots are not present in all the species of the genus. Therefore the phylogenetic significance of this character requires further studies.

Knowledge about immature traits is essential to understanding the Lepidoptera evolution (Balcázar-Lara and Wolfe 1997; Freitas and Brown 2004). In this study, we describe for the first time the morphology of all immature stages of one Olceclostera species and illustrate for the first time images of the egg of an Apatelodidae species using SEM. In dorsal view, the egg of O. jairana sp. nov. is similar to that described for O. seraphica, Apatelodes, and Thelosia species (Packard 1895; Peigler 1994; Thomazini 2010; Herbin 2020; Orlandin et al. 2022a, 2022b) appearing to be flattened as cited for these species. However, the lateral view of the eggs of O. jairana sp. nov. shows that the micropylar pole is concave (pie-shaped) and not flattened (coin-shaped). This finding is important because it demonstrates that there are a greater variety of egg shapes in Apatelodidae than previously thought since we found dome-shaped eggs in Drepatelodes (Orlandin et al. 2021) and coin-shaped eggs in Apatelodes (Packard 1895; Orlandin et al. 2022a, 2022b). To infer the evolutionary significance of the egg morphology in Apatelodidae, eggs of more species of the different genera should be obtained, and these eggs must be studied in detail (e.g. using SEM), including records in lateral view.

The chaetotaxy of O. jairana sp. nov. is very similar to the chaetotaxy of other Apatelodidae species, which were described in detail (Orlandin et al. 2021, 2022a, 2022b). The head shows only primary setae, as the other species already studied. On the body, the known species have secondary setae, mostly arranged on large verrucae. However, the most striking difference in the chaetotaxy of O. jairana sp. nov. is the lack of L2 on A1-A8 isolated, while in Apatelodes je Orlandin, Piovesan & Carneiro, 2022, Apatelodes kotzschi Draudt, 1929, and D. friburgensis, L2 is isolated (Orlandin et al. 2021, 2022a, 2022b). Additionally, first instar larvae of Apatelodidae also seem to present a great variation in the number of secondary setae grouped with D1 and D2 on the thorax and the abdomen. Olceclostera jairana sp. nov. has on T2-T3 segments, D1 seta solitary and D2 seta arranged with only one secondary seta, while A. je also has D1 seta solitary but D2 seta on a verruca with about six secondary setae (Orlandin et al. 2022a). On the other hand, in A. kotzschi, D1 seta is on a verruca with three secondary setae, and D2 seta is on a verruca with about six secondary setae (Orlandin et al. 2022b), while D. friburgensis has D1 seta on a verruca with two or three secondary setae, and D2 seta next to a secondary seta (Orlandin et al. 2021). Between A1-A7, O. jairana sp. nov. presents D1 seta on a verruca with two more secondary setae and D2 seta on a verruca with about six secondary setae. Unlike this configuration, A. je and A. kotzschi have both D1 and D2 setae on each verruca with five to seven secondary setae, and D. friburgensis has D1 seta on a vertuca with about five secondary setae and D2 seta solitary (Orlandin et al. 2021, 2022a, 2022b). In A8, the configuration of dorsal setae is very similar, and all species have the D1 seta with one to three secondary setae and solitary D2 seta. Here we emphasize that the chaetotaxy of Apatelodiade still requires a broader study to evaluate its diagnostics and phylogenetic relevance to delimit monophyletic groups in the family.

The available information on the immature stages of Olceclostera allows us to make some comparisons between the species of the genus. In the last larval instars, O. jairana sp. nov. is very similar to O. bifenestrata and O. seraphica, differing from them by the more brownish color, by the dorsal region of the body with smaller yellow spots (larger in O. bifenestrata) and by the absence of orange spots on the dorsal region of the body (present in O. seraphica) (Peigler 1994; Janzen and Hallwachs 2009). The color pattern, the slightly flattened dorsoventrally body, and the pattern in the setae distribution make these species very similar to larvae of some species of Euglyphis (Lasiocampidae) (Janzen and Hallwachs 2009). Euglyphis caterpillars are known for having urticating setae (Specht et al. 2008), absent in the Olceclostera genus, which may constitute a Batesian-type mimicry. In Apatelodidae, this type of mimicry seems to be common, with caterpillars mimicking larvae of Megalopygidae, Arctiinae, and Lasiocampidae (Orlandin et al. 2021, 2022a).

Olceclostera jairana **sp. nov.** pupates in the soil like other known Apatelodidae species. This trait appears to be shared by most species in the family, as, so far, only *Prothysana* is known to make cocoons (Becker and Orlandin in press). The pupa of *O. jairana* **sp. nov.** is similar to that described for *O. seraphica* and to the other species of Apatelodidae (Peigler 1994; Janzen and Hallwachs 2009; Thomazini 2010; Orlandin et al. 2021, 2022a, 2022b). However, the absence of a rudimentary cremaster and the almost smooth body (few punctuations) makes it much more similar to the pupae of *D. friburgensis* (Orlandin et al. 2021).

Apatelodidae, like most moth groups, has been neglected in worldwide collections, although they belong to one of the most collectible moth groups in the world, the Bombycoidea (e. g. Saturniidae, Sphingidae, Brahmaeidae). As for these groups, detailed investigation on male genitalia showed that some old recognized wide-spread species comprise a set of cryptic species whose geographic distribution is much more restricted than previously thought. This is the case of South American specimens formerly identified as O. bifenestrata, that currently are the type series of O. jairana sp. nov., O. quilombola sp. nov., O. xeta sp. nov., and O. wayana sp. nov., whose differences in genitalia morphology are corroborated by great genetic distances in COX1 (barcode region, 658pb) sequences. The accumulated knowledge, especially in recent years, shows that there is still much to be studied to understand the evolutionary relationships between Apatelodidae lineages.

#### 5. Competing interests

The authors have declared that no competing interests exist.

#### 6. Acknowledgments

We are grateful to Dr Mirna Martins Casagrande, from Laboratório de Estudos de Lepidoptera Neotropical (UFPR), for making available the stereomicroscope used to illustrate the specimens. To Dr Sharon de Toledo Martins from the Center of Electronic Microscopy (CME) of the Universidade Federal do Paraná (UFPR) for the SEM images. To Dr Vitor Osmar Becker (VOB) and Dr Diego Dolibaina (DD) for loan specimens of Apatelodidae. To Dr Thamara Zacca (MNRJ) for dissecting specimens of Olceclostera. To Dr Pablo Mulieri, Dr Darío Lijtmaer, Dr Pablo Lavinia, and Dr Ezequiel Osvaldo Núñez Bustos (MACN) for providing data on Olceclostera from Argentina. To Dr Alessandro Giusti (NHMUK); Dr Scott Miller and Dr Talitta Simões (USNM); Dr Axel Hausmann and Dr Mei-Yu Chen (ZSM); Dr Théo Léger and Dr Viola Richter (MfN); Dr Jason J. Dombroskie and Jennifer Elizabeth Campos (CUIC) for making available photos of Olceclostera Types. Elton Orlandin (141475/2020-0) and Mônica Piovesan (150962/2022-3) thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for graduate and post-graduate fellowship, respectively.

### 7. References

- Balcázar-Lara MA, Wolfe KL (1997) Cladistics of the Ceratocampinae (Lepidoptera: Saturniidae). Tropical Lepidoptera Research 8(2): 1–53.
- Ballesteros-Mejia L, Arnal P, Hallwachs W, Haxaire J, Janzen D, Kitching I, Rougerie R (2020) A global food plant dataset for wild silkmoths and hawkmoths and its use in documenting polyphagy of their caterpillars (Lepidoptera: Bombycoidea: Saturniidae, Sphingidae). Biodiversity Data Journal 8: e60027. https://doi.org/10.3897/ BDJ.8.e60027
- Becker VO, Orlandin E (in press) A review of the Neotropical Apatelodidae genus *Prothysana* Walker (Lepidoptera: Bombycoidea). SHILAP Revista de lepidopterología.
- Biezanko CM (1986) Adelocephalidae, Saturniidae, Mimallonidae, Lasiocampidae, Eupterotidae e Lymantriidae da região missioneira do Rio Grande do Sul. Revista Centro Ciências Rurais 16(2): 89–112.
- Biezanko CM, Ruffnelli A, Link D (1974) Plantas y otras sustancias alimenticias de las orugas de los lepidopteros uruguayos. Revista Centro Ciências Rurais 4(2): 107–148.
- Borror DJ, deLong DM (1971). An introduction to the study of insects. Holt, Rinehart and Winston, New York, 812 pp.
- Butler AG (1879) On the Lepidoptera of the Amazons, collected by Dr James W. H. Trail, during the years 1873 to 1875. Transactions of the Royal Entomological Society of London 26(1): 39–84. https:// doi.org/10.1111/j.1365-2311.1878.tb01942.x
- CCDB (2013) The Canadian Centre for DNA Barcoding (CCDB). CCBD, Centre for Biodiversity Genomics, University of Guelph, Guelph, Ont. https://ccdb.ca/solutions
- Costa-Lima A (1950) Família Eupterotidae. In: Costa-Lima A (Ed.) Insetos do Brasil. Lepidópteros. 2ª parte. Imprensa Nacional, Brasil, 281–282.
- Diniz IR, Braga L, Lepesqueur C, Silva N, Morais H (2013) Lagartas do Cerrada: Guia de Campo. Technical Books, 200 pp.
- Draudt MWK (1929) Familia Bombycidae. Zanolinae. In: Seitz A. (Ed.) Die amerikanischen Spinner und Schwärmer. Die Groß Schmetterlinge der Erde. Verlag des Seitzschen Werkes: 693–711.

- Druce H (1890) Descriptions of new Species of Lepidoptera Heterocera from Central and South America. Proceedings of the Zoological Society of London: 493–520. https://doi.org/10.1111/j.1469-7998.1885. tb07859.x
- Dyar H (1906) Description of new North American moths and larvae. Bulletin of the Brooklyn Institute of Arts and Sciences 1: 185–197
- Edwards H (1886) Notes on North American Zygaenidae and Bombycidae with descriptions of new forms. Entomologica Americana 2(1): 8–15.
- Franclemont JG (1973) Mimallonoidea (Mimallonidae) and Bombycoidea (Apatelodidae, Bombycidae, Lasiocampidae). Classey, 86 pp.
- Freitas AVL, Brown KS (2004) Phylogeny of the Nymphalidae (Lepidoptera). Systematic Biology 53(3): 363–383. https://doi.org/10. 1080/10635150490445670
- Grote AR (1864) Descriptions of North American Lepidoptera. Proceedings of the Entomological Society of Philadelphia 4: 322–326.
- Hamilton CA, St Laurent RA, Dexter K, Kitching IJ, Breinholt JW, Zwick A, Timmermans MJTN, Barber JR, Kawahara AY (2019) Phylogenomics resolves major relationships and reveals significant diversification rate shifts in the evolution of silk moths and relatives. BMC Evolutionary Biology 19(1): 182. https://doi.org/10.1186/s1-2862-019-1505-1
- Hebert PDN, Cywinska A, Ball SL, de Waard JR (2003) Biological identifications through DNA barcodes. Proceedings of the Royal Society of London. Series B: Biological Sciences 270(1512): 313–321. https://doi.org/10.1098/rspb.2002.2218
- Herbin D (2015) Description de nouvelles espèces d'Hétérocères de Guyane Française avec notes taxinomiques. Antenor 2(1): 81–105.
- Herbin D (2017) Description de nouveaux Hétérocères néotropicaux (Lepidoptera Apatelodidae). Antenor 4(2): 114–134.
- Herbin D (2018) Nouveaux Apatelodidae du Muséum d'Histoire Naturelle de Genève (Lepidoptera Bombycoidea). Antenor 5(1): 37–42.
- Herbin D (2021a) New Bombycoidea species from French Guiana (Lepidoptera Apatelodidae and Bombycidae). Antenor 8 (2): 159–166.
- Herbin D (2021b) New species of Apatelodidae (Lepidoptera: Bombycoidea) from Andean countries. The European Entomologist 13(1): 55–64.
- Herbin D, Beccacece HM (2018) Description of three new species of Apatelodidae from the southern neotropical region (Lepidoptera, Bombycoidea). ZooKeys 788: 3–17. https://doi.org/10.3897/zookeys.788.25323
- Herbin D, Mielke CGC (2018) Preliminary list of Apatelodidae (Lepidoptera: Heterocera, Bombycoidea) from the Brazilian state of Maranhão (Ne Brazil) with description of ten new species. The European Entomologist 9(3): 105–140.
- Herbin D, Monzon Sierra J (2015) Description of four new species of Mimallonidae and four new species of Apatelodidae from the El Mirador Basin in northern Guatemala (Lepidoptera Mimallonidae and Apatelodidae). Antenor 2(2): 176–197.
- Herbin D, Orlandin E, Carneiro E (2021) Taxonomic revision of *Arotros* Schaus, 1892 with description of seven new species (Lepidoptera: Apatelodidae). Nachrichten des Entomologischen Vereins Apollo 42(3): 125–144.
- Janzen D, Hallwachs W (2009). Dynamic database for an inventory of the macrocaterpillar fauna, and its food plants and parasitoids, of Área de Conservación Guanacaste (ACG), northwestern Costa Rica. http://janzen.sas.upenn.edu
- Jones ED (1908) Descriptions of New Species of Lepidoptera-Heterocera from South-east Brazil. Transactions of the Entomological Society of London 1: 143–176.

- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS (2017) ModelFinder: Fast model selection for accurate phylogenetic estimates. Nature Methods 14(6): 587–589. https://doi.org/10.1038/ nmeth.4285
- Katoh K, Standley DM (2013) MAFFT Multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30(4): 772–780. https://doi.org/ 10.1093/molbev/mst010
- Kirby WF (1892) A synonymic catalogue of Lepidoptera Heterocera. (Moths). Vol. 1. Sphinges and Bombyces. Taylor and Francis, London, 951 pp.
- Kitching I, Rougerie R, Zwick A, Hamilton C, St Laurent R, Naumann S, Ballesteros-Mejia L, Kawahara A (2018) A global checklist of the Bombycoidea (Insecta: Lepidoptera). Biodiversity Data Journal 6: e22236. https://doi.org/10.3897/BDJ.6.e22236
- Kristensen NP (2003) 4. Skeleton and muscles: Adults. In: Kükenthal W (Ed) Band 4: Arthropoda 2 Hälfte: Insecta, Lepidoptera, Moths and Butterflies, Teilband/Part 36, Vol 2: Morphology, physiology, and development. de Gruyter, Berlin, Boston: 39–132.
- Lintner JA (1874) Descriptions of the larvae of some Bombycidae. Entomological Contributions 3: 129–134.
- Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Molecular Biology and Evolution 32(1): 268–274. https://doi.org/10.1093/molbev/msu300
- Orlandin E, Carneiro E (2021) A new species of *Pantelodes* (Lepidoptera: Bombycoidea: Apatelodidae) from Southern Bahia, Brazil. Zootaxa 5047(5): 589–595. https://doi.org/10.11646/zootaxa.5047. 5.9
- Orlandin E, Piovesan M, Carneiro E (2021) Biological and morphological aspects of *Drepatelodes* Draudt 1929, the genus sister to all other Apatelodidae (Lepidoptera: Bombycoidea). Studies on Neotropical Fauna and Environment: 1–14. https://doi.org/10.1080/01650521.2 021.2003585
- Orlandin E, Piovesan M, Carneiro E (2022a) From molecular data to natural history: A new species of *Apatelodes* (Lepidoptera: Bombycoidea: Apatelodidae) from southern Brazil. Canadian Journal of Zoology 100(3): 161–175. https://doi.org/10.1139/cjz-2021-0160
- Orlandin E, Piovesan M, Herbin D, Carneiro E (2022b) Immature stages, description of female, and redescription of male of *Apatelodes kotzschi* Draudt 1929 (Lepidoptera: Bombycoidea: Apatelodidae). Neotropical Entomology 52(1): 92–103 https://doi.org/10.1007/ s13744-022-01003-3
- Packard SA (1864) Synopsis of the Bombycidae of the United States. Proceedings of the Entomological Society of Philadelphia 3: 331– 396.
- Packard SA (1895) Monograph of Bombycine moths of America North of Mexico including their transformations and origem of larval markings and armature. Parte I. Family 1. Notodontidae. Memoirs of the National Academy of Sciences 7: 291.
- Peigler RS (1994) Life history of *Olceclostera seraphica* (Lepidoptera: Bombycidae, Apatelodinae). Nachrichten des Entomologischen Vereins Apollo 15: 245–252.
- Petersen W (1904) Die Morphologie der Generationsorgane der Schmetterlinge und ihre Bedeutung f
  ür die Artbildung. M
  émoires de L'Acad
  émie Imp
  ériale d
  ês Sciences de St. P
  étersbourg 8: 1–84.
- Peterson A (1962) Larvae of Insects. I (Lepidoptera and Hymenoptera). Ohio State University, Columbus, Ohio; Edwards Brothers, Ann Arbor, Michigan: 183 pp.

- Ratnasingham S, Hebert PDN (2007) Bold: The Barcode of Life Data System. Molecular Ecology Notes 7(3): 355–364. https://doi.org/ 10.1111/j.1471-8286.2007.01678.x
- Robinson GS, Ackery PR, Kitching IJ, Beccaloni GW, Hernández LM (2010) HOSTS – A database of the world's Lepidopteran Hostplants. http://www.nhm.ac.uk/hosts
- Schaus W (1892) Descriptions of new species of Lepidoptera Heterocera from Brazil, Mexico and Peru-Part II. Proceedings of the Zoological Society of London: 318–340.
- Schaus W (1894) On new species of Heterocera from Tropical America. Proceedings of the Zoological Society of London 1: 225–243.
- Schaus W (1900) New species of Heterocera from tropical America. Journal of the New York Entomological Society 8(4): 225–234.
- Schaus W (1905) Descriptions of new South American moths. Proceedings of the United States National Museum 39: 179–345.
- Schaus W (1910) New species of Heterocera from Costa Rica—III. Annals and Magazine of Natural History 6(36): 561–585.
- Schaus W (1912) New Species of Heterocera from Costa Rica—XII. Annals and Magazine of Natural History 9(49): 34–56.
- Schaus W (1920) New species of Lepidoptera in the United States National Museum. Proceedings of the United States National Museum 57: 107–152.
- Schaus W (1927) New species of Heterocera (Lepidoptera) from Central and South America. Proceedings of the Entomological Society of Washington 29(5): 101–111.
- Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nature Methods 9: 671–675.
- Scoble MJ (1992) The Lepidoptera: Form, function, and diversity. Oxford University Press: 404 pp.
- Specht A, Corseuil E, Abelha HB (2008) Lepidópteros de importância médica: Principais espécies no Rio Grande do Sul. Editora Useb: 220pp.
- Snodgrass RE (1935). Principles of insect morphology. McGraw Hill Book Co: 667 pp.
- Stehr FW (1987) Order Lepidoptera, introduction part. In: Stehr FW (Ed) Immature insects. Kendall/Hunt: 208–305.
- Tamura K, Nei M (1993) Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. Molecular Biology and Evolution 10(3): 512–526. https://doi.org/10.1093/oxfordjournals.molbev.a040023
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. Molecular Biology and Evolution 38(7): 3022–3027. https://doi.org/10.1093/molbev/msab120
- Thomazini MJ (2010) Coleta e criação da lagarta da erva-mate, para estudos em laboratório. Colombo: Embrapa Florestas.
- Wagner DL (2005) Caterpillars of Eastern North America: A Guide to identification and natural history. Princeton University Press, 513 pp.
- Walker F (1855) List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. Lepidoptera Heterocera. Part IV. Edward Newman, London.
- Walker F (1865) List of the Specimens of Lepidopterous Insects in the Collection of the British Museum. Part XXXI. Supplement. Edward Newman, London.
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang GT (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. Molecular Ecology Resources 20(1): 348–355. https://doi.org/10.1111/1755-0998.13096
- Zwick A (2009) The principal structure of male genital sclerites and muscles of bombycoid moths, with special reference to Anthelidae

(Lepidoptera: Bombycoidea). Arthropod Structure & Development 38(2): 147–161. https://doi.org/10.1016/j.asd.2008.07.006

Zwick A, Regier JC, Mitter C, Cummings MP (2011) Increased gene sampling yields robust support for higher-level clades within Bom-

### **Supplementary Material 1**

#### Table S1

Authors: Orlandin E, Piovesan M, Herbin D, Carneiro E (2023) Data type: .xlsx

**Explanation note:** GenBank and BOLD Systems accession numbers of Olceclostera specimens used in the Maximum likelihood phylogenetic tree, based on cytochrome c oxidase subunit 1 (COX1) (DNA barcode).

**Copyright notice:** This dataset is made available under the Open Database License (http://opendatacommons.org/ licenses/odbl/1.0). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: https://doi.org/asp.81.e107507.suppl1

bycoidea (Lepidoptera). Systematic Entomology 36(1): 31–43. https://doi.org/10.1111/j.1365-3113.2010.00543.x

# **ZOBODAT - www.zobodat.at**

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Arthropod Systematics and Phylogeny

Jahr/Year: 2023

Band/Volume: 81

Autor(en)/Author(s): Orlandin Elton, Piovesan Monica, Herbin Daniel, Carneiro Eduardo

Artikel/Article: More than Olceclostera bifenestrata: New species and morphology of immature stages of Olceclostera Butler, 1879 (Lepidoptera: Bombycoidea, Apatelodidae) 1063-1088