

Distribution, conservation and prognosis for *Gampsocleis glabra* (HERBST, 1786) (Insecta: Ensifera) in Slovakia and the Czech Republic

Peter J. Fedor, Jaroslav Holuša, Oto Majzlan and Pavol Prokop

Abstract

The Heath Bush-cricket (*Gampsocleis glabra*) can be considered as a really xerothermophilous species preferring dry and warm steppe-like formations, quite intolerant to different or changed ecological conditions. Although this Euro-Asian bush-cricket occurs in many European countries, the distribution in Slovakia can be still classified as "mystic and unclear". However several records have been published on its presence from Slovakia (E and SW) and the Czech Republic (N). The references hinted at the fact that *G. glabra* seemed to appear just in Eastern Slovakia on its South close to the boundary with Hungary. However in the 90's the species was collected in South-Western Slovakia, especially in the surroundings of the Danube River. Due to a lack of sufficient information on the occurrence of *G. glabra* in Slovakia as well as in the Czech Republic it seems to be difficult to define the exact ecosozological status of the species and to determine relevant activities for its conservation. Since the main threat seems to be connected with the destruction of real and potential localities, conservation efforts should be proposed from this point of view.

Zusammenfassung

Die Heideschrecke (*Gampsocleis glabra*) ist eine ausgesprochen xerothermophile Art, die trockenwarme, steppenähnliche Lebensräume bewohnt und sehr empfindlich auf Änderungen der Habitatbedingungen reagiert. Obwohl diese eurasische Laubheuschrecke in einigen Ländern Europas vorkommt, ist ihre Verbreitung in der Slowakischen und Tschechischen Republik noch weitgehend unklar. Es liegen einzelne publizierte Nachweise der Art aus beiden Ländern vor. Ursprünglich wurde angenommen, *G. glabra* käme in der Slowakei nur im Osten nahe der ungarischen Grenze vor. In den 1990er-Jahren gelangen jedoch Aufsammlungen der Art im Südwesten des Landes in den Donauniederungen. Die nach wie vor unklare Verbreitungssituation in beiden Ländern erschwert eine genaue Bestimmung des Gefährdungsstatus und die Formulierung geeigneter Schutz- und Fördermaßnahmen. Da die Gefährdungsursachen für die Art zu einem erheblichen Teil in der Zerstörung bestehender und potenziell geeigneter Lebensräume liegen dürften, sollten sich Schutzbemühungen für *G. glabra* v.a. auf diese Problematik konzentrieren.

Introduction

The Heath Bush-cricket (*Gampsocleis glabra*) is a xerothermophilous species preferring dry and warm steppe-like formations, quite intolerant to different or changed ecological conditions. ČEJCHAN (1959) has even classified it as a deserticolous species. It usually inhabits various meadows with sparse bush or tree stands. GULIČKA (1992) underlined sites with poor grass vegetation as well as dry soils. BELLMAN (1985) considered as suitable habitats those steppe regions with high grass (*Stippa* spp.). However, it obviously depends on the availability of habitats, geographical, and ecological aspects. HARZ (1957) and KNIPPER (1958), for example, emphasized even heather (*Callunetum*) as the only alternative of habitat for the Heath bush-cricket in the Lüneburger Heide (Germany).

It feeds on various foods – plant as well as animal organic matter. As a heliophilous species, it is active especially in sunshine; the stridulation is similar to that of *Tettigonia viridissima* or *Metrioptera roeselii* (BELLMAN 1985). More detailed information on biology and ecology of *Gampsocleis glabra* can be seen in HARZ (1957).

Distribution in Slovakia and the Czech Republic

This Euro-Asian Bush-Cricket occurs in many European countries. KALTENBACH (1970) presents its distribution from Western Siberia and Kazakhstan to Spain, including Eastern and Southern Europe. However, it is absent from the British Isles, Portugal and Italy and occurs only in isolated spots in Spain, Germany and Poland (LIANA 1992, MAAS et al. 2002).

The distribution of *Gampsocleis glabra* in Slovakia can be still considered as unclear. GULIČKA (1992) hints at its "sporadic occurrence" in former Czechoslovakia. However several records have been published on its presence either in Slovakia or and rather than in Moravia and the Czech Republic (Figure 1, 2). In the 50's MAŘAN (1954) observed the species in Eastern Slovakia (Svätá Mária) while he was dealing with xerothermophilous orthopterans historically immigrating from the pannonian lowlands of Hungary and Austria. And finally at the end of 50's ČEJCHAN (1959) updated the previous knowledge by the record from Veľký Kamenec (mentioned as well as Veľký Kevežd) (Eastern Slovakia). This information came from Dr. A. Hoffer, who had found 22 males and 14 females here in July 11, 1953. Therefore ČEJCHAN (1959) considered this locality as the one with the most abundant population of *Gampsocleis glabra* in Slovakia. In 1985 he summarized all the available data on the occurrence of the species and added the new record from Somotor (Eastern Slovakia), when Dr. Hoffer had collected one female in 1951 (ČEJCHAN 1985). GULIČKA (1992) dealt with the conservation of this bush-cricket and as for its occurrence he mentioned "for example" Bodrog and Veľký Kamenec (Eastern Slovakia) with no more detailed data or reference.

All the available references hinted at the fact that *Gampsocleis glabra* seemed to appear just in Eastern Slovakia on its South close to the boundary with Hungary. ČEJCHAN (1985) wrote that it has not been recorded out of Eastern Slovakia yet. However in the 90's the species was collected in South-Western Slovakia, espe-

cially in the surroundings of the Danube river. This region, although formed mainly by floodplain forests, sporadically provides very dry and warm ecological conditions suitable for xerothermophilous species such as the Heath Bush-cricket. In July 17, 1991 Majzlan found one specimen in Dobrohošť (MAJZLAN, 1992) and in July 26, 1998 Fedor captured one female in Gabčíkovo – Istragov – in meadow communities of Dauco-Melilotion on gravel terraces (FEDOR 2000, 2001). Another specimen of *Gampsocleis glabra* from South-Western-Slovakia was recorded in Abov-Hurbanovo, a sandy xerothermous site, in August 1999 (MAJZLAN et al. 2000a, b).

The first record of *Gampsocleis glabra* from the Czech Republic was published by SEIDEL (1836), from Sedlo hill near Litoměřice. Although, this record was marked by GULIČKA (1992) as disputable, from the zoogeographical point of view the occurrence is possible. The area of northwestern Bohemia is very dry and warm with the occurrence of several xerothermic species on the northern edge of their ranges (MAŘAN 1964, ČEJCHAN 1981, 1982, HONCÚ 1993, HOLUŠA & HOLUŠA 2002). Recently, the existence of population in this locality of Sedlo is impossible, due to reforestation of hill.

The only record of the Heath Bush-cricket from Moravia comes from the locality of "Větrníky" at the town of Vyškov mentioned by GINTER (1951) (recently a national natural reserve Větrníky, 15.09.1943, 6M, ??F, 02.08.1946, 2M, revid. J.Holuša, coll. MJM). The local occurrence of the species in Southern Moravia has been recorded by GULIČKA (1992) as well. Recently, we do not know any locality with its population.

The record by HYKEŠ (1925) at the town of Osoblaha in Silesia (former historical country) is very questionable (ČEJCHAN 1959). It is not clear whether it is an original or taken record. In spite of other improbable species (*Anechura bipunctata*) noted from Silesia, the record seems to be unreliable. ZACHER (1907) cited only localities from Upper Silesia.

The following survey presents a list of localities of *Gampsocleis glabra* in Slovakia and the Czech Republic with a grid reference number and relevant literature references.

Slovakia

Svätá Mária (7596): MAŘAN (1954).

Veľký Kamenec (Veľký Kevežd or Kevežd) (7696): ČEJCHAN (1959) GULIČKA (1992).

Somotor (7596): ČEJCHAN (1985).

Dobrohošť (8070): MAJZLAN (1992).

Gabčíkovo – Istragov (8171): FEDOR (2000, 2001).

Abov – Hurbanovo (8175): MAJZLAN et al. (2000 a, b).

The Czech Republic

Sedlo hill (5451): SEIDEL (1836).

Větrníky u Vyškova (6768): GINTER (1951).




Fig. 1: Distribution of *Gampsocleis glabra* in Slovakia.




Fig. 2: Occurrence of *Gampsocleis glabra* in the Czech Republic.

Conservation and prognosis for future

Due to a lack of sufficient information on the occurrence of *Gampsocleis glabra* in Slovakia as well as in the Czech Republic it seems to be difficult to define the exact ecosozological status of the species, and to determine the relevant activities for its conservation absolutely reliably. The species has been listed in the Red Data Book of Slovakia and the Czech republic as an endangered insect (GULIČKA 1992). Generally, the Heath Bush-cricket is usually included in national Red Data Books in Central Europe. Even in Hungary, where the population is much more abundant than in Slovakia, the species has been under the nature conservancy (SZÖVÉNYI & NAGY 1999). In Niederösterreich (Austria) it has been considered as an endangered species classified into the category 1 ("vom Aussterben bedroht"; BERG & ZUNA-KRATKY 1997). The species has been included into the Polish red list as well (LIANA 1992).

Gampsocleis glabra occurs on the northern edge of its range in Slovakia and the Czech republic. The known data are concentrated in the southern parts of the countries mentioned. Therefore, the occurrence is very local and may be ephemeral. But even in much more warm areas the fact is that the Heath Bush-cricket finds its suitable ecological conditions sporadically, on remarkably dry and warm sites. Hence the threat must be intensively interacted with the destruction of real and potential localities. This is actually the problem of many other xerothermophilous orthopterans, e.g. *Saga pedo* or *Acrida ungarica*. GULIČKA (1992) presented as main reasons of the threatening the radical impact on original steppe and wooden-steppe communities, especially their change into farmland with intensive management (application of chemicals, deep tilling etc.). Today, *Gampsocleis glabra* escaped to refugia, that have not been under significant intensive human land use, especially due to its low value for agriculture (sand dunes, gravel terraces, dams etc.).

The conservation of the species is only possible in protection in situ. Nature reserves of various character and status may save or keep the populations locally. A need for conservation of steppe habitats as potential biotops for *Gampsocleis glabra* has been presented by BERG & ZUNA-KRATKY (1997). This seems to be a general plea for nature conservation in endangered habitats of xerothermous formations. Southern Slovakia is the most northern point of the formerly large pannonian steppe with its center in Hungary and Austria. Thus, the study of its species diversity is necessary for effective conservation activities. The same consequences appear from the paper by BERG & BIERINGER (1998) from Niederösterreich (Austria).

On the other hand an interesting question appears in effect of aridization that significantly leads to increase in abundance of some xerothermophilous species and their infiltration into suitable biotopes. This phenomenon may cause a significant increase in population density of *Gampsocleis glabra* in future, actually as it has been observed at several other species such as *Oecanthus pellucens* (GEREND & PROESS 1994, FEDOR & MAJZLAN 2001).

Acknowledgement

This project has been supported by VEGA: 1/9109/02.

Authors:

Dr. Peter J. Fedor
Comenius University, Faculty of Natural Sciences
Department of Ecosozology
Mlynská dolina
Bratislava, SK - 842 15, Slovakia
E-Mail: fedor@fns.uniba.sk

Dr. Jaroslav Holuša
Forestry and Game Management Research Institute Jíloviště-Strnady
Office Frýdek-Místek
Nádražní 2811
73801 Frýdek-Místek, Czech Republic
E-Mail: holusaj@seznam.cz

Prof. Dr. Oto Majzlan
Comenius University, Faculty of Education
Račianska 59
Bratislava, SK - 813 34, Slovakia
oto.majzlan@fedu.uniba.sk

Dr. Pavol Prokop
Institute of Zoology, Slovak Academy of Sciences
Dúbravská 9
Bratislava, SK - 842 06, Slovakia
pprokop@post.sk

References

- BELLMANN, H. (1985): Heuschrecken beobachten – bestimmen. – Neumann – Neudamm, Mel-sungen, 210 S.
- BERG, H.M. & BIERINGER, G. (1998): Bemerkenswerte Neu- und Wiederfunde zur Heuschreckenfauna Niederösterreichs (Österreich). – *Articulata* 13 (2): 163–172.
- BERG, H.M. & ZUNA-KRATKY, T. (1997): Heuschrecken und Fangschrecken. Eine Rote Liste der Niederösterreich gefährdeten Arten. – Herausgegeben von der NÖ Landesregierung/Abteilung Naturschutz, Wien, 112 S.
- ČEJCHAN, A. (1959): Príspevek k rozšírení nektých vzácných druhu orthopteroidního hmyzu v Čechách a na Slovensku. – *Acta mus. Reginaehrad., S.A. Sc. Nat.* 2: 173–182.
- ČEJCHAN A. (1981): K poznání orthopteroidního hmyzu (s.l.) ČSSR II. On the orthopteroid insects (s.l.) of Czechoslovakia II. – *Čas. Nár. Muz. Praha, Řada přírodověd.* 150: 147–151 (in Czech, English abstract).
- ČEJCHAN A. (1982): K poznání orthopteroidního hmyzu (s.l.) ČSSR III. On the orthopteroid insects (s.l.) of Czechoslovakia III. – *Čas. Nár. Muz. Praha, Řada přírodověd.* 151: 1–13 (in Czech, English abstract).
- ČEJCHAN, A. (1985): K poznání orthopteroidního hmyzu (s. l.) ČSSR IV. (Dermaptera, Grylloptera, Orthoptera s. str.). – *Čas. Nár. Muz., řada přírodov.* 154: 133–144.
- FEDOR, P.J. (2000): K biodiverzite rovnokřídlovců (Ensifera a Caelifera) v okolí geo-technického systému Dunajského vodného diela. – *Acta environmentalistica* 10: 171–176.
- FEDOR, P.J. (2001): Bioindikačný význam rovnokrídleho hmyzu (Ensifera et Caelifera) vo vzťahu k zmenenej krajine v okolí Vodného diela Gabčíkovo. – *Folia Faunistica Slovaca* 6, supplement 1, 91 S.
- FEDOR, P.J. & MAJZLAN, O. (2001): Distribution and Infiltration of the tree cricket *Oecanthus pellucens* (SCOPOLI, 1763) to Unoriginal Conditions in Slovakia. – *Bull. Soc. Nat. Luxemb.* 102: 103–108.
- GEREND, R. & PROESS, R. (1994): Aktueller Nachweis des Weinhähnchens *Oecanthus pellucens* (SCOPOLI, 1763) im Süden Luxemburgs (Saltatoria, Gryllidae). – *Bull. Soc. Nat. Luxemb.* 95: 245–246.
- GINTER, O. (1951): Výskyt vzácnějších Orthopter na Moravě. – *Fol. Ent., XIV:* 167–169.
- GULIČKA, J. (1992): Rovnokřídlovce (Caelifera, Ensifera). – In: ŠKAPEC, L. (ed.): Červená kniha ohrozených a vzácných rostlin a živočichův ČSFR 3 Bezstavovce: 70–73, Bratislava, Příroda, 152 S.
- HARZ, K. (1957): Die Geradflügler Mitteleuropas. – Jena, VEB G. Fischer Verlag, 949 S.
- HOLUŠA, J. & HOLUŠA, O. (2002): Occurrence of grasshopper *Stenobothrus eurasius bohemicus* (Caelifera: Acrididae) in the Czech Republic. – *Articulata* 17: 89–93.
- HONCÚ, M. (1993): Príspevek k poznání orthopteroidního hmyzu severozápadních Čech. – Beitrag zur Kenntnis der Orthopteren-Fauna aus Nordwestböhmen. – *Sbor. Okres. Muz. Most, Řada přírodověd.* 13–14 (1992): 89–108 (in Czech, German abstract).
- HYKEŠ, O.V. (1925): Zvířena našeho Slezska. [Fauna of Czech Silesia]. – In: LHOTSKÝ, A. (ed.): Vlastivědný Sborník Slezský, Část I. – Přírodní popis a hospodářské poměry. Školství, Statistika. Slezská Grafia, Opava: 51–66 (in Czech).
- IHSSEN, G. (1957): Die Verbreitung von *Gampsocleis glabra* (HERBST 1786) in der Lüneburger Heide (Orthopt., Ensifera, Tettigoniidae). – *Mitt. dtsh. ent. Ges.* 16 (1): 8–9.
- KALTENBACH, A. (1970): Zusammensetzung und Herkunft der Orthopterenfauna im pannonischen Raum Österreichs. – *Ann. Naturhist. Mus. Wien* 74: 159–186.
- KNIPPER, H. (1958): Weitere Beiträge zum Vorkommen von *Gampsocleis glabra* (HERBST, 1786) in der Lüneburger Heide (Orthopt., Ensif., Tettigoniidae). – *Verh. Ver. naturw. Heimatsforsch. Hamburg* 33: 16–24.
- LIANA, A. (1992): Owady Prostoskrzydłe Orthoptera. – In: GLOWACINSKI, Z. (ed.): Czerwona Lista Zwierząt Ginacych i Zagrozonych w Polsce. – Polska Akademia Nauk, Krakow: 85–91.
- MAAS, S., DETZEL, P., & STAUDT, A. (2002): Gefährdungsanalyse der Heuschrecken Deutschlands – Verbreitungsatlas, Gefährdungseinstufung und Schutzkonzepte. – Schriftenreihe Bundesamt für Naturschutz, 401 S.
- MAJZLAN, O. (1992): Význam niektorých skupín hmyzu pre monitoring Podunajska. – *Spr. Slov. ent. Spoloč. SAV*, 4 (3): 1–9.
- MAJZLAN, O., RYCHLÍK, I. & FEDOR, P. (2000a): Stav biodiverzity psamofilů (Ectognatha) na vltých pieskoch lokality Aba pri Hurbanove. – *Acta environmentalistica* 10: 215–220.

- MAJZLAN, O., RYCHLÍK, I., & FEDOR, P.J. (2000b): Chrobáky (Coleoptera), kobylky (Ensifera), koníky (Caelifera) na viatych pieskoch lokality Aba pri Hurbanove, Liščie diery pri Nesvadoch a Balvany pri Kameničnej (južné Slovensko). – *Rosalia*, Nitra, 15: 155–174.
- MAŘAN, J. (1954): Rovnokřídly hmyz státních přírodních rezervací v okolí Štúrova na jižním Slovensku. – *Ochrana přírody* 9: 132–139.
- MAŘAN, J. (1964): Výsledky orthopterologického výzkumu v Lounském středohoří. – Einige Ergebnisse der orthopterologischen Forschungen im Launer Mittelgebirge. – In: Referáty entomologického sympozia (22. –24. září 1964) u příležitosti oslav 150let trvání Slezského muzea v Opavě. Slezské muzeum, Opava: 165–178 (in Czech, German abstract)
- SEIDEL, W. (1836): Die Orthopteren Böhmens. – *Weitenwebers Beiträge*: 205–223.
- SZÖVÉNYI, G., & NAGY, B. (1999): Szikes és löszpuszta élőhelyek egyenesszárnyú rovar (Orthoptera) együtteseinek összehasonlító elemzése a Körös-Maros Nemzeti Park területén. – *Crisicum* 2: 115–122.
- ZACHER, F. (1907): Beitrag zur Kenntnis der Orthopteren Schlesiens. – *Ztsch. Wiss. Insektenbiol.* 3 (6): 179–185; 3 (7): 211–217.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Articulata - Zeitschrift der Deutschen Gesellschaft für Orthopterologie e.V. DGfO](#)

Jahr/Year: 2004

Band/Volume: [19_2004](#)

Autor(en)/Author(s): Fedor Peter J., Holusa Jaroslav, Majzlan Oto, Prokop Pavol

Artikel/Article: [Distribution, conservation and prognosis for *Gampsocleis glabra* \(HERBST, 1786\) \(Insecta: Ensifera\) in Slovakia and the Czech Republic 217-224](#)