

Recent expansions of the bush-crickets *Phaneroptera falcata* and *Phaneroptera nana* (Orthoptera: Tettigoniidae) in the Czech Republic

Petr Kočárek, Jaroslav Holuša, Robert Vlk, Pavel Marhoul & Thomas Zuna-Kratky

Abstract

Phaneroptera falcata was known to be widespread only in southern and central Moravia till the year 2000. During the last years, quick expansion to northern Moravia and Silesia (Moravian-Silesian region) has been observed. Based on the sequence of findings, consecutive spreading through Bohemia is also supposed. *Phaneroptera nana* was published from southern Moravia firstly in 2002 with findings from 1992 and 2001. At the present time, occurrence in nine localities is confirmed with the northernmost finding in Brno. Climatic changes associated with the current global warming trend in the last decades together with the increase of abandoned fields and meadows are probably the main reasons of the expansions.

Zusammenfassung

Phaneroptera falcata war bis in das Jahr 2000 in Tschechien lediglich aus Süd- und Zentralmähren als weit verbreitet bekannt. In den letzten Jahren wurde jedoch eine rasche Ausbreitung nach Nordmähren und Schlesien beobachtet. Die zeitliche Aufeinanderfolge der Nachweise macht es wahrscheinlich, dass die Einwanderung durch Böhmen erfolgte. *Phaneroptera nana* wurde in Tschechien erstmals in den Jahren 1992 und 2001 in Südmähren gefunden. Bis 2007 gelangen Nachweise an neun verschiedenen Stellen, die nördlichste davon bei Brno. Klimaveränderungen in Zusammenhang mit der derzeit beobachteten Erderwärmung sowie die Zunahme brachgefallener landwirtschaftlicher Flächen dürften die Hauptgründe für diese Ausbreitungen darstellen.

Introduction

Species of the genus *Phaneroptera* are long-winged thermophilous bush-crickets which are able to quickly colonize new sites (NAGY 1992). Two species of this genus, *P. falcata* (Poda, 1761) and *P. nana* Fieber, 1853, occur in the Czech Republic. Both species occupy a wide variety of open habitats, from xerothermic vegetation (esp. forest steppes) to wetlands (esp. lowland meadows and floodplain forest edges) including ruderal habitats in towns; but they demand taller herbaceous or shrubby vegetation (or trees) everywhere (KÖHLER 2001).

P. falcata is an Euro-Siberian species, distributed from France and Belgium to Japan (KOČÁREK et al. 2005). In the Czech Republic, this species is historically widespread within southern and central Moravia, which is documented by many

published records (survey see CHLÁDEK 1993). Only one single specimen from Horní Radíkov (6857) has been published from Bohemia (ČEJCHAN 1981). From the northern part of Moravia and Silesia (Moravian-Silesian region) the only occasional occurrence of a single specimen has been found in Háj ve Slezsku (6074) (DOBŠÍK & CHLÁDEK 1975). Both records have not been verified during the last 30 years.

The expansion of *P. falcata* to the north has been noticed in several countries of Western Europe (e.g. Germany, Belgium and Russia; see KLEUKERS et al. 1996, DECLER et al. 2000, LANDECK et al. 2005, STOLZENBURG 2005, BOLSHAKOV 2006).

P. nana is a circummediterranean species, distributed in Europe, Africa and Asia Minor (KOČÁREK et al. 2005). From the Czech Republic, only records of two single specimens caught near Strachotín (7065) in 1992 and Pouzdřany (7065) in 2001 in the most southern part of Moravia were published by VLK (2002). In the neighboring south-western part of Slovakia, the population of this species has been found for the first time in 1994 by HOLUŠA & VIDLIČKA (1997). At the present time it is known from many localities in southern Slovakia (e.g. MAJZLAN & FEDOR 1997, MAJZLAN et al. 2000, FEDOR 2000, 2001, GAVLAS 2001, CHLÁDEK & GAVLAS 2004 and own unpublished data). The expansion of this species has been observed also in northern Austria (BIERINGER & ROTTER 2001, BERG et al. 2005, ZECHNER & KOSCHUH 2005, archive of Orthopterenkartierung Ostösterreich), France, Switzerland and Germany (CORAY 2003, SARDET et al. 2005, BAUR et al. 2006).

The data about spreading of both *Phaneroptera* species in the Czech Republic are analyzed in this contribution.

Data base and methods

Data of *P. falcata* and *P. nana* presented in this contribution are based mainly on author's collecting activities carried out during the period 1995-2007. Unpublished data preserved in the collection of O. Ginther in the Museum of Zlín, Czech Republic have also been processed. The majority of *P. falcata* faunistic data were published in a previous conference contribution (KOČÁREK & HOLUŠA 2006). Most of the mentioned material is kept in author's collections. Original data are arranged as follow:

Čejkovice¹ – Špidlák Nature Reserve² (7167)³, 48°55'1"N, 16°57'33"E⁴, 210⁵, 6.ix.2006⁶, 0/1 and 1n⁷, PK⁸;

¹ = village

² = specification of locality

³ = the grid mapping square code used for faunistic research in the Czech Republic (PRUNER & MÍKA 1996)

⁴ = GPS coordinates

⁵ = altitude in m a.s.l.

⁶ = date of collection (finding)

⁷ = number of males, females and nymphs collected

⁸ = leg., det. et coll.

Abbreviations: PK – Petr Kočárek, Ostrava; JH – Jaroslav Holuša, Frýdek-Místek; RV – Robert Vlček, Brno; PM – Pavel Marhoul, Praha; TZK – Thomas Zuna-Kratky, Wien (A); JD – Jan Dolanský, Pardubice; OK – Ondřej Konvička, Veselí nad Moravou; BM – Bohuslav Mocek, Hradec Králové (Mocek leg. et det., Regional Museum Hradec Králové coll.).

Results and discussion

Published faunistic data

P. falcata: individual faunistic records published till 1990 were summarized by CHLÁDEK (1993), all the later by KOČÁREK & HOLUŠA (2006).

P. nana: VLK (2002), CHLÁDEK (2006).

Unpublished faunistic data

P. falcata:

Horní Lipová – Smrk Mt. (**5768**), 50°13'52.5"N, 17°2'3"E, 1060, 29.viii.2007, 0/1, JH; Krnov – Staré hliniště (**5872**), 50°6'29"N, 17°41'38"E, 325, 14.ix.2007, 0/1 observ., PK & JH.; Kunětická hora – Kunětická horahill (**5960**), 50°4'47.5"N, 15°48'42.5"E, 300, 30.viii.2007, cca 10 ex. observ., 0/1 and 50°4'51"N, 15°48'34.5"E, 240, 30.viii.2007, 0/1, BM; Lázně Bohdaneč – Na Kovárně (**6059**), 50°5'24"N, 15°42'26.5"E, 220, 6.viii.2007, 0/1 ex. documented by photo, JD; Skačice – Profil Morávky Nature Monument (**6376**), 49°39'58.5"N, 18°23'41"E, 20.vi.2007, 370, 6 n., PK; Halenkov – Lušová (**6674**), 49°21'27.5"N, 18°9'5.5"E, 525, 18.viii.2006, 1/0, PM; Maletice (**6751**), 49°13'45"N, 14°11'22"E, 375, 18.vii.2007, 1 ex. observ., PM; Vladislav – syenite slopes (**6761**), 49°12'58.5"N, 15°59'10"E, 420, 2.viii.2007, 3/0, RV; Zahrádka – Častotice (**6762**), 49°14'24"N, 16°5'9"E, 470, 2.ix.2007, 0/1 documented by photo – L. Jeřábková, PM det.; Zdounky (**6770**), 49°13'0"N, 17°19'13"E, 335, 19.viii.2007, 1/1, JH; Vlachovice – Za Hájem (**6873**), 49°7'28.5"N, 17°54'53"E, 415, 30.ix.2007, 0/1, OK; Nedašov – Kaňoury Nature Monument, (**6874**), 49°6'51.5"N, 18°6'24"E, 625, 15.viii.2007, 1 ex., OK; Nedašova Lhota (**6874**), 49°7'18"N, 18°4'48.5"E and 49°7'24.5"N, 18°4'54.5"E, 440, 19.viii.2007, 1 and 3 ex. resp., OK; Jaronice – hill Vráže (**7051**), 48°58'51"N, 14°19'44"E, 405, 30.vii.2006, documented by photo – P. Kouba, PM det.; Starý Hrozenkov (**7073**), 48°58'22"N, 17°52'11.5"E, 490, 8.vii.2007, cca 20 ex. observ., PK; Javorník nad Veličkou – Kománkův Mlýn (**7171**), 48°51'53"N, 17°32'38.5"E, 355, 26.vii.2007, cca 10 ex. observ., OK; Suchov – Suchovské mlýny, (**7171**), 48°53'7"N, 17°34'40.5"E, 385, 26.vii.2007, cca 20 ex. observ., OK; Velká nad Veličkou – Zahrady pod Hájem Nature Reserve (**7171**), 48°53'1"N, 17°31'46"E, 344, 14.ix.2007, 1 ex., OK; Lanžhot – Ranšpurk Nature Reserve (**7367**), 48°40'8"N, 16°56'24.5"E, 150, 5.ix.2006, cca 10 ex. observ., JH.

P. nana:

Brno – Špilberk (**6865**), 49°11'39"N, 16°35'48.5"E, 230, 20.x.2006, 1/0, RV; Pouzdřany (**7065**), 48°56'39.5"N, 16°38'39"E, 250, 16.ix.2006, 3/0, JH; Drnholec (**7164**), 48°51'41"N, 16°29'44.5"E, 170, 19.x.2006, 1/0 and 48°51'37"N, 16°29'5.5"E, 175, 21.x.2006, 0/1, RV; Hrušovany nad Jevišovkou – centre of the

village (**7164**), 48°49'47.5"N, 16°24'3.5"E, 185, 12.x.2007, 0/1, RV; Mikulov (**7165**), 48°48'33.5"N, 16°38'36.5"E, 295, 18.ix.2006, 3/0, PK; Čejkovice – Špidlák Nature Reserve (**7167**), 48°55'1"N, 16°57'33"E, 210, 6.ix.2006, 0/1 and 1n, PK; Břeclav (**7267**), 48°44'N, 16°54'E, 160, 31.viii.2006, several males calling in gardens and bushes, TZK; Lanžhot (**7267**), 48°43'N, 16°56'E and 48°44'N, 16°55'E, 150, 31.viii.2006, several males calling in fallow land and edges of street, TZK.

Recent expansion of *P. falcata* and *P. nana* in the Czech Republic

Phaneroptera falcata has been widespread historically within southern and central Moravia (CHLÁDEK 1993). In 2000, the first breeding population of this species in northern Moravia was found in Kunín (6373) (KOČÁREK & NĚMEČKOVÁ 2000). Next populations were found during the two successive years in other sites of Poodří Protected Landscape Area – Kunín (6373), Bartošovice-Hukovice (6374), Bartošovice na Moravě (6374) and consecutively many populations were found in the surroundings of the city of Ostrava, and towns of Opava, Karviná and Frýdek-Místek during 2004-2005 (KOČÁREK & HOLUŠA 2006). Recently, records in 11 squares of the faunistic grid of the Moravian-Silesian region are known.

Based on the sequence of findings, we suppose a consecutive spreading through the Moravian Gate (see Fig. 1), which forms the hypothetical bridge between the two warmer Moravian Silesian areas, the southern Moravian lowlands and the Ostravian Basin. The species probably crossed this a bit cooler area, followed by the quick expansion and establishment of many new populations in the Ostravian Basin. There are two main reasons, which could allow the penetration through the gate. (i) Climatic changes associated with higher temperatures in the last decades are generally used as an explanation of the recent expansion of this species in Western Europe (e.g. KLEUKERS et al. 1996, DECLER et al. 2000, BAH et al. 2003, LANDECK et al. 2005). (ii) The declining of large-scale farming in the Czech Republic after 1989 has been followed by the escalation of local ruderalization of abandoned fields, meadows and pastures (see also HOLUŠA 2003). As *P. falcata* requires taller herbaceous vegetation, this factor could play an important role during the dispersal through the Moravian Gate and could escalate the expansion.

Recently (2006-2007), permanent populations have also occurred in Bohemia, although we have known only several single specimens from this territory: Kunětica – Kunětická hora – hill (5960), Lázně Bohdaneč – Na Kovárně (6059), Malečice (6751) and Jaronice – hill Vráže (7051). Based on these data, several directions of the expansion can be discussed: (i) around the north-eastern edge of the Českomoravská Vrchovina highland along the town of Litomyšl, (ii) along river valleys through the Českomoravská Vrchovina highland and (iii) along the Danube River into the South Bohemian Basin (see Fig. 1). Some of these directions were probably also used during the last post glacial migration by many animal species. Although some localities in the river Jihlava valley, e.g. at Vladislav (6761), had been still considered to be the edge of termophilous southeastern (pannonian) fauna (CULEK 1996), other quite recent expansions of orthopteran species along the river valleys were documented, e.g. by HOCHKIRCH (2002).




Fig. 1: Distributional map of *Phaneroptera falcata* in the Czech Republic. Arrows mark the probable direction of spreading through the Moravian Gate and into Bohemia.

The occurrence of *Phaneroptera nana* has not been recorded from the Czech Republic up to 1992 (VLK 2002). The author published records of single specimens caught near Strachotín (7065) in 1992 and Pouzdřany (7065) in 2001. Despite the extensive faunistic research carried out in southern Moravia during 2002-2006, other specimens of *P. nana* were not observed till 2006. Finally, during 2006-2007, some specimens of *P. nana* were found in Brno (6865), Pouzdřany (7065), Drnolec (7164), Hrušovany nad Jevišovkou (7164), Mikulov (7165), Čejkovice (7167), Břeclav (7267) and Lanžhot (7267). A very abundant population was observed in the town of Mikulov, where *P. nana* was the dominant member of nightly singing orthopterans in bushes and trees. In comparison with *P. falcata*, this species is more thermophilous and the current northern limit of its range is probably not crossing the latitude of 49°15' in the Czech Republic, which also limits the biogeographical region of the Pannonicum (CULEK 1996).

P. nana probably penetrated into the Czech Republic from neighboring Slovakia and Austria in the last two decades (see Fig. 2). Most probably this species followed mainly the lowlands of the valleys of Morava and Dyje rivers that offer suitable habitats like thermophilous ruderal vegetation. But mapping of the occurrence was complicated by the requirements of this species, because it occupies deciduous trees and shrubs and may be only occasionally found on the ground (especially in the second half of October, when temperature may firstly decline to the freezing point during nights; see also HOLUŠA ET VIDLIČKA 1997). So, the species could have penetrated into Moravia earlier, without being detected. However, in the neighboring area of the Austrian Weinviertel the widespread

distribution of this species was also not detected earlier than the second half of the 1990s (archive of Orthopterenkartierung Ostösterreich).

As well as in *P. falcata*, we suppose that the probable reasons of spreading were climatic changes associated with higher temperatures as well as the increase in area of fallow fields and meadows.




Fig. 2: Distributinal map of *Phaneroptera nana* in the Czech Republic. Arrows mark the probable direction of spreading from Austria and Slovakia.

Acknowledgements

This paper was partially supported by the grant of the Ministry of Agriculture of the Czech Republic No. 0002070201. We thank Ondřej Konvička (Veselí nad Moravou, CR), Jan Dolanský (Pardubice, CR) and Bohuslav Mocek (Hradec Králové, CR) for additional information concerning the occurrence of *P. falcata* in the CR.

Authors:

Dr. Petr Kočárek

Department of Biology and Ecology, Faculty of Sciences

University of Ostrava

Chittussiho 10

CZ-710 00 Ostrava, Czech Republic

E-mail: petr.kocarek@osu.cz

Doc. Dr. Jaroslav Holuša
Forestry and Game Management Research Institute
Office Frýdek-Místek
Nádražní 2811
CZ-738 01 Frýdek-Místek, Czech Republic
Czech University of Life Science Prague
Faculty of Forestry, Wildlife and Wood Sciences
Kamýcká 129
CZ-16521 Praha – Suchdol
E-mail: holusaj@seznam.cz

Dr. Robert Vlk
Department of Biology, Faculty of Education
Masaryk University
Poříčí 7
CZ-603 00 Brno, Czech Republic
E-mail: vlk@ped.muni.cz

Mgr. Pavel Marhoul
Agency for Nature Conservation and Landscape Protection of the Czech
Republic
Nuselská 34
CZ-140 00 Praha 4, Czech Republic
E-mail: pavel.marhoul@nature.cz

D.I. Thomas Zuna-Kratky
Lange Gasse 58/20
A-1080 Wien, Austria
E-mail: thomas.zuna-kratky@blackbox.at

References

- BAH, M.P., KORSAKOVA, S., GRINGORYAN, V., HENDRICKSON, O. & BAIER, W. (2003): Interactions between climate and biological diversity. CAgM report no. 91. – World meteorological organisation, Commission for agricultural meteorology, Geneva, Switzerland, 105 pp.
- BAUR, B., BAUR, H., ROESTI, C. & ROESTI, D. (2006): Die Heuschrecken der Schweiz. – Haupt, Bern. 352 pp.
- BERG, H.-M., BIERINGER, G. & ZECHNER, L. (2005): Rote Liste der Heuschrecken (Orthoptera) Österreichs. – In: ZULKA, K.P. (Red.): Rote Listen gefährdeter Tiere Österreichs. Checklisten, Gefährdungsanalysen, Handlungsbedarf. Teil 1. Grüne Reihe des Lebensministeriums, Bd. 14/1, Wien. 167-209.
- BIERINGER, G. & ROTTER, D. (2001): Verzeichnis der österreichischen Heuschrecken-Belege (Orthoptera: Ensifera und Caelifera) der Sammlung Wilhelm Kühnelt (1905-1988). – Beiträge zur Entomofaunistik (Wien) 2: 15-47.
- BOLSHAKOV, L.V. (2006): *Phaneroptera falcata* (Poda, 1761) (Hexapoda: Orthoptera: Tettigoniidae) – a dispersing species. – Biologicheskoe Raznoobrazie Tulsogo na Rubezhe Vekov 5: 3-4.

- ČEJCHAN, A. (1981): K poznání orthopteroidního hmyzu (s. I.) ČSSR II. – Čas. Nár. Muz. Praha, Řada přírodověd. 149: 147-151.
- CHLÁDEK, F. (1993): K rozšíření kobylky *Phaneroptera falcata* (Poda), (Orthoptera, Phaneropterinae) na Třebíčsku a v Československu. – Přírodověd. Sbor. Západoslov. Muz. Třebíč 19: 101-104.
- CHLÁDEK, F. (2006): Zajímavé nálezy rovnokřídých (Insecta, Orthoptera) v ČR. – Tetrax 2 (2): 8.
- CHLÁDEK, F. & GAVLAS, V. (2004): Zajímavé nálezy rovnokřídého hmyzu (Orthoptera s.l.) na Slovensku. – Tetrax 1 (12): 69-71.
- CORAY, A. (2003): *Phaneroptera nana* Fieber, 1853 (Ensifera: Phaneropteridae) überwundet der Rhein bei Basel. – Articulata 18: 247-250.
- CULEK, M. (ed.) (1996): Biogeografické členění České republiky. – Enigma, Praha, 347 pp.
- DECLLEER, K., DEVRIESE, H., HOFMANS, K., LOCK, K., BARENBRUG, B. & MAES, D. (2000): Voorlopige atlas en "rode lijst" van de sprinkhanen en krekels van België (Insecta, Orthoptera). – Werkgroep Saltabel i.s.m. I.N. en K.B.I.N., Rapport Instituut voor Natuurbehoud 2000/10, Brussel, 76 pp.
- DOBŠIK, B. & CHLÁDEK, F. (1975): K současnému stavu znalostí o rovnokřídém hmyzu (Saltatoria) Slezska. – Ochr. Fauny Severomorav. Kraje 7: 31-50.
- FEDOR, P.J. (2000): K biodiverzite rovnokřídlovců (Caelifera a Ensifera) v okolí geotechnického systému Dunajského vodného diela. – Acta environmentalistica Universitatis Comenianae (Bratislava) 10: 171-176.
- FEDOR, P.J. (2001): Bioindikačný význam rovnokřídého hmyzu (Ensifera et Caelifera) vo vzťahu k zmenenej krajine v okolí Vodného diela Gabčíkovo. – Folia Faunistica Slovaca 6(Suppl. 1): 1-91.
- GAVLAS, V. (2001): Kobylky, koníky a modlivky Borskej nížiny. – Záhorie 10: 21-24.
- HOCHKIRCH, A. (2002): Recent range expansions of Orthoptera in North-western Germany. – Verhandlungen der Gesellschaft für Ökologie 32: 346.
- HOLUŠA, J. (2003): Výsledky faunistického průzkumu sarančí (Orthoptera: Caelifera) a kobytek (Orthoptera: Ensifera) na území Chráněné krajinné oblasti Kokořínsko. – Bezděz 12: 307-334.
- HOLUŠA, J. & VIDLIČKA, L. (1997): Orthoptera in the Devínská Kobyla hill. – Entomofauna Carpathica 9: 54-59.
- KLEUKERS, R.M.J.C., DECLLEER, K., HAES, E.C.M., KOLSHORN, P. & THOMAS, B. (1996): The recent expansion of *Conocephalus discolor* (Thunberg) (Orthoptera: Tettigoniidae) in western Europe. – Entomologist's Gazette 47: 37-49.
- KOČÁREK, P. & HOLUŠA, J. (2006): Recent expansion of bush-cricket *Phaneroptera falcata* (Orthoptera: Tettigoniidae) in northern Moravia and Silesia (Czech Republic), pp. 207-211. In: Kočárek P., Plášek V. & Malachová K. (eds): Environmental changes and biological assessment III. Scripta Facultatis Rerum Naturalium Universitas Ostraviensis Nr. 163, Ostrava, 355 pp.
- KOČÁREK, P. & NĚMEČKOVÁ, I. (2000): Orthopteroidní hmyz (Orthoptera, Blattaria & Dermaptera) CHKO Poodří a nejbližšího okolí (Česká republika). – Čas. Slez. Muz. Opava (A), 49: 111-116.
- KOČÁREK, P. & HOLUŠA, J. & VIDLIČKA, L. (2005): Blattaria, Mantodea, Orthoptera & Dermaptera of the Czech and Slovak Republics. Illustrated key 3. Blattaria, Mantodea, Orthoptera & Dermaptera České a Slovenské republiky. Ilustrovaný klíč 3. Kabourek, Zlín, 349 pp.
- KÖHLER, G. (2001): Fauna der Heuschrecken (Ensifera et Caelifera) des Freistaates Thüringen. – Naturschutz Report, Heft 17, Jena, 378 pp.

- LANDECK, I., BRUNK, I., RÖDEL, I. & VORWALD, J. (2005): Neue Nachweise der Gemeinen Sichel-schrecke *Phaneroptera falcata* (Poda 1761) für des Land Brandenburg (Saltatoria: Tettigoniidae). – Märkische Ent. Nachr. 7: 113-122.
- MAJZLAN, O. & FEDOR, P.J. (1997): Kobylyky (Ensifera) NPR Drieňová hora pri Novej Vieske sledované Malaiseho pascou. – Folia faunistica Slovaca 2: 53-55.
- MAJZLAN, O., FEDOR, P.J. & RYCHLÍK, I. (2000): Vybrané skupiny hmyzu (Coleoptera, Ensifera, Caelifera, Dermaptera, Blattodea a Mantodea) na viatych pieskoch južného Slovenska. – Rosalia (Nitra) 15: 155-174.
- NAGY, B. (1992): Role of activity pattern in colonisation by Orthoptera. – Pp. 351-363, in: Proceedings of the Fourth European Congress of Entomology and the XIII. Internationale Symposium für Entomofaunistik Mitteleuropas, 1-6 September 1991, Gödöllő, Hungary, Hungarian Natural History Museum, Budapest.
- PRUNER, L. & MÍKA, P. (1996): Seznam obcí a jejich částí v České republice s čísly mapových polí pro síťové mapování fauny. – Klapalekiana 32 (Suppl.): 1-175.
- VLK, R. (2002): First record of the bush-cricket *Phaneroptera nana nana* (Orthoptera: Tettigoniidae) in the Czech Republic. – Articulata 17: 101-102.
- SARDET, E., HUGEL, S., SCHWEBEL, L., CARRON, G. & TREIBER, R. (2005): Nouvelles observations de *Phaneroptera nana* Fieber (1853) (Orthoptera, Phaneropteridae) en Alsace et dans le reste de l'Europe occidentale et centrale. – Materiaux Orthopteriques et Entomologiques 10: 73-81.
- STOLZENBURG, U. (2005): Erneuter Nachweis der Sichel-schrecke *Phaneroptera falcata* (Poda, 1761) im Land Brandenburg (Saltatoria). – Entomologische Nachrichten und Berichte 49: 68.
- ZECHNER, L. & KOSCHUH, A. (2005): Aktuelle und historische Nachweise ausgewählter Heu-schreckenarten in der Steiermark, Österreich – Teil I: Ensifera (Saltatoria). – Joannea Zool. 7: 179-201.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Articulata - Zeitschrift der Deutschen Gesellschaft für Orthopterologie e.V. DGfO](#)

Jahr/Year: 2008

Band/Volume: [23_1_2008](#)

Autor(en)/Author(s): Kocarek Petr, Holusa Jaroslav, Vlk Robert, Marhoul Pavel, Zuna-Kratky Thomas

Artikel/Article: [Recent expansions of the bush-crickets *Phaneroptera falcata* and *Phaneroptera nana* \(Orthoptera: Tettigoniidae\) in the Czech Republic 67-75](#)