Atalanta (Mai 1992) 23(1/2):255-258, Würzburg, ISSN 0171-0079

A new species of the genus Chamaesphecia SPULER, 1910 from the Transcaucasus and Northern Iran

(Lepidoptera, Sesiidae) by OLEG G. GORBUNOV & KAREL SPATENKA received 24.II.1992

During our work on a monograph of the palaearctic Sesiidae and based on a revision of type-material from various european museums, we discovered that the paratype of *Chamaesphecia morosa* LE CERF, 1937, deposited in the Naturhistorska Riksmuseet (Stokholm), is different from the holotype of this species deposited in the collection of the Museum national d'Histoire naturelle (Paris). Such a situation threatens the stability of the nomenclature, the more so as this paratype represents a yet undescribed species wide-spread in the Transcaucasus (Armenia, Azerbaijan).

Below, we present the description of this new species and a redefinition of *Ch. morosa* and other allied species.

The types have been deposited in the following collections:

- RMS Naturhistorska Riksmuseet, Stokholm, Sweden.
- ZISP Zoological Institute, Russian Academy of Sciences, St-Petersburg, Russia.
- CG collection of Dr. O. GORBUNOV, Moscow, Russia.
- CS collection of Dr. K. SPATENKA, Prague, CSFR.

Chamaesphecia zarathustra spec. nov. (figs. 1-5)

Chamaesphecia morosa LE CERF, 1937a:175 (partim); LE CERF, 1937b: pl. 5, fig. 8.

Holotype ♂^{*}: "Perse, Elburgsgebirge, Tarsee, 2600m, 14./17.VII.1936, coll. L. SCHWINGEN-SCHUSS" – paratype of *Ch. morosa* (RMS).

Paratypes: 1 \eth , N-Iran, Afigod, leg. N. FILIPPOV (ZISP); 1 \wp , Iran, Elburs Mts., Tehran Prov. vic. Gajereh, 2400m alt., 20.VII.1976, leg. C. NAUMANN, Nr. 1685 (CS); 1 \eth , 1 \wp , USSR, Azerbaijan, Ashagi Buzgov, 39°31'N, 43°24'E, VII-VIII. 1988, e.l., leg. K. SPATENKA (CS); 1 \eth , 2 $\wp \varphi$, same locality and date, e.l., leg. O. GORBUNOV (CG); 1 \eth , USSR, Azerbaijan, Talysh Mts., Zuvand, Gilidara, 1500m alt., 6.VII.1980, leg. A. DANTCHENKO (CG); 1 \eth , 1 \wp , same locality, 5.VI.1985, e.p., leg. O. GORBUNOV (CG); 2 $\eth \eth$, 2 $\wp \varphi$, USSR, Armenia, Khosrov Nature Reserve, 1400m alt., 15.-16.VI.1988, e.p., leg. O. GORBUNOV (CG).

Description

Holotype ♂ (LE CERF, 1937b: pl. 5, fig. 8).

Body length 12.5; forewing 10.0; antenna 6.0 mm.

Head: labial palp pale yellow with a narrow black strip posteriorly, apical segment yellow; antenna black with blue lustre, with a narrow yellow strip posteriorly; frons pale yellow with

a grey-brown spot centrally; vertex black with a few orange scales; pericephalic hairs yellow-orange dorsally and white laterally.

Thorax: patagia black with bronze lustre; tegula black with bronze lustre, with a broad yellow strip anteriorly and a pale yellow spot near base of forewing; mesothorax black with a narrow yellow strip medially; metathorax yellow; in adition, all parts of thorax dorsally covered by dense, long, hairy-like greenish-yellow scales masking background colouration; pleura of thorax yellow.

Legs: completely yellow with a few black scales; spurs yellow.

Forewing: brown-black, covered by dense scales, yellow with greenish hue; transparent areas covered by dense scales, colourless with greenish lustre; external transparent area divided into five cells, about twice as broad as discal spot.

Hindwing: veins and discal spot brown-black; discal spot narrow, trapeziform, reaching to base of veins M3-Cu1.

Abdomen: brown-black, covered by dense scales, yellow with greenish hue, completely masking background colouration; tergites nos. 2, 4, 6 and 7 with a narrow, pale yellow, distal strip, anal tuft orange yellow centrally and brown-black laterally.

Female: tegula with an orange-yellow spot near base of forewing; tergites nos. 2, 4 and 6 of abdomen with a narrow, pale yellow, distal strip; all sternites and anal tuft yellow-orange.

Variability; The fresh specimens are brighter and with a more intense greenish hue in the colouration. Sometimes the males are with a narrow, pale yellow, distal strip on tergit no. 5 of the abdomen. Besides, there are slight variations in size.

Genitalia (male): tegumen-uncus complex (fig. 1) narrow; scopula androconialis small; crista gnathi lateralis long, oval; crista gnathi medialis extremely small, practically absent; valva (fig. 2) trapeziform; crista sacculi absent; sclerotized crista long and high; saccus right, slightly shorter than vinculum (fig. 3); aedoeagus (fig. 4) somewhat longerthan valva; vesica with two cornuti.

Genitalia (female) (fig. 5): 8th tergite narrow; apophysis anterior as long as apophysis posterior; ostium bursae wide, funnel-shaped; antrum thick, about half as long as apophysis anterior; ductus bursae short; bursa copulatrix oval.

Differential diagnosis: Habitually, *Ch. zarathustra* spec. nov. is very closely related to *Ch. doryceraeformis* (LEDERER, 1853), but can be easily distinguished by the yellow colouration with greenish hue of the wings and the abdomen. From *Ch. morosa* LE CERF, 1937, *Ch. sefid* LE CERF, 1938, and *Ch. ophimontana* GORBUNOV, 1991, this new species differs by the yellow colouration with greenish hue of the discal spot of the forewing (in the other species brown or dark brown).

Bionomics: The host plant of this species is *Phlomis caucasicus* (Lamiaceae). The larva lives inside a very hard and ligneous root, where it hibernates twice. After the second hibernation, approximately from the end of May until early July, the larva constructs anexit tube about 1-5 cm from the root to the surface of the soil, and then it constructs a cocoon in a gallery in the root. The imago is on the wing in July.



Figs. 1-4: Male genitalia of *Chamaesphecia zarathustra* spec. nov., paratype. 1) tegumenuncus complex; 2) valva; 3) saccus; 4) aedoeagus. Scale bar: 0.5 mm. Fig. 5: Female genitalia of *Chamaesphecia zarathustra* spec. nov., paratype. Scale bar: 0.5 mm.

Habitat: The host plant of this species grows in highland xerophytes, a typical biotope in the southern Transcaucasus and northern Iran. This habitat is distinguished by numerous perennial plants, in particular various species of Lamiaceae, Fabaceae, Euphorbiaceae etc. As a rule, such biotopes yield good collections of clearwing moths. For example, we have collected a lot of Sesiidae in similar biotopes near Buzgov and Khosrov Nature Reserve, e.g. three species of the genus *Tinthia (T. brosiformis, T. myrmosaeformis* and *T. hoplisiformis)*, *Pyropteron minianiformis*, two species of the genus *Bembecia (B. zuvandica* and *B. pontica)*, and not less than ten species of the genus *Chamaesphecia (Ch. aurifera, Ch. elampiformis, Ch. turbida, Ch. mirza*, etc.).

Acknowledgements

We express our sincere thanks to Dr. B. GUSTAFSSON (RMS) and Dr. G. LUQUET (MNHP) for loaning us the type-material of F. LE CERF. We also thank Dr. S. GOLOVATCH (Moscow) for the correction of the English, and Dr. U. EITSCHBERGER (Marktleuthen, FRG) for editing this paper.

Literature

LE CERF, F. (1937a): Aegeriidae nouvelles d'Asie Mineure. – Bull. Soc. ent. Fr. **42**:172-176. LE CERF, F. (1937b): Aegeriidae nouvelles ou peu connues d'Asie antérieure. – Zt. Öst. Ent.-Ver. **22**:81-89, 96-99, 103-109, pl. 5.

Adresses of the authors

Dr. OLEG GORBUNOV Institute of Evolutionary Morphology and Ecology of Animals Russian Academy of Sciences Leninsky prospect 33 117071 Moscow, V-71 Russia

Dr. KAREL SPATENKA Research Institute of Food Industry and Agricultural Service Entomological Laboratory CS-28911 Pecky CSFR

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: Atalanta

Jahr/Year: 1992

Band/Volume: 23

Autor(en)/Author(s): Gorbunov Oleg G., Spatenka Karel

Artikel/Article: <u>A new species of the genus Chamaesphecia Spuler, 1910</u> from the Transcaucasus and Northern Iran 255-258