

Atalanta (November 1998) 28 (3/4): 327–334, Würzburg, ISSN 0171-0079

## The species of casebearer moths described from the Volgo-Ural region

(Lepidoptera, Coleophoridae)

by

V. V. ANIKIN

received 20.I.1998

**Summary:** 22 species of Coleophoridae have been described from the Volgo-Ural region. 4 species are synonymized at present day.

**Резюме:** 22 вида чехлоносок описаны из Волго-Уральского региона. 4 вида являются синонимами на сегодняшний день.

During our work with the STAUDINGER collection material from the Volgo-Ural region, which is kept at the Humboldt-University Museum in Berlin, we investigated the type specimens of casebearer moths with labels "Origin". These are:

*Valvulalongia falcigerella* (CHR.), *Ecebalia quadrifariella* (STGR.), *Casignetella occatella* (STGR.), *Carpochena asperginella* (CHR.), and *Carpochena aequalella* (CHR.).

We also looked at the Zoological Institut of Russian Academy of Sciences (St. Petersburg, Russia) into EVERSMANN's collection of moths from Sarepta (now the vicinity of Volgograd), Orenburg and "Ural".

Having at hand our own moth material of the before cited species, gathered during the years 1986–97 in the Volgo-Ural region, and the casebearers which were described later by TOLL (1957, 1961), BALDIZZONE (1988), and FALKOVITSH (1993) on the base of old collected material (by H. CHRISTOPH, M. BARTEL, V. KUZNETSOV), and the results of our determination of V. ZOLOTUHIN's casebearer material (in ZFMK) from the vicinity of Volgograd, we have the possibility to extend our knowledge about the distribution of these species in Russia.

Taxonomic order of the Coleophoridae corresponds mainly to TOLL's [1953, 1962] scheme in addition to which we use contemporary generic names. The specimens collected by the author are kept at the Zoological Institute of the Russian Academy of Sciences (St. Petersburg) and at the University of Saratov.

The author wishes to express his gratitude to the curator of Lepidoptera of the Natural Museum of the Humboldt-University Dr. W. MEY (Berlin, Germany) for using the material of the STAUDINGER collection, to Dr. D. STÜNING (Bonn, Germany), curator of the Lepidoptera section of the Zoological Museum Alexander Koenig for help in the work with the museum material of casebearer moths, Dr. M. FALKOVITSH (Zoological Institute RAS, St. Peterburg, Russia) for using the EVERSMANN collection and for comments during the preparation of this article and Dr. V. ZOLOTUHIN (Ulyanovsk University, Russia) for help and critical remarks on the manuscript.

## Abbreviations for museums:

MHUB	Museum für Naturkunde der Humboldt-Universität, Berlin, Germany
ZFMK	Zoologisches Forschungsinstitut und Museum Alexander Koenig, Bonn, Germany
ZISP	Zoological Institut of Russian Academy of Sciences, St. Petersburg, Russia
BMHM	British Museum of Natural History, London, England
IZPC	Institute of Systematic and Experimental Zoology PAS, Cracow, Poland
ZMSU	Zoological Museum of Saratov State University, Saratov

An asterisk (\*) before a species name means, that it is synonymized at present day.

### *Orthographis uralensis* (TOLL, 1961)

*Coleophora uralensis* TOLL, Annls. Zool. Warsz., **19** (6): 217, figs.

Locus typicus: Russia, Sarepta.

Material examined: 2 ♂♂, 28.V.1992, Seleznikha, Saratov. Prov., V. ANIKIN (ZMSU); 3 ♂♂, 1.-3.V.1994, lake Elton, Volgograd Distr.; 1 ♂, 24.V.1996, vil. Akulovka, Ul'yanovsk Prov., V. ANIKIN (ZMSU); 2 ♂♂, 10.VIII.1997, Zhiguli Preserve, Samara Prov., V. ANIKIN (ZMSU).

Distribution: W Palaearctic (southern part).

Biology: Steppe biotopes. Flight period (FP) in V–VI. Larval foodplants (L): *Artemisia sericea* WEB. ex STECHM.\* (\* – original data).

### *Valvuloncia falcigerella* (CHRISTOPH, 1872)

*Coleophora falcigerella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 31, pl. 2A, fig. 27

Locus typicus: Russia, Sarepta.

Material examined: Holotype ♂, "Origin", Sarepta, GP N164; 7 specimens, Sarepta, H. CHRISTOPH (MHUB); 1 ♀, 28.VIII.1997, r. Kizan, Astrakhan Prov., V. ANIKIN (ZMSU).

Distribution: Russia (Lower Volga region), Kazakhstan, Middle Asia.

Biology: Steppe biotopes. FP in VII–VIII. L: *Glycyrrhiza glabra* L. 1 generation (G).

### *Chnoocera botarella* (HERRICH-SCHÄFFER, 1861)

*Coleophora botarella* HERRICH-SCHÄFFER, 1861, (CorrespBl. Samml. Schmett. **2**: 143.

Locus typicus: Russia, Sarepta, Ural.

Material examined. 6 specimens, Sarepta, H. CHRISTOPH (MHUB); 5 specimens, Sarepta, H. CHRISTOPH (ZISP); 11 ♂♂, 11 ♀, 8.VIII.1991, Dyakovka, Nature Reserve, Saratov Prov., V. ANIKIN (ZMSU); 1 ♀, 26.VII.1997, vil. Raslovka-2, Saratov Prov., O. SINICHKINA (ZMSU).

Distribution: Russia (Lower Volga region), Kazakhstan, Middle Asia, Iran.

Biology: Steppe biotopes. FP in VI–VIII. 1 G.

*Oedicaula serinipennella* (CHRISTOPH, 1872)

*Coleophora serinipennella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 31, fig. 32.

Locus typicus: Russia, Sarepta.

*Coleophora stefanii* JOANNIS, 1899, Bull. Soc. ent. Fr. **1899**: 331, fig.

*Coleophora novella* CHRÉTIEN, 1926, Amat. Papillons **3**: 9.

*Coleophora caliacraella* CARADJA, 1931, Mem. St. Acad. Rom. (3) **7**(8): 331.

*Coleophora caliacraella lucidella* CARADJA, 1932, Bull. Sec. Sc. Acad. Rom. **15** (1/2): 43.

*Coleophora jerichoella* AMSEL, 1935, Mitt. Zool. Mus. Berl. **20** (2): 306, fig. 55.

*Coleophora jordanella* AMSEL, 1935, Mitt. Zool. Mus. Berl. **20** (2): 306, fig. 56.

*Coleophora sosisperma* MEYRICK, 1936, Exot. Microl. **4**: 261.

*Coleophora deserticola* TOLL, 1944, Z. Wien. ent. Ges. **29**: 292, figs.

*Coleophora soffneri* TOLL, 1944, Z. Wien. ent. Ges. **29**: 292, fig. 6.

Material examined: 2 specimens, 7.-24.V.1967, vicinity Volgograd, V. ZOUHAR (ZFMK); 5 ♂♂, 2 ♀♀, 9.VIII.1991, Dyakovka, Nature Reserve, Saratov Prov., V. ANIKIN (ZMSU); 2 ♂♂, 1 ♀, 14.VIII.1991 Pallasovka, Volgograd Prov., V. ANIKIN (ZMSU); 3 ♂♂, 1 ♀, 24.-26.VI.1997, vil. N. Bannovka, Saratov Prov., V. ANIKIN (ZMSU); 2 ♀♀, 27.VIII.1997, r. Kizan, Astrakhan Prov., V. ANIKIN (ZMSU).

Distribution: All over the Palaearctic in the steppe zone.

Biology: Steppe biotopes. FP in VI-VIII. The larva in the swelling galls on the haulms of *Atriplex*. 1 G.

*Apista impalella* (TOLL, 1961)

*Coleophora impalella* TOLL, 1961, Annals. Zool. Warsz. **19** (6): 213, figs.

Locus typicus: Russia, Sarepta.

Material examined: Not examined. Only known after the type material (TOLL, 1961).

Distribution: Lower Volga region (Volgograd Prov.).

Biology: Steppe biotopes. FP in VI. 1 G. L: unknown.

*Apista lacera* FALKOVITSH, 1993

*Apista lacera* FALKOVITSH, 1993, Proc. Zool. Inst. St. Petersburg, **251**: 40, figs.

Locus typicus: Ural, Yanvarcevo.

Material examined: Holotype ♂, Ural, Yanvarcevo, 9.III.1951 (KUZNETSOV) (ZISP); paratype: 2 ♂♂, Russia, Novosibirsk Distr., Akademgorodok, 10.-20.VI.1983, K. MIKKOLA (ZISP); 1 ♂, Altay, r. Katun, 22.-27.VI.1983, K. MIKKOLA (ZISP); 1 ♀ Russia, Belgorod Distr., vil. Borisovka, 21.V.1973, A. LVOVSKY (ZISP).

Distribution: European part of Russia (Belgorod Distr.), Ural, Altay.

Biology: Steppe biotopes. FP in VII. 1 G. L: *Caragana frutex*.

\* *Multicoloria gypsophilae* (CHRISTOPH, 1872)

*Coleophora gypsophilae* CHRISTOPH, 1872, Stett. ent. Ztg. **23**: 223.

Locus typicus: Russia, Sarepta.

Material examined. In MHUB 3 specimens without abdomen collected by H. CHRISTOPH. BALDIZZONE (1994) synonymized this species with *M. vicinella* after predetermined type(?) material from the British Museum.

\* *Multicoloria paraspumosella* (TOLL, 1957)

*Coleophora paraspumosella* TOLL, 1957, Acta zool. Cracov. **2**(6): 126, figs.

Locus typicus: Russia, Sarepta.

Material examined: 1 ♀?, 22.VII.1994, Cherkaskoe, Khvalynsk Distr., Saratov Prov., V. ANIKIN (ZMSU). BALDIZZONE (1994) synonymized this species with *M. cartilaginella* after predetermined material (type?) from the TOLL collection (IZPC).

Biology: Forest-steppe biotopes. FP in VI–VII. 1 G.

*Multicoloria cartilaginella* (CHRISTOPH, 1872)

*Coleophora cartilaginella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 33, pl. 2A, fig. 30.

Locus typicus: Russia, Sarepta.

*Coleophora echinella* STAUDINGER, 1880, Horae Soc. ent. ross. **15**: 364.

*Coleophora dubiosa* TOLL, 1952, Bull. Soc. ent. Mulhouse **1952**: 43.

?*Coleophora paraspumosella* TOLL, 1957, Acta zool. Cracov. **2**(6): 126, figs.

*Coleophora medicagivora* TOLL, 1961, Schr. bayer. Akad. Wiss. Mat.-Nat. **170**(7/10): 286, figs.

Material examined: 2 specimens, Sarepta, H. CHRISTOPH (MHUB); 3 ♂♂, 2 ♀♀, 7.–24.V.1967, vicinity Volgograd, V. ZOUHAR (ZFMK); 1 ♀, 3.VI.1990, vil. Burkino, Saratov Prov., V. ANIKIN (ZMSU); 2 ♂♂, 1 ♀, 24.–26.VI.1997, vil. N. Bannovka, V. ANIKIN (ZMSU).

Distribution: W Palaearctic (southern part).

Biology: Forest-steppe biotopes. FP in VI. 1 G.

L: *Astragalus albicaulis* GROSSH., *A. glycyphylloides* L. \*, *Medicago romanica* PROD.

*Multicoloria stachi* (TOLL, 1957)

*Coleophora stachi* TOLL, 1957, Acta zool. Cracov. **2**(6): 125, figs.

Locus typicus: Russia, Sarepta.

Material examined: Not examined.

Distribution: Russia: Lower Volga region (Volgograd Prov.), Far East.

Biology: Steppe biotopes. FP in VI. 1 G. L: *Artemisia*.

*Perygra numeniella* (BALDIZZONE, 1988)

*Coleopora numeniella* BALDIZZONE, 1988, Beitr. Ent. **38**: 74, figs.

Locus typicus: Ural, Uralisk.

Material examined: Not examined.

Only known after 1 ♀ collected by MAX BARTEL (coll. Museum d'Historie de Bucarest) (BALDIZZONE, 1988).

Distribution: Russia (Volga-Ural region).

Biology: Steppe biotopes. FP in VII. 1 G.

*Ecebalia charadriella* (BALDIZZONE, 1988)

*Coleophora charadriella* BALDIZZONE, 1988, Beitr. Ent. **38**: 81, figs.

Locus typicus: Ural, Uralsk.

Material examined: 1 ♂, 22.V.1987, Saratov (ZMSU); 1 ♂, 15.V.1991, vil. Dyakovka, Nature Reserve, Saratov Prov., V. ANIKIN (ZMSU).

Distribution: Russia (Lower Volga Region, S. Ural), Central Kazakhstan.

Biology: Steppe and semi-desert biotopes. FP in VII. 1 G.

*Ecbalia attalicella* (ZELLER, 1871)

*Coleohpora attalicella* ZELLER, 1871, Stett. ent. Ztg. **1871**: 77

Locus typicus: Russia, Sarepta.

*Coleophora unistriella* CARADJA, 1920, D. ent. Z. Iris **34**: 152.

Material examined: 2 specimens, Sarepta, H. CHRISTOPH (MHUB).

Distribution: Russia (Lower Volga region), Kazakhstan (S Ural), Afghanistan.

Biology: Forest-steppe and steppe biotopes. FP in VI–mid VII. 1 G.

*Ecbalia quadrifariella* (STAUDINGER, 1880)

*Coleophora quadrifariella* STAUDINGER, 1880, Horae Soc. ent. ross. **15**: 373, fig. 45.

Locus typicus: Russia, Sarepta.

Material examined: Holotype 1 ♂, "Origin", Sarepta, GP N190 (MHUB); paratype: 1 ♀, Sarepta, GP N191 (MHUB).

Distribution: Russia (Lower Volga Region), Kazakhstan (S Ural).

Biology: Steppe biotopes. FP in VI–VII. 1 G.

*Casignetella occatella* (STAUDINGER, 1880)

*Coleophora occatella* STAUDINGER, 1880, Horae Soc. ent. ross. **15**: 376.

Locus typicus: Russia, Sarepta.

Material examined: Lectotype ♂, "Origin" Sarepta, GP N296 (MHUB); 1 ♀, 19.VIII.1968, Astrakhan Distr., PENCHUKOVSKAYA (ZISP).

Distribution: S Europe (local).

Biology: Steppe biotopes. FP in VIII.

*Casignetella tringella* (BALDIZZONE, 1988)

*Coleophora tringella* BALDIZZONE, 1988, Beitr. Ent. **38**: 81, figs.

Locus typicus: Ural, Uralsk.

Material examined: 3 ♂♂, 27.VII.1996, Engels, Saratov Prov., V. ANIKIN (ZMSU).

Distribution: Steppe zone of E Europe.

Biology: Steppe and semi-desert biotopes. FP in VII. 1 G. L: *Kochia prostrata*\*

\* *Carpochena delibutella* (CHRISTOPH, 1872)

*Coleophora delibutella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 35, pl. 2A, fig. 31.

Locus typicus: Russia, Sarepta.

Material examined: Holotype ♂, "Origin", Sarepta, GP N133; paratype ♀, Sarepta, GP N134 (MHUB). The Type material was predetermined and synonymized with *C. squalorella* (ZELLER, 1849) by TOLL (1962).

*Carpochena asperginella* (CHRISTOPH, 1872)

*Coleophora asperginella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 31, pl. 2A, fig. 28.

Locus typicus: Russia, Sarepta.

*Coleophora nigrosquamella* FILIPJEV, 1925, Rev. russe. ent. **19**: 51.

Material examined: Holotype ♂, "Origin", Sarepta, GP N6 (MHUB); 1 specimen, 7.–24.V. 1967, vicinity Volgograd V. ZOUHAR (ZFMK); 2 ♂♂, 1 ♀, 8.VIII.1991, Dyakovka, Nature Reserve, Saratov Prov., V. ANIKIN (ZMSU); 1 ♀, 26.VIII.1997, r. Kizan, Astrakhan Prov., V. ANIKIN (ZMSU).

Distribution: Russia (Lower Volga region), Kazakhstan, Middle Asia.

Biology: Desert biotopes. FP in VIII. 1 G. The larva lives on the carpels of *Coryspermum*.

\* *Carpochena sareptella* (TOLL)

*Coleophora sareptella* TOLL, in litt. nom. nudum.

Locus typicus: Russia, Sarepta.

Material examined: Holotype ♀, Sarepta, GP N132 (MHUB). We predetermined the type material and synonymized it with *C. salicorniae* (HEINEMANN & WOCKE, 1877).

*Carpochena trientella* (CHRISTOPH, 1872)

*Coleophora trientella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 37, pl. 2A, fig. 33.

Locus typicus: Russia, Sarepta.

Material examined: 1 specimen, Sarepta, H. CHRISTOPH (MHUB); 1 ♂, 7.–24.V.1967, vicinity Volgograd, V. ZOUHAR (ZFMK); 1 ♂, 3 ♀♀, 8.VIII.1991, vil. Dyakovka, Nature Reserve, Saratov Distr., V. ANIKIN (ZMSU).

Distribution: S Palaearctic, from Hungary to the Far East.

Biology: Desert biotopes, salt-marshes. FP in VIII. 1 G. The larva lives on the carpels of *Coryspermum*.

*Carpochena aequalella* (CHRISTOPH, 1872)

*Coleophora aequalella* CHRISTOPH, 1872, Horae Soc. ent. ross. **9**: 32, pl. 2A, fig. 29.

Locus typicus: Russia, Sarepta.

*Coleophora heratella* TOLL & AMSEL, 1967, Beitr. naturk. Forsch. SüdwDtL. **26** (3): 11, figs. 16, 43.

Material examined: Lectotype ♂, "Origin", Sarepta, GP N9 (MHUB); paratype ♀, Sarepta, GP N8 (MHUB).

Distribution: Russia (Lower Volga region), Kazakhstan, Middle Asia, Iran, Afghanistan.

Biology: Steppe and half-desert biotopes. FP in V, VI. 1 G.

*Klinzigedia phlomidella* (CHRISTOPH, 1862)

*Coleophora phlomidella* CHRISTOPH, 1862, Stett. ent. Ztg. **23**: 222.

Locus typicus: Russia, Sarepta.

Material examined: 3 specimens, Sarepta, H. CHRISTOPH (MHUB); 2 ♂♂, 1 ♀, 9.VIII.1993, vil. Dyakovka, Saratov Prov., V. ANIKIN (ZMSU); 1 ♀, 26.VIII.1997, vil. Raslovka-2, Saratov Prov., V. ANIKIN (ZMSU).

Distribution: Russia (Lower Volga region), Middle Asia (mountains), Asia Minor.

Biology: Steppe biotopes. VIII. FP in 1 G. L: *Phlomis*.

## References

- BALDIZZONE, G. (1988): Contributions a la connaissance des Coleophoridae XLVIII. Guadre nouvelles especies du genre *Coleophora* HÜBNER de l'URSS. – Beitr. Ent. **38**: 74–82.
- BALDIZZONE, G. (1994): Coleophoridae dell'Area Irano-Anatolica e regioni limitrofe (Lepidoptera). – Memorie Associazione Naturalistica Piemontese **3**: 423 pp.
- CHRISTOPH, H. (1862): Vier neue südrussische Schmetterlinge. – Stett. Ent. Zeit. **23**: 220–224.
- CHRISTOPH, H. (1872): Neue Lepidoptera des Europäischen Faunengebietes. – Horae Soc. Ent. Ross. **9**: 3–39, pl. 1, 2A.

- FALKOVITSH, M. I. (1993): New species of casebearer moths (Lepidoptera, Coleophoridae) from Altai. – Proc. Zool. inst. Acad. Sci. of USSR **251**: 40–52 (in russian).
- HERRICH-SCHÄFFER, G. A. W. (1861): Revision der europäischen Schmetterlingsfauna. – Korrespondenzblatt für Sammler von Insekten, Regensburg **2**: 189 pp.
- STAUDINGER, O. (1880): Lepidopteren-Fauna Kleinasiens. – Horae Soc. ent. ross. **15**: 369–435.
- TOLL, S. (1957): Studien über die Genitalien einiger Coleophoriden, XIV, Lepidoptera. – Acta Zool. Cracov. **2** (6): 119–148. 41 figs.
- TOLL, S. (1961): Studien über die Genitalien einiger Coleophoridae, XVIII (Lepidoptera). – Ann. Zool. **19** (6): 209–277 43 figs.
- TOLL, S. (1962): Materialien zur Kenntnis der paläarktischen Arten der Familie Coleophoridae (Lepidoptera). – Acta Zool. Cracoviensia **7** (16): 577–720, 133 pls.
- ZELLER, H. (1871): Lepidopterologische Beobachtungen im Jahre 1870. – Stett. Ent. Ztg. **32**: 49–81. Pl. 2.

Address of the author

V. V. ANIKIN

Department of ecology and morphology of animals  
Biology faculty of Saratov University  
Astrakhanskay Str. 83  
410026, Saratov  
Russia

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Atalanta](#)

Jahr/Year: 1997-1998

Band/Volume: [28](#)

Autor(en)/Author(s): Anikin Vasily Victorovich

Artikel/Article: [The species of casebearer moths described from the Volgo-Ural region \(Lepidoptera, Coleophoridae\) 327-334](#)