

Atalanta (May 1999) 29(1/4):295–336, Würzburg, ISSN 0171-0079

**"Fauna Lepidopterologica Volgo-Uralensis" 150 years later:
changes and additions. Part 4. Coleophoridae, Gelechiidae,
Symmocidae and Holcopogonidae**

(Insecta, Lepidoptera)
by

VASILY V. ANIKIN, SERGEY A. SACHKOV & VADIM V. ZOLOTUHIN

received 8.XII.1998

Summary: 415 species of Coleophoridae, Gelechiidae, Symmocidae and Holcopogonidae are listed for the modern fauna of the Volgo-Ural area. 5 species of Coleophoridae are deleted from the previous lists as synonyms (= *gypsophilae* CHRISTOPH, 1862; = *paraspumosella* TOLL, 1957; = *unicolorella* TOLL, in litt.; = *delibutella* CHRISTOPH, 1872; *soreptella* TOLL, in litt.).

This paper is the fourth in a series of publications¹ dealing with the composition of the present day fauna of the Gelechioids in the Lower and Middle Volga and the south-western Cisurals. This region comprises the administrative divisions of Astrakhan, Volgograd, Saratov, Samara, Uljanovsk, Orenburg, Uralsk and Atyraus (= Gurjev) Districts, together with Bashkiria.

As in the previous parts of this series only material reliably labelled was accepted, and the time span used for this study are the last 20 years. The main collections are those of the authors: V. ANIKIN (Saratov, Astrakhan and Volgograd Districts), S. SACHKOV (Samara District) and V. ZOLOTUHIN (Uljanovsk district and southern Tataria).

All the data from the 19th and early 20th century were taken into account but only as a reference (HERRICH-SCHÄFFER, 1856, 1861; CHRISTOPH, 1862, 1872, 1888; STAUDINGER, 1870, 1879, 1880; STAUDINGER & WOCKE, 1871; ZELLER, 1871; REBEL, 1901). Whilst compiling this list we also took advantage of the information from recent papers on this region (MARTYNova, 1952; TOLL, 1957, 1961; POVOLNY, 1966; SACHKOV, 1995; ANIKIN, 1995, 1998a, 1998b; ANIKIN & PISKUNOV, 1995, 1996; ANIKIN & FALKOVITSH, 1997; ANIKIN & SACHKOV, 1998). The collection material of the Zoological Institute of the Russian Academy of Sciences at St.-Petersburg, Moscow State University, Zoologisches Museum an der Humboldt Universität (Berlin) and Museum Alexander Koenig (Bonn) were examined for our study.

We owe special thanks to the curators of the lepidoptera collections at the institutions and museums listed above – namely Dr. W. MEY (Berlin), Dr. D. STÜNING (Bonn) and Dr. A. Lvovsky (St.-Petersburg), for their help in our work with the museum funds. Cordial thanks to Dr. M. FALKOVITSH (St.-Petersburg, Russia) and Dr. V. PISKUNOV (Vitebsk, Byelorussia) for their valuable advices concerning the taxonomy and nomenclature of the Coleophoridae and Gelechiidae accordingly.

1 This series started in Atalanta (1993) 24(1/4): 89–120.

The private collection of Mr. D. KOMAROV (Volgograd) could be studied, to whom we also express our sincere thanks.

A general checklist of the casebearers and gelechiids from the Volga-Ural region is given below. Taxonomic order of the Coleophoridae corresponds mainly to TOLL's (1953, 1962) scheme in addition to which we use contemporary generic names, and of the Gelechiidae mainly to LERAUT's (1980) scheme with some additions and remarks after the works of HODGES (1986), POVOLNY (1989) and PISKUNOV (1990). The Latin names of plants were taken according to CZEZEPANOV (1995).

For the ease of use, information is given in the form of a table, with the principal data on all species mentioned for the Volgo-Ural region. Many localities have been renamed during the last 150 years, the most important ones being listed below:

Samara – later Kuibyshev – now Samara

Simbirsk – now Uljanovsk

Sarepta – now a suburb of Volgograd

Zarizyn or Tsarizyn – later Stalingrad – now Volgograd.

Note: Spassk, usually interpreted as EVERSMANN's estate not far from Orenburg, really might be also a town being flooded under Volga's water during the erection of hydroelectrostations and following increasing of waters levels, before that Spassk had been situated about 82 km ESE of Kasan on the left bank of the Volga.

Notes on the table and maps

Column 1: Species number

species is deleted from the list

species needs guarding

Column 2: Species name

Column 3: Species listed by EVERSMANN (1844) within the regional limits of that paper

Column 4–10: Administrative units

4 Astrakhan District (centre is Astrakhan)

5 Volgograd District (Volgograd)

6 Saratov District (Saratov)

7 Samara District (Samara)

8 Uljanovsk District (Uljanovsk)

9 Bashkiria (Ufa)

10 Uralsk District (Uralsk)

+ species is present

– species not found during this study

? species is known from old or doubtful data

o type locality

ø species now unknown in its type locality

Column 11: Flight periods

IV-XI - months

b, m, e - beginning, middle, end of month

1 (2) G - species develops 1 (2) generation(s)

Column 12: Comments and larval foodplants

L larval foodplants, *indicating original data

TL type locality

E E. EVERSMANN

ZMHUB - Zoologisches Museum der Humboldt Universität (Berlin)

ZISP - Zoological Institut of Russian Academy of Sciences, St.-Petersburg, Russia

BMHM - British Museum of Natural History, London, England.

On the maps, the filled circles indicate the places in which the species concerned is to be found. Hollow circles indicate those places in which the species concerned have not been found during the last 15–50 years, possibly having disappeared due to the changes in the environment. A plot size we accepted is 30x30 km.

N	Species	Flight period										Comments
		E	A	V	S	S	U	B	U			
1	2	3	4	5	6	7	8	9	10	11		12

Coleophoridae

- | | | |
|---|-----------------------|--|
| 1. <i>Augasma aeratella</i> (ZELLER, 1839) | + eVI-mVIII
in 1 G | It is typical of opened landscapes of forest-steppe and steppe biotopes. The larva lives in fusiform galls on stems of <i>Polygonum aviculare</i> *, <i>P. arenarium</i> , <i>P. lapathifolium</i> . |
| 2. <i>Augasma atraphaxidellum</i> KUZNEsov, 1957 | - V
In 1 G | Semi desert biotopes. L: <i>Atraphaxis spinosa</i> . Noted only as galls. |
| 3. <i>Papyrosiphia zhusguni</i> FALKOVITSH, 1972 | - V
In 1 G | Desert biotopes. L: <i>Calligonum aphyllum</i> * |
| 4. <i>Metriotes lutarea</i> (HAWORTH, 1828) (= <i>modestella</i> DUPONCHEL, 1839) | - eIV-VI
in 1 G | Forest biotopes. L: <i>Stellaria holostea</i> . |
| 5. <i>Casas albella</i> (THUNBERG, 1788) (= <i>leucapennella</i> HÜBNER, 1796) | + VI-VII
in 1 G | Was cited by E as <i>Leucapennella</i> . Forest-steppe biotopes. L: <i>Viscaria vulgaris</i> , <i>Silene nutans</i> . |
| 6. <i>Paravalvia spiraeella</i> (REBEL, 1916) | - V-VI
in 1 G | Noted the cases from the vicinity of Saratov on Spiraeae. |

1	2	3	4	5	6	7	8	9	10	11	12
7.	<i>Postvinculia lutipennella</i> (ZELLER, 1838)			+	+	+	+	+	+	VI-VII in 1 G	Forest biotopes. L: <i>Quercus</i> *
8.	<i>Orghidania gryphipen- nella</i> (HÜBNER, 1796)			-	eV-VI in 1 G						Forest-steppe biotopes. L: <i>Rosa canina</i> .
9.	<i>Scleriductia ochripennella</i> (ZELLER, 1849)					VI-mVII in 1 G					Forest-steppe biotopes. L: <i>Ballota, Betonica, Lamium, Stachys</i> .
10.	<i>Frederickoenigia flavipen- nella</i> (DUPONCHEL, 1843)			+	+	+	+	+	-	eVI-VII in 1 G	Forest biotopes. L: <i>Quercus</i> *
11.	<i>Haploptilia serratella</i> (LINNÆUS, 1761) (= <i>fusce- dinella</i> ZELLER, 1849)			+	+	+	+	+	+	VI-VII in 1 G	Forest biotopes. L: <i>Alnus, Betula, Ulmus</i> .
12.	<i>Haploptilia prunifolia</i> (DOETS, 1944)					-	VI-VII in 1 G				Forest and forest-steppe biotopes. L: <i>Prunus spinosa, Cerasus frutescens</i> *
13.	<i>Haploptilia spinella</i> (SCHRANK, 1802) (= <i>cera- sivorella</i> PACKARD, 1870, = <i>serratella</i> auct.)					-	mVI-VII in 1 G				Forest and forest-steppe biotopes. L: ligneous Rosaceae (<i>Cerasus, Crataegus, Malus</i> *, <i>Pirus, Prunus, Sorbus</i>).
14.	<i>Haploptilia hemerobiella</i> (SCOPOLI, 1763)					-	eVI- mVIII in 1 G				Was cited by E as <i>Anseripennella</i> . Forest and forest-steppe biotopes. L: ligneous Rosaceae (<i>Crataegus, Pirus, Prunus, Sorbus, Cotoneaster</i>).
15.	<i>Systrophoea siccifolia</i> (STAINTON, 1856)					-	VI-VII in 1 G				Forest-steppe biotopes. L: different trees - <i>Ulmus, Betula, Tilia, Sorbus, Crataegus</i> .
16.	<i>Dumitrescumia cecido- phorella</i> (OUDJEANS, 1972) (= <i>icterella</i> TOLL, 1949)					-	VI-VII in 1 G				Forest-steppe biotopes. L: galls of <i>Falllopia convolvulus</i> and <i>F. dumetorum</i> .
17.	<i>Dumitrescumia hydro- lopathella</i> (HERING, 1921)					-	VII-VIII in 1 G				Near the water biotopes, beside rivers. L: <i>Rumex hydrolathum</i> .
18.	<i>Kasyfia orbitella</i> (ZELLER, 1849)					+	mV-VI in 1 G				Was cited by E as <i>Leucapennella</i> . Forest and forest-steppe biotopes. L: <i>Alnus, Betula, Corylus</i> .
19.	<i>Kasyfia binderella</i> (KOLLAR, 1832)						V-VI in 1 G				Only known old material from Sarepta (ZMHUB). L: <i>Alnus, Corylus</i> .
20.	<i>Tollsia violacea</i> (STROM, 1783) (= <i>hornigi</i> TOLL, 1952)					-	VI-VII in 1 G				Forest and forest-steppe biotopes. L: <i>Betula, Alnus, Carpinus, Corylus, Myrica, Ulmus, Spiraea, Filipendula, Rubus, Rosa, Fragaria, Potentilla, Crataegus, Prunus, Tilia, Rhamnus, Viburnum</i> .
21.	<i>Agapalsa lusciniapennella</i> - (TREITSCHKE, 1833) (= <i>vimi- netella</i> ZELLER, 1849)					+	VI-VII in 1 G				Only known from Ural river (MARTYNOVA, 1952). Forest and inundated biotopes. L: <i>Salex caprea, S. cinerea</i> .
22.	<i>Agapalsa idaeella</i> (HOFMANN, 1869)					-	mV-VI in 1 G				Forest-steppe biotopes. L: <i>Vaccinium vitis-idaea</i> .

1	2	3	4	5	6	7	8	9	10	11	12
23.	<i>Rhamnia ahenella</i> (HEINEMANN & WOCKE, 1877)				+	+	+		VI-IX in 2 G	Forest and steppe-forest biotopes. L: <i>Geum</i> , <i>Potentilla</i> , <i>Rubus</i> , <i>Lonicera</i> , <i>Rhamnus</i> , <i>Symporicarpus</i> , <i>Viburnum</i> , <i>Alnus</i> , <i>Corylus</i> .	
24.	<i>Quadratia fuscocuprella</i> (HERRICH-SCHAFFER, 1854)						VI		in 1 G	Only known from mines (GALASIEVA, 1986). L: <i>Alnus</i> , <i>Carpinus</i> , <i>Corylus</i> .	
25.	<i>Protocryptis sibiricella</i> FALKOVITSH, 1972 (= <i>sibirica</i> FALKOVITSH, 1964, nom. praeocc.)				+	+	+	+	+ VI-VII in 1 G	Only in the planting larch forests in the region. L: <i>Larix</i> *	
26.	<i>Orthographis uralensis</i> (TOLL, 1961) (fig. 1)					o	V-VI in 1 G			TL: Uralsk. Steppe biotopes. L: <i>Artemisia sericea</i> *	
27.	<i>Orthographis ptarmicia</i> (WALSINGHAM, 1910) (fig. 1)					-	VI-VII in 1 G			Forest-steppe biotopes. L: <i>Achillea millefolium</i> *, <i>A. ptarmica</i> .	
28.	<i>Orthographis virgatella</i> (ZELLER, 1849) (fig. 1)					+	VI-VII in 1 G			From Spassk only known from old material of STAUDINGER's collection (ZMHUB). L: <i>Salvia</i> , <i>Stachys</i> , <i>Achillea</i> , <i>Globularia</i> .	
29.	<i>Halvalbia lineolea</i> (HAWORTH, 1828) (= <i>crocogrammos</i> ZELLER, 1849)					-	eVI-VII in 1 G			Forest and forest-steppe biotopes. L: <i>Ballota</i> , <i>Stachys</i> , <i>Lamium</i> .	
30.	<i>Polystrophia calligoni</i> (FALKOVITSH, 1972) (fig. 1)						V-VIII in 1 G			Desert-steppe biotopes. L: on the carpels of <i>Callygonum</i> *	
31.	<i>Ascleriducta lithargyrinella</i> (ZELLER, 1849) (= <i>solitariella</i> HERRICH-SCHAFFER, 1855)						eVI-eVII in 1 G			Forest-steppe biotopes. L: <i>Cerastium</i> , <i>Stellaria</i> .	
32.	<i>Aporiptura ochroflava</i> (TOLL, 1961)						- V, VIII-IX in 2 G			Desert-steppe biotopes, salt-marches. L: <i>Atriplex nitens</i> *, <i>A. verrucifera</i> *	
33.	<i>Aporiptura klimeschiella</i> (TOLL, 1952) (fig. 2)						- VI, VIII in 2 G			Steppe and desert-steppe biotopes. L: <i>Salsola australis</i> * and other contiguous species.	
34.	<i>Aporiptura eurasistica</i> (BALDIZZONE, 1989) (fig. 2)						- VI, IX in 2 G			Desert-steppe and steppe biotopes. L: <i>Kochia prostrata</i> *	
35.	<i>Aporiptura lonchodes</i> FALKOVITSH, 1994 (fig. 2)						- eV-VI in 1 G			Desert-steppe and desert biotopes. L: <i>Suaeda physophora</i> *	
36.	<i>Aporiptura physophorae</i> FALKOVITSH, 1994 (fig. 2)						- V-VI in 1 G			Desert-steppe biotopes. L: <i>Suaeda physophora</i> *	
37.	<i>Aporiptura dissecta</i> FALKOVITSH, 1989 (fig. 2)						IX in 1 G			Desert-steppe and desert biotopes. L: <i>Halocnemum strobilaceum</i> *	
38.	<i>Aporiptura macilenta</i> FAKKOVITSH, 1972						- V-VI in 1 G			Desert biotopes. L: <i>Ceratooides papposa</i> *	
39.	<i>Aporiptura hypoxantha</i> FALKOVITSH, 1982						- V-VI in 1 G			Desert biotopes. L: <i>Kalidium foliatum</i> *	
40.	<i>Aporiptura nigridorSELLA</i> (AMSEL, 1935) (fig. 2)						VIII in 1 G			Dry steppe biotopes.	

1	2	3	4	5	6	7	8	9	10	11	12
41. <i>Amselghia argyrella</i> (HERRICH-SCHÄFFER, 1856)		+ o +		VII in 1 G							TL: Sarepta. Forest-steppe and steppe biotopes. L: <i>Althagi pseudo-alhagi</i> .
42. <i>Amselghia fringillella</i> (ZELLER, 1839)		±		VI in 1 G							Noted from Sarepta by CHRISTOPH (1872), there the presence of this species needs to be confirmed.
43. <i>Ardania trifariella</i> (ZELLER, 1849)				VI-VII in 1 G							Forest-steppe biotopes. L: <i>Cytisus ruthenicus*</i> , <i>Genista</i> , <i>Sarothamnus</i> .
44. <i>Ardania saturatella</i> (STAINTON, 1850)				VI-VII in 1 G							Forest-steppe biotopes. L: <i>Cytisus*</i> , <i>Genista</i> .
45. <i>Valvuloniga falcigerella</i> (CHRISTOPH, 1872)			-	VII-VIII in 1 G							TL: Sarepta. Steppe biotopes. L: <i>Glycyrrhiza glabra</i> .
46. <i>Damophila mayrella</i> (HÜBNER, 1813) (= <i>spissicornis</i> HAWORTH, 1828)	-	+	+	+	+	+	+	+	VI in 1 G		Wet meadows, edges of forest, meadow-steppe. L: on the carpels of <i>Trifolium</i> .
47. <i>Damophila deauratella</i> (LIENIG & ZELLER, 1846)	+	+	+	+	+	+	+	+	eV-VI, VII in 1-2 G		Biotopes and food plants as for the previous species.
48. <i>Damophila alcyonipennella</i> (KOLLAR, 1832) (= <i>cupariella</i> ZELLER, 1847)	-	+	+	+	+	+	+	+	mV-VIII in 1-2 G		Biotopes and food plants as for the previous species.
49. <i>Damophila frischella</i> (LINNAEUS, 1758) (= <i>auronitella</i> TOLL, 1962)			-	VI-VII in 1 G							Forest-steppe biotopes. L: on the carpels of <i>Trifolium</i> .
50. <i>Damophila trifolii</i> CURTIS, 1832 (= <i>melilotella</i> SCOTT, 1861)				VI-VII in 1 G							Forest-steppe biotopes. L: <i>Melilotus altissimus</i> , <i>M. officinalis*</i>
51. <i>Calcomarginia ballotella</i> (FISCHER VON RÖSLERSTAMM, 1839)			-	VI-VII in 1 G							Forest-steppe biotopes. L: <i>Lamiaceae</i> (<i>Ballota</i> , <i>Lamium</i> , <i>Teucrium</i> , <i>Marrubium</i> , <i>Stachys*</i>).
52. <i>Chnoocera botarella</i> (HERRICH-SCHÄFFER, 1861) (fig. 3)			o	VI-VIII in 1 G							TL: Sarepta, Ural. Steppe biotopes.
53. <i>Sympypoda parthenica</i> (MEYRICK, 1891) (= <i>cyanipennella</i> TOLL, 1956, = <i>transcaspica</i> TOLL, 1959)			-	VI-VIII in 1 G							Steppe and desert-steppe biotopes. L: <i>Salsola</i> .
54. <i>Oedicaula serinipennella</i> (CHRISTOPH, 1872) (fig. 3)			+	VI-VIII in 1 G							TL: Sarepta. Steppe and desert-steppe biotopes. L: in the swelling galls on stems of <i>Atriplex</i> .
55. <i>Coleophora anatipennella</i> (HÜBNER, 1796) (= <i>beroulliella</i> GOEZE, 1783)			+	VI, VII-VIII in 2 G							Forest and forest-steppe biotopes. L: different trees (<i>Cerasus</i> , <i>Malus</i> , <i>Prunus</i> , <i>Alnus</i> , <i>Betula</i> , <i>Corylus</i> , <i>Crataegus</i> , <i>Populus</i> , <i>Quercus</i> , <i>Salix</i> , <i>Tilia</i>).
56. <i>Coleophora albidella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				VI-VII in 1 G							Forest-steppe biotopes. L: <i>Salix caprea</i> , <i>S. cinerea</i> .

1	2	3	4	5	6	7	8	9	10	11	12
57.	<i>Coleophora kuehnella</i> (GOEZE, 1783) (= <i>pallipennella</i> TREITSCHKE, 1833)		+ +		+ VI-VII in 1 G					Forest and forest-steppe biotopes. L: <i>Quercus robur</i> *	
58.	<i>Coleophora ibipennella</i> ZELLER, 1849 (= <i>nemorum</i> HEINEMANN, 1854)				VI-VII in 1 G					Forest and forest-steppe biotopes. L: <i>Quercus robur</i> *, <i>Ulmus</i> .	
59.	<i>Coleophora betulella</i> HEINEMANN & WOCKE, 1877				VI-VII in 1 G					Forest biotopes. L: <i>Betula pubescens</i> *	
60.	<i>Coleophora zelleriella</i> HEINEMANN, 1854 (= <i>pan-nicella</i> GOZMANY, 1956; = <i>platyphyllae</i> OKU, 1965)				- V-VI in 1 G					Forest and forest-steppe biotopes. L: <i>Salix caprea</i> , <i>S. cinerea</i> .	
61.	<i>Coleophora curruccinella</i> ZELLER, 1839 (= <i>nemorum</i> HEINEMANN, 1854; = <i>cristinae</i> CAPUSE, 1971)				- VI-VII in 1 G					Forest and forest-steppe biotopes. L: <i>Quercus robur</i> *, <i>Corylus</i> .	
62.	<i>Phagolamia auricella</i> (FABRICIUS, 1794) (= <i>paucinotella</i> TOLL, 1961)				- VI-VII in 1 G					Forest and forest-steppe biotopes. L: <i>Stachys recta</i> *, <i>Betonica officinalis</i> .	
63.	<i>Phagolamia virgatella</i> (ZELLER, 1849)				+ VI-VII in 1 G					Forest and forest-steppe biotopes. L: <i>Salvia</i> , <i>Stachys</i> , <i>Achillea</i> , <i>Globularia</i> .	
64.	<i>Apista impalella</i> (TOLL, 1961)				- VI in 1 G					Known from TL: Sarepta. Steppe biotopes. L: unknown.	
65.	<i>Apista lacera</i> FALKOVITSH, 1993				o VII in 1 G					TL: Yanvartsevo of Uralsk Distr. Known only after type material. L: <i>Caragana frutex</i> .	
66.	<i>Apista gallipennella</i> (HÜBNER, 1796)				- VI-VII in 1 G					Forest-steppe biotopes. L: <i>Astragalus glycyphyllos</i> .	
67.	<i>Razowskia coronillae</i> (ZELLER, 1849)				- VI-mVII in 1 G					Forest-steppe biotopes. L: <i>Coronilla</i> *	
68.	<i>Multicoloria astragalella</i> (ZELLER, 1849)				+ VI-VII in 1 G					Noted from the Ural river by MARTYNOVA (1952) but erroneously. Known from Guberli and Cisural (REZNIK, 1976). Forest-steppe and steppe biotopes. L: <i>Astragalus</i> .	
	- <i>Multicoloria gypsophilae</i> (CHRISTOPH, 1862)									TL: Sarepta. In ZMHUB are kept 3 examples collected by CHRISTOPH BALDIZZONE (1994) synonymized this species with <i>M. vicinella</i> after redetermination of the type(?) material in BMNH.	
69.	<i>Multicoloria tshiligella</i> REZNIK, 1976				o eVI-VII in 1 G					TL: Guberli (CHRISTOPH collection, ZISP). From Zhiguli preserv known as the case from L: <i>Caragana frutex</i> .	
70.	<i>Multicoloria vicinella</i> (ZELLER, 1849)				eVI-VII in 1 G					Local in steppe and forest-steppes. L: <i>Medicago</i> .	

1	2	3	4	5	6	7	8	9	10	11	12
71. <i>Multicoloria cracella</i> (VALLOT, 1835)		+ +		VII in 1 G	Forest-steppe biotopes. L: <i>Vicia cracca</i> .						
<i>Multicoloria paraspumosella</i> (TOLL, 1957)					BALDIZZONE (1994) synonymized this species with <i>M. cartilaginella</i> after redetermination of the type material in BMNH. LT for <i>paraspumosella</i> TOLL is Sarepta.						
72. <i>Multicoloria spumosella</i> (STAUDINGER, 1859)		+ VII in 1 G		Steppe biotopes. Only known from Guberli (REZNIK, 1977).							
73. <i>Multicoloria cartilaginella</i> (CHRISTOPH, 1872) (= <i>echinella</i> STAUDINGER, 1880; = <i>paraspumosella</i> TOLL, 1957)		- eVI-VII in 1 G		TL: Sarepta. Forest-steppe biotopes. L: <i>Astragalus albicaulis</i> , <i>A. glycyphyllos</i> *, <i>Chamaecytisus rutenicus</i> *, <i>Medicago romanica</i> .							
74. <i>Multicoloria eremosparti</i> FALKOVITSH, 1974		- VI-VII in 1 G		Semi-desert biotopes. L: <i>Eremosparton aphyllum</i> *							
75. <i>Multicoloria vibicella</i> (HÜBNER, [1813]) (= <i>vibicennella</i> TREITSCHKE, 1833)		+ VI-VI in 1 G		Was cited by E as <i>Vibicennella</i> . Forest-steppe biotopes. L: <i>Genista tinctoria</i> *							
76. <i>Multicoloria changaica</i> REZNIK, 1975 (fig. 4)		- V, IX in 1-2G		Steppe biotopes. L: <i>Artemisia nigra</i> *, <i>A. pauciflora</i> *							
77. <i>Multicoloria fuscociliella</i> (ZELLER, 1849) (= <i>medicaginis</i> HERRICH-SCHAFFER, 1861)		- VI in 1 G		Forest-steppe and steppe biotopes. L: <i>Medicago Coronilla</i> .							
78. <i>Multicoloria singreni</i> (FALKOVITSH, 1973) (fig. 4)		- VII-VIII in 1 G		Steppe and desert-steppe biotopes. L: <i>Astragalus varius</i> *							
79. <i>Multicoloria caelebi-pennella</i> (ZELLER, 1839)		+ eVI-VIII in 1 G		Noted from Ural by REZNIK (1976). Forest-steppe biotopes. L: <i>Artemisia campestris</i> , <i>Gnaphalium</i> , <i>Heli-chrysum</i> .							
80. <i>Multicoloria solenella</i> (STAUDINGER, 1859)		+ VII in 1 G		Noted from Guberli by E as <i>Gallipennella</i> .							
81. <i>Multicoloria conspicuella</i> (ZELLER, 1849) (= <i>similis</i> STAUDINGER, 1880)		+ VII-VIII in 1 G		Noted from Guberli by REZNIK (1976). Forest-steppe and steppe biotopes. L: <i>Aster</i> , <i>Achillea</i> , <i>Centaurea</i> , <i>Linosyris</i> .							
82. <i>Multicoloria incostans</i> REZNIK, 1975 (fig. 4)		mV-VI in 1 G		Desert-steppe biotopes.							
83. <i>Multicoloria stachi</i> (TOLL, 1957)		- VI		Only known from TL: Sarepta. Steppe biotopes. L: <i>Artemisia</i> .							
84. <i>Multicoloria pseudoditella</i> - (BALDIZZONE & PATZAK, 1983)		VI-VII in 1 G		Rare in steppes. L: <i>Artemisia</i> .							
85. <i>Multicoloria ditella</i> (ZELLER, 1849)		+ VI-VII in 1 G		Forest-steppe and steppe biotopes. L: <i>Artemisia</i> .							

1	2	3	4	5	6	7	8	9	10	11	12
86.	<i>Multicoloria vibicigerella</i> (ZELLER, 1839) (= <i>didyma</i> TOLL, 1957)		+ + +				+ V-VI in 1 G			Steppe and semidesert biotopes. L: <i>Artemisia campestris</i> *, <i>Achillea millefolium</i> .	
87.	<i>Multicoloria partitella</i> (ZELLER, 1849)						VI-VII in 1 G			Forest-steppe and steppe biotopes. L: <i>Artemisia absinthium</i> .	
88.	<i>Suireia milvipennis</i> (ZELLER, 1839)						VI in 1 G			Forest biotopes, near the sphagnum bogs.	
89.	<i>Suireia badiipennella</i> (DUPONCHEL, 1843)						+ VI-VII in 1 G			Forest-steppe biotopes. L: <i>Ulmus</i> , <i>Corylus</i> , <i>Prunus spinosa</i> , <i>Fraxinus</i> , <i>Acer</i> .	
90.	<i>Suireia limosipennella</i> (DUPONCHEL, 1842)						V-VI in 1 G			Known from mines only (GALASJEVA, 1986). L: <i>Ulmus</i> .	
91.	<i>Argyraclinia kautzi</i> (REBEL, 1933)						- VIII in 1 G			Rare in desert and semi-desert biotopes. The first data from Russia.	
92.	<i>Argyraclinia necessaria</i> (STAUDINGER, 1880)						- VIII-IX in 1 G			Desert biotopes. L: <i>Allyssum turkestanicum</i> *	
93.	<i>Argyraclinia ochrea</i> (HAWORTH, 1828)						- VI, VIII in 1-2 G			Rare and local in forest-steppes. L: <i>Helianthemum</i> .	
94.	<i>Eupista ornatipennella</i> (HÜBNER, 1796)	+ + + + + + +					+ VI-VII in 1 G			Forest-steppe biotopes. L: in the carpels of Lamiaceae, then on Poaceae (<i>Holcus</i> , <i>Briza</i> , <i>Dactylis</i> , <i>Bromus</i>).	
95.	<i>Eupista lixella</i> (ZELLER, 1849)						- VII-VIII in 1 G			Steppe biotopes with various steppe-herbaceous plants. L: the young larva on <i>Thymus</i> , then on Poaceae (<i>Anthoxanthum</i> , <i>Holcus</i> , <i>Alopecurus</i> , <i>Avena</i> , <i>Koeleria</i> , <i>Briza</i> , <i>Bromus</i>). TL: Zhiguli preserv (Samara Distr.). Steppe biotopes. L: unknown.	
96.	<i>Eupista samborensis</i> ANIKIN (in press)						VI-VII in 1 G				
97.	<i>Klimeschja oriolella</i> (ZELLER, 1849) (= <i>mongettella</i> CHRETIEN, 1900)						- VI-VII in 1 G			Forest-steppe biotopes. L: <i>Coronilla</i> .	
98.	<i>Klimeschja vulnerariae</i> (ZELLER, 1839)						- VII in 1 G			Local in forest-steppes. L: <i>Anthyllis</i> .	
99.	<i>Bourgogneja pennella</i> ([DENIS & SCHIFFERMÜLLER], 1775) (= <i>lomosmella</i> BRAHM, 1791)						+ VI-VII in 1 G			Was cited by E as <i>Struthiopennella</i> . Forest biotopes. L: <i>Hieracium</i> *, <i>Stachys</i> , <i>Anchusa</i> , <i>Symphtym</i> , <i>Pulmonaria</i> , <i>Lithospermum</i> , <i>Myosotis</i> , <i>Cynoglossum</i> .	
100.	<i>Perygra murinipennella</i> (DUPONCHEL, 1844)						- V-VI in 1 G			Was cited by E as <i>Otidipennella</i> , but it was determined only by external characters; this determination has to be confirmed. Forest-steppe biotopes. L: <i>Luzula</i> .	
101.	<i>Perygra coracipennella</i> (HÜBNER, 1796) (= <i>nigricella</i> STEPHENS, 1835; = <i>varii</i> PATZAK, 1969)						+ V in 1 G			Steppe with bush. L: <i>Cerasus</i> .	

1	2	3	4	5	6	7	8	9	10	11	12
102.	<i>Perygra numeniella</i> (BALDIZZONE, 1988)								?	o VII in 1 G	TL: Uralsk. From Ulianovsk known after 1 ♂ which may be identified previously as this species.
103.	<i>Perygra alticolella</i> (ZELLER, 1849) (= <i>caespititiella</i> auct.)								- VI-VII in 1 G	Forest and forest-steppe biotopes. L: <i>Juncus</i> .	
104.	<i>Perygra caespititiella</i> (ZELLER, 1839)								- VI in 1 G	Forest biotopes with fens. L: <i>Juncus articulatus</i> .	
105.	<i>Perygra glaucicolella</i> (WOOD, 1892)								- mV-VII in 1 G	Forest-steppe biotopes; wet meadows, meadow-steppe. L: on the carpels of <i>Juncus</i> .	
106.	<i>Perygra adjunctella</i> (HODGKINSON, 1882) (= <i>paludicola</i> STANTON, 1886)								- VI-VII in 1 G	Forest-steppe biotopes. L: <i>Juncus</i> .	
107	<i>Perygra taeniipennella</i> (HERRICH-SCHÄFFER, 1855)								- mVI-VII in 1 G	Forest-steppe biotopes; wet meadows. L: <i>Juncus obtusiflorus</i> .	
108.	<i>Tritemachia teredo</i> FALKOVITSH, 1994								- V-VI in 1 G	Desert and desert-steppe biotopes. L: <i>Anabasis aphylla</i> *	
109.	<i>Ecebalia therinella</i> (TENGSTROM, 1848)								- VI, VII-VIII in 1-2 G	Steppe biotopes and anthropogenic landscapes. L: <i>Carlina</i> , <i>Carduus</i> , <i>Cirsium</i> .	
110.	<i>Ecebalia protella</i> (ZELLER, 1871) (fig. 5)								- eV-VI, IX in 1-2 G	Forest-steppe biotopes. L: <i>Carduus</i> , <i>Cirsium</i> , <i>Corispermum hyssopifolium</i> *	
111.	<i>Ecebalia bagarella</i> (FALKOVITSH, 1977) (fig. 5)								- VIII in 1 G	Along the left side of the Volga. Before known from Mongolia only.	
112.	<i>Ecebalia vestianella</i> (LINNAEUS, 1758) (= <i>laripennella</i> ZETTER-STEDT, 1839)	+ + + + + +							+ VI-VIII in 1 G	The species occurs everywhere in the region in the places of the host plants distribution, in cultural landscapes. L: <i>Atriplex</i> *, <i>Chenopodium</i> *	
113.	<i>Ecebalia gaviaepennella</i> (TOLL, 1952)								- VIII-IX in 1 G	Common in desert-steppe, semi-desert biotopes and anthropogenic landscapes.	
114.	<i>Ecebalia magyarica</i> (BALDIZZONE, 1983)								- VIII in 1 G	Rare in desert and semi-desert biotopes. L: <i>Kochia</i> , <i>Camphorosma</i> .	
115.	<i>Ecebalia superlonga</i> (FALKOVITSH, 1989) (fig. 5)								- VIII in 1 G	Dry steppe biotopes. L: <i>Kochia prostrata</i> *, <i>Salsola australis</i> *	
116.	<i>Ecebalia lunensis</i> FALKOVITSH, 1975								- VIII in 1 G	Not common in desert-steppe biotopes. L: unknown.	
117.	<i>Ecebalia eichleri</i> (PATZAK, 1977)								IX in 1 G	Desert-steppe biotopes. L: <i>Kochia prostrata</i> f. <i>viridens</i> *	
118.	<i>Ecebalia saxicolella</i> (DUPONCHEL, 1843) (= <i>bennardi</i> KANERVA, 1941, = <i>flaginella</i> auct.)								+ VII-VIII in 1 G	The moth occurs in the places of the host plant's distribution. L: <i>Atriplex</i> *, <i>Chenopodium</i> .	

1	2	3	4	5	6	7	8	9	10	11	12
119. <i>Ecebalia motacillella</i> (ZELLER, 1849)		+ + + +		VII-VIII in 1 G	The moth occurs in the places of the host plants distribution. L: <i>Atriplex*</i> , <i>Chenopodium*</i>						
120. <i>Ecebalia sternipennella</i> (ZETTERSTEDT, 1839) (= <i>flavaginella</i> LIENIG & ZELLER, 1846)		+ VII-VIII in 1 G		Steppe and desert biotopes, in wildlands and as well as in the anthropogenic landscapes. L: <i>Atriplex*</i> , <i>Chenopodium*</i>							
121. <i>Ecebalia nomgona</i> (FAKOVITSH, 1975)		VIII in 1 G		Rare in dry steppes. The first note from Russia.							
122. <i>Ecebalia versurella</i> (ZELLER, 1849) (= <i>amoranthivora</i> Oku, 1965)	+ + + + + + +	V-VIII in 1-2 G		The moth occurs in the places of the host plants distribution. L: <i>Atriplex*</i> , <i>Amaranthus</i> , <i>Chenopodium*</i>							
123. <i>Ecebalia pseudolinosyris</i> KASY, 1979		- VI-VII in 1 G		Forest-steppe biotopes. L: <i>Galatella villosa*</i>							
124. <i>Ecebalia thyrrhaenica</i> (AMSEL, 1951)		- VIII in 1 G		Steppe biotopes. L: unknown.							
125. <i>Ecebalia adspersella</i> (BENANDER, 1939)		- VII-VIII in 1 G		Forest and forest-steppe biotopes, anthropogenic landscapes. L: <i>Chenopodium*</i> , <i>Atriplex*</i>							
126. <i>Ecebalia aestuariella</i> (BRADLEY, 1975)		- IX in 1 G		Desert-steppe biotopes. L: <i>Suaeda maritima*</i>							
127. <i>Ecebalia lassella</i> (STAUDINGER, 1859)		- V in 1 G		Rare and local in stepped biotopes.							
128. <i>Ecebalia tornata</i> (FAKOVITSH, 1989)		- IX in 1 G		Desert-steppe biotopes. L: <i>Kochia prostrata f. viridens*</i>							
129. <i>Ecebalia anabaseos</i> (FAKOVITSH, 1975)		- V-VI in 1 G		Desert-steppe biotopes. L: <i>Anabasis aphylla*</i>							
130. <i>Ecebalia charadiella</i> (BALDIZZONE, 1988)	+ V-VI in 1 G			Local and rare in forest-steppe, steppe and dry steppe biotopes.							
131. <i>Ecebalia ottalicella</i> (ZELLER, 1871) (= <i>unistriella</i> CARADJA, 1920)	- VI-mVII in 1 G			TL: Sarepta. Forest-steppe and steppe biotopes.							
132. <i>Ecebalia quadrifariella</i> (STAUDINGER, 1880)	+ VI-VII in 1 G			TL: Sarepta.							
133. <i>Ecebalia kargani</i> (FAKOVITSH, 1989)	- VI-VII in 1 G			Semidesert biotopes. L: <i>Kochia prostrata f. viridens*</i> , <i>Salsola dendroides</i> .							
134. <i>Ecebalia virgaureae</i> (STAINTON, 1857)	- VII-VIII in 1 G			Forest-steppe biotopes. L: <i>Solidago virgaurea*</i> , <i>Aster amellus</i> .							
135. <i>Ecebalia halophylella</i> (ZIMMERMANN, 1926)	+ VIII-IX in 1 G			Forest-steppe biotopes.							
136. <i>Ecebalia halocnemi</i> (FAKOVITSH, 1994)	- V-VI in 1 G			Desert-steppe biotopes. L: <i>Halocnemum strobilaceum*</i> .							
137. <i>Ecebalia halostachydis</i> (FAKOVITSH, 1994)	- V-VI in 1 G			Desert-steppe biotopes. L: <i>Halostachys caspica*</i>							

1	2	3	4	5	6	7	8	9	10	11	12
138.	<i>Casignetella absinthii</i> (HEINEMANN & WOCKE, 1877)			+ +		VII in 1 G				Steppe and forest-steppe biotopes. L: <i>Artemisia campestris</i> *	
139.	<i>Casignetella ancistron</i> (FALKOVITSH, 1976) (fig. 6)					V, VII-VIII in 2 G				Steppe biotopes. L: unknown.	
140.	<i>Casignetella occatella</i> (STAUDINGER, 1880) (fig. 6)					V, VIII in 1-2 G				TL. Sarepta. Steppe and desert- steppe biotopes. L: unknown.	
141.	<i>Casignetella amarchana</i> (FALKOVITSH, 1975)					VII-VIII in 1 G				Steppe and desert-steppe biotopes. L: unknown.	
142.	<i>Casignetella argentula</i> (STEPHENS, 1834) (= <i>cothurn-</i> <i>nella</i> DUPONCHEL, 1843)			+ VI-VII in 1 G						The moth occurs in meadow-steppe and dry meadows. L: <i>Achillea</i> <i>millefolium</i> *	
143.	<i>Casignetella tanaceti</i> (MÜHLIG, 1865)					VI-VII in 1 G				Forest and forest-steppe biotopes. L: <i>Tanacetum vulgare</i> *	
144.	<i>Casignetella erratella</i> (TOLL & AMSEL, 1967)			- V in 1 G						Not rare, but local in semi-desert biotopes. The first note for Russia.	
145.	<i>Casignetella pilion</i> FALKOVITSH, 1992					V in 1 G				Rare in semi-desert and dry steppe biotopes and salt-marshs.	
146.	<i>Casignetella directella</i> (ZELLER, 1849)					VII-VIII in 1 G				Forest and forest-steppe biotopes. L: <i>Artemisia campestris</i> *	
147.	<i>Casignetella artemisiella</i> (SCOTT, 1861)	+ + + + + +		+ V-VII in 1-2 G						Steppe and forest-steppe biotopes. L: <i>Artemisia campestris</i> *	
148.	<i>Casignetella gnaphalii</i> (ZELLER, 1839)			- VII-VIII in 1 G						Forest-steppe biotopes. L: <i>Gnapha-</i> <i>lium, Helichrysum arenarium</i> .	
149.	<i>Casignetella granulatella</i> (ZELLER, 1849) (= <i>arte-</i> <i>misiae</i> MÜHLIG, 1864)			+ VI-VII in 1 G						Steppe and forest-steppe biotopes. L: <i>Artemisia campestris, A. scop-</i> <i>aria</i> .	
150.	<i>Casignetella peisoniella</i> (KASY, 1965)			- VIII in 1 G						Dry steppe biotopes.	
151.	<i>Casignetella ramosella</i> (ZELLER, 1849)			eVI-VII in 1 G						Forest-steppe biotopes. L: <i>Solidago</i> <i>virgaurea, Aster amellus</i> .	
152.	<i>Casignetella trochilella</i> (DUPONCHEL, 1843)			- VI-VII in 1 G						Steppe and forest-steppe biotopes. L: <i>Inula, Achillea, Tanacetum, Arte-</i> <i>misia, Carduus</i> .	
153.	<i>Casignetella inulae</i> (HEI- NEMANN & WOCKE, [1876])			- VII in 1 G						Forest-steppe biotopes. L: <i>Inula</i> .	
154.	<i>Casignetella gardesanella</i> - (TOLL, 1953) (= <i>machinella</i> BRADLEY, 1971)			- VI-VII in 1 G						Forest-steppe biotopes. L: <i>Artemisia</i> <i>maritima, Centaurea jacea</i> .	
155.	<i>Casignetella peribenonderi</i> (TOLL, 1943) (= <i>benonderi</i> TOLL, 1942)			- VI-VII in 1 G						Forest-steppe biotopes. L: <i>Carduus,</i> <i>Cirsium arvense</i> .	
156.	<i>Casignetella kyffhusana</i> (PETRY, 1898) (Fig. 5)			- VI-VII, VIII-IX in 1-2 G						Forest-steppe and steppe biotopes. L: <i>Gypsophila fastigiata</i> *	

1	2	3	4	5	6	7	8	9	10	11	12
157.	<i>Casignetella spec.</i> (pr. <i>kudrosella</i> BALDIZZONE & OKU, 1988)			+		-	VII in 1 G				Stoned steppe biotopes in Zhituli preserv. Known after 1 ♂ and more material is necessary for exact identification. L: unknown.
158.	<i>Casignetella spec.</i> (pr. <i>ammophora</i> FALKOVITSH, 1989)						VII-VIII in 1 G				Steppe biotopes. L: unknown.
159.	<i>Casignetella dentatella</i> (TOLL & AMSEL, 1967)					V-VI in 1 G					Semidesert biotopes with sand. L: <i>Acantophyllum</i> *
160.	<i>Casignetella dianthi</i> (HERRICH-SCHÄFFER, 1855)					-	VI-VII in 1 G				Forest and forest-steppe biotopes. L: <i>Dianthus</i> .
161.	<i>Casignetella silenella</i> (HERRICH-SCHÄFFER, 1855)					-	V-VI in 1 G				Forest biotopes. L: <i>Viscaria, Silene</i> *
162.	<i>Casignetella ciconiella</i> (HERRICH-SCHÄFFER, 1855)					VI-VII in 1 G					Steppe biotopes. L: <i>Silene</i> *
163.	<i>Casignetella graminecolella</i> (HEINEMANN, 1876)					-	VI in 1 G				Forest-steppes. L: <i>Silene</i> .
164.	<i>Casignetella pseudo-ciconiella</i> (TOLL, 1952)					-	VIII in 1 G				Steppe and forest-steppe biotopes. L: <i>Silene</i> *
165.	<i>Casignetella albilineella</i> (TOLL, 1960)					-	VI-VII in 1 G				Forest-steppe biotopes. L: unknown.
166.	<i>Casignetella niveistrigella</i> (HEINEMANN & WOCKE, 1877)					-	mVI-VII in 1 G				Steppe and forest-steppe biotopes. L: <i>Gypsophila fastigiata</i> *, <i>Gypsophila juzepczukii</i> *
167.	<i>Casignetella solitariella</i> (ZELLER, 1849)					-	VI-VII in 1 G				Forest biotopes. L: <i>Stellaria, Silene, Cerastium</i> .
168.	<i>Casignetella deviella</i> (ZELLER, 1847)					-	IX in 1 G				Steppe biotopes. L: <i>Halostachys caspica</i> *
169.	<i>Casignetella tremula</i> FALKOVITSH, 1989					V-VI in 1 G					Steppe biotopes. The larva was marked on <i>Arthropodium iliense</i> (FALKOVITSH, 1989) (this plant isn't native for the Lower Volga region). In deserts of Kazakhstan and Middle Asia it has 2-3 generations.
170.	<i>Casignetella lebedella</i> FALKOVITSH, 1982					-	V, VIII-IX in 2 G				Steppe and desert-steppe biotopes. L: <i>Atriplex</i> *
171.	<i>Casignetella hungariae</i> (GOZMANY, 1955)					-	VIII in 1 G				Dry steppe biotopes. L: <i>Kochia prostrata</i> *
172.	<i>Casignetella stepposa</i> (FALKOVITSH, 1975)					-	VII in 1 G				Steppe and desert-steppe biotopes. L: unknown.
173.	<i>Casignetella artemisicolella</i> (BRUAND, [1855])					-	VII-VIII in 1 G				Forest-steppe and steppe biotopes. L: <i>Artemisia vulgaris</i> *
174.	<i>Casignetella galatellae</i> (HERING, 1942)					VI in 1 G					Local and rare in forest-steppe biotopes. L: <i>Crinitaria</i> .

1	2	3	4	5	6	7	8	9	10	11	12
175.	<i>Casignetella galbulipennella</i> (ZELLER, 1838) (= <i>otitae</i> ZELLER, 1838)						+	+	VII-VIII in 1 G	Forest-steppe and steppe biotopes. L: <i>Silene</i> .	
176.	<i>Casignetella scabrida</i> (TOLL, 1959)							V	in 1 G	Steppe biotopes. L: <i>Herniaria glabra</i> .	
177.	<i>Casignetella striatipennella</i> (TENGSTRÖM, [1848])							VI-VIII in 1 G	Forest-steppe biotopes. L: <i>Cerastium, Stellaria</i> .		
178.	<i>Casignetella tringella</i> (BALDIZZONE, 1988)						o	eV, VIII in 1 G	TL: Uralsk. Steppe and semi-desert biotopes. L: <i>Kochia prostrata</i> *		
179.	<i>Casignetella remisella</i> (BALDIZZONE, 1983) (fig. 6)						-	IX in 1 G	Steppe biotopes. L: <i>Kochia prostrata</i> *		
180.	<i>Ionescumia clypeiferella</i> (HOFMANN, 1871)	+	+	+	+	+	+	+	VII-VIII in 1 G	Steppe and cultured landscapes. L: <i>Chenopodium, hibernates</i> .	
181.	<i>Ionescumia dilabens</i> FALKOVITSH, 1982							VII in 1 G	Steppe and semidesert biotopes. L: <i>Salsola orientalis</i> .		
182.	<i>Ionescumia acerosa</i> FALKOVITSH, 1989						-	VIII in 1 G	Local in semidesert biotopes.		
183.	<i>Carpochena trientella</i> (CHRISTOPH, 1872) (fig. 6)						-	VIII in 1 G	TL: Sarepta. Desert biotopes, salt-marshs and chalk-steppes. L: <i>Corispermum</i> *		
184.	<i>Carpochena aequalella</i> (CHRISTOPH, 1872) (= <i>heratella</i> TOLL & AMSEL, 1967) (fig. 6)						o	V-VI, VIII in 1-2 G	TL: Sarepta, Ural. Steppe and half-desert biotopes.		
185.	<i>Carpochena unipunctella</i> (ZELLER, 1849)						-	VI-VII in 1 G	Forest-steppe biotopes. L: <i>Chenopodium, Atriplex</i> .		
186.	<i>Carpochena binotapennella</i> (DUPONCHEL, 1843)						+	VII-VIII in 1 G	Steppe biotope. L: <i>Chenopodium*, Atriplex*</i>		
187.	<i>Carpochena ceratoidis</i> FALKOVITSH, 1979						-	VIII in 1 G	Chalk hills with stepped biotopes. L: <i>Ceratooides papposa</i> *		
188.	<i>Carpochena squalorella</i> (ZELLER, 1849) (= <i>delibutella</i> CHRISTOPH, 1872)						+	+	VIII in 1 G	Steppe and semidesert biotopes. L: <i>Chenopodium*, Atriplex</i> . TL for <i>delibutella</i> CHR. is Sarepta.	
189.	<i>Carpochena armeniae</i> (BALDIZZONE & PATZAK, 1991)						-	VIII in 1 G	Steppe and semidesert biotopes. L: unknown.		
190.	<i>Carpochena salicorniae</i> (HEINEMANN & WOCKE, 1877)						-	VIII in 1 G	Desert biotopes; salt-marshs. L: carpels of <i>Salicornia europaea</i> , boring the summit of a carpel.		
191.	<i>Carpochena asperginella</i> (CHRISTOPH, 1872) (= <i>nigrosquamella</i> FILIPJEV, 1925) (fig. 6)						-	VIII in 1 G	TL: Sarepta. Desert biotopes. L: carpels of <i>Corispermum</i> .		
192.	<i>Klinzigidia phlomidella</i> (CHRISTOPH, 1862) (fig. 6)						-	VIII in 1 G	TL: Sarepta. Steppe biotopes. L: <i>Phlomis</i> .		

1	2	3	4	5	6	7	8	9	10	11	12
193. <i>Klinzigedia phlomidis</i> (STAINTON, 1867)		+			mVII- mVIII in 1 G					Noted from Sarepta by REBEL (1901). Steppe biotopes. L: <i>Phlomis</i> <i>pungens</i> .	
194. <i>Klinzigedia wockeella</i> (ZELLER, 1849) (= <i>italia</i> TOLL, 1960; = <i>fumigatella</i> CAPUSE, 1974)					VI-VII in 1 G					Forest biotopes. L: <i>Stachys</i> .	
195. <i>Goniodoma auroguttella</i> ZELLER, 1849					eVI-VIII in 1 G					Steppe and semidesert biotopes. L: <i>Atriplex</i> .	
Gelechiidae											
196. <i>Pycnostola bohemica</i> (NICKERL, 1864)		-	V-VI, VIII in 2 G							Rare in forest-steppe biotopes.	
197. <i>Metzneria ehikeella</i> GOZMANY, 1954			V-VIII in 1-2G							Rare in forest-steppe biotopes. L: <i>Centaurea scabiosa</i> , <i>C. calcitrapa</i> .	
198. <i>Metzneria metzneriella</i> (STAINTON, 1951)			V-VII in 1 G							Meadows-steppe biotopes. L: <i>Centaurea jacea</i> , <i>C. nigriceps</i> .	
199. <i>Metzneria santolinella</i> AMSEL, 1951		-	VI-VII in 1 G							Rare in forest-steppe biotopes. L: <i>Anthemis tinctoria</i> .	
200. <i>Metzneria lappella</i> (LINNAEUS, 1758)			V-VIII in 2 G							The moth occurs in the places of the host plants distribution. L: <i>Arctium lappa</i> .	
201. <i>Metzneria littoralis</i> DOUGLAS, 1850		-	VI-VII in 1 G							Forest-steppe biotopes. L: <i>Plantago</i> <i>coronopus</i> .	
202. <i>Metzneria artificella</i> (HERRICH-SCHÄFFER, 1861) (= <i>litigiosella</i> MILLIERE, 1879; = <i>pannonicella</i> REBEL, 1915)										TL: Sarepta.	
203. <i>Metzneria aestivella</i> ZELLER, 1939		-	V-VIII in 1 G							Dry meadows. L: <i>Centaurea</i> <i>diffusa</i> , <i>C. solstitialis</i> , <i>C. scabiosa</i> .	
204. <i>Metzneria neuropterella</i> ZELLER, 1839			VI-IX in 2 G							From Volgograd Distr. known from old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Forest-steppe biotopes. L: <i>Cirsium esculentum</i> , <i>Dipsacus</i> , <i>Campanula</i> .	
205. <i>Metzneria aprilella</i> (HERRICH-SCHÄFFER, 1854)		+	V-VIII in 2 G							From Volgograd region known from the old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Steppe biotopes. L: <i>Centaurea diffusa</i> , <i>C. scabiosa</i> .	
206. <i>Metzneria paucipunctella</i> ZELLER, 1839		-	V-VII in 1 G							Dry steppe biotopes. L: <i>Centaurea</i> , <i>Anthemis tinctoria</i> .	
207. <i>Metzneria subflavella</i> ENGLERT, 1974			VI-VIII in 2 G							Forest-steppe and steppe biotopes.	

1	2	3	4	5	6	7	8	9	10	11	12
208.	<i>Isophrictis striatella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				+ + +		VII in 1 G			Forest and forest-steppe biotopes.	
209.	<i>Isophrictis anthemidella</i> (WOCKE, 1871)						VI-VII in 1 G			Forest-steppe biotopes. L: <i>Anthemis subtinctoria</i> .	
210.	<i>Eulamprotes wilkella</i> (LINNAEUS, 1758) (= <i>pictella</i> ZELLER, 1839)						VI-VIII in 1-2 G			Steppe and desert-steppe biotopes. L: <i>Cerastium holosteoides</i> .	
211.	<i>Eulamprotes superbella</i> (ZELLER, 1839)						V-VI in 1 G			Noted from Sarepta by REBEL (1901). Local in dry biotopes. L: <i>Thymus serpyllum</i> .	
212.	<i>Eulamprotes atrella</i> ([DENIS & SCHIFFERMÜLLER], 1775)						VI-VIII in 1 G			Rare in forest-steppe biotopes. L: <i>Hypericum perforatum</i> , <i>H. maculatum</i> .	
213.	<i>Eulamprotes unicolorella</i> (DUPONCHEL, 1843)				-		V-VII in 1 G			Not common in forest-steppe. L: un-known.	
214.	<i>Argolamprotes micella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				-		VI-VIII			Forest biotopes. L: <i>Rubus idaeus</i> , <i>R. caeius</i> .	
215.	<i>Monochroa tenebrella</i> (HÜBNER, [1817])				+	-	VI-VII			Rare in forest-steppe biotopes. L: <i>Rumex acetosella</i> .	
216.	<i>Monochroa palustrella</i> (DOUGLAS, 1850)				-	mV-VIII in 1-2 G				Near the water biotopes. L: <i>Rumex aquaticus</i> , <i>R. crispus</i> .	
217.	<i>Monochroa lucidella</i> (STEPHENS, 1834)				+		VI-VIII in 1 G			Near the water biotopes. L: <i>Scirpus</i> , <i>Juncus</i> .	
218.	<i>Monochroa nomadella</i> (ZELLER, 1868)				-	VII in 1 G				Very rare in forest-steppe biotopes. L: unknown.	
219.	<i>Monochroa hornigi</i> (STAUDINGER, 1883)				-		VI-VII in 1 G			Steppe biotopes. L: <i>Polygonum aviculare</i> .	
220.	<i>Monochroa saltenella</i> (BENANDER, 1928)				-	VII in 1 G				Forest-steppe biotopes.	
221.	<i>Monochroa elongella</i> (HEINEMANN, 1870)					mV-VI in 1 G				Near the water biotopes. L: <i>Potentilla</i> .	
222.	<i>Monochroa sepicolella</i> (HERRICH-SCHÄFFER, 1854)					VI-VII in 1 G				Rare in steppe biotopes. L: <i>Rumex</i> .	
223.	<i>Ptocheuusa inopella</i> (ZELLER, 1847)				-	eV-VII in 1 G				Rare in dry steppe biotopes. L: <i>Helichrysum arenarium</i> .	
224.	<i>Ptocheuusa sublutella</i> CHRISTOPH, 1872					?				TL: Sarepta. Biology unknown.	
225.	<i>Chrysoesthia drurella</i> (FABRICIUS, 1775) (= <i>hermannella</i> auct.)					V-VIII in 2 G				The moth occurs in the places of the host plants distribution. L: <i>Atriplex*</i> , <i>Chenopodium*</i>	
226.	<i>Chrysoesthia sexguttella</i> (THUNBERG, 1794)					V-VIII in 2 G				The moth occurs in the places with host-plants. L: <i>Chenopodium alba*</i> , <i>Atriplex patula*</i>	
227.	<i>Psamathocrita osseella</i> (STAINTON, 1861)					V-VII in 1-2 G				Very rare in steppe biotopes. L: <i>Origanum</i> .	

1	2	3	4	5	6	7	8	9	10	11	12
228.	<i>Caulastrocecis furfurella</i> (STAUDINGER, 1870) (= <i>cryptexena</i> GOZMANY, 1952) (fig. 7)		o	+		V-VII in 1 G		TL: Sarepta. Forest-steppe biotopes. L: <i>Galatella linosyris</i> .			
229.	<i>Aristotelia subdecurtella</i> (STAINTON, 1859) (= <i>pro-</i> <i>haskaella</i> REBEL, 1911)					VII-VIII in 1 G		Rare and local, meadow-steppe biotopes. L: <i>Lythrum</i> .			
230.	<i>Aristotelia subericinella</i> (HERRICH-SCHÄFFER, 1853)					VI in 1 G		Chalk hills with forest-steppe biotopes. L: <i>Lytrum salicaria</i> .			
231.	<i>Aristotelia decurtella</i> (HÜBNER, [1813])					- VII-VIII in 1 G		Forest-steppe and steppe biotopes. L: <i>Eryngium campestre</i> , <i>Sanguisorba officinalis</i> , <i>Rosa</i> .			
232.	<i>Aristotelia interstratella</i> (CHRISTOPH, 1872) (= <i>sali-</i> <i>natrix</i> MEYRICK, 1926)					+ VII in 1 G		TL: Sarepta. TL for <i>salinatrix</i> MEYRICK: Ural.			
234.	<i>Aristotelia panchaliella</i> (STAUDINGER, 1870)					- V-VIII in 1-2 G		Steppe and desert-steppe biotopes.			
235.	<i>Aristotelia cervinella</i> (EVERSMANN, 1844) (fig. 7)					o VII in 1 G		TL: Ural (Guberli). Very rare in meadow-steppe biotopes.			
236.	<i>Aristotelia calastomella</i> (CHRISTOPH, 1872)							TL: Sarepta. We know one specimen of moth only collected by CHRISTOPH in Sarepta, which is kept in BMNH. Biology unknown.			
237.	<i>Aristotelia mirabilis</i> (CHRISTOPH, 1888)							TL: Sarepta.			
238.	<i>Evippe pseudolella</i> (CHRISTOPH, 1888)					ø ?		TL: Sarepta and Guberli (Ural).			
239.	<i>Xystophora pulveratella</i> (HERRICH-SCHÄFFER, 1854)					+ IV-V in 2 G		Forest-steppe biotopes. L: <i>Medicago sativa</i> , <i>M. minima</i> , <i>Lotus corniculatus</i> , <i>Coronilla varia</i> , <i>Trifolium</i> , <i>Onobrychis</i> .			
240.	<i>Xystophora orthagonella</i> (STAUDINGER, 1870)							TL: Sarepta. Steppe biotopes. Biol- ogy unknown.			
241.	<i>Chilopselaphus fallax</i> MANN, 1867					+ VI-VIII in 1 G		From Volgograd Distr. known from the old collection material (CHRIS- TOPH) from Sarepta, which is kept in ZMHUB. Steppe biotopes. L: <i>Stipa</i> .			
242.	<i>Megacraspedus separatellus</i> (FISCHER VON RÖSLERSTAMM, 1844)					- V-mVI in 1 G		Steppe biotopes.			
243.	<i>Megacraspedus argyroneuellus</i> (STAU- DINGER, 1870) (fig. 7)					+ eVI-VII in 1 G		TL: Sarepta. Steppe biotopes.			
244.	<i>Megacraspedus attritellus</i> - (STAUDINGER, 1870) (fig. 7)					+ V in 1 G		TL: Sarepta. Steppe and desert- steppe biotopes.			
245.	<i>Megacraspedus binotellus</i> - (FISCHER VON RÖSLERSTAMM, 1844)					IV-VI in 1 G		Steppe biotopes. L: <i>Poaceae</i> .			

1	2	3	4	5	6	7	8	9	10	11	12
246. <i>Megacraspedus dolosellus</i> ZELLER, 1839				+ V-VI in 1 G		Rare in steppe biotopes. L: <i>Cynodon dactylon</i> , <i>Poa bulbosa</i> , cultural Poaceae.					
247 <i>Stenolechia gemmella</i> (LINNAEUS, 1758)				- VII-IX in 1-2 G		Oak forests. L: <i>Quercus</i> .					
248. <i>Parachronistis albiceps</i> (ZELLER, 1839)				V-VII in 1 G		Parks and green zone of landscapes, forests. L: <i>Corylus avellana</i> , <i>Ulmus</i> , <i>Malus</i> , <i>Persica</i> .					
249. <i>Recurvaria nanella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				VI-VIII in 2 G		The moth occurs in the places with host-plants. L: woody Rosaceae.					
250. <i>Recurvaria leucatella</i> (CLERCK, 1759)				- VI-VIII in 1-2 G		Data as for <i>Recurvaria nanella</i> .					
251. <i>Exoteleia dodecella</i> (LINNAEUS, 1758)				- VI-VII in 1 G		Forest biotopes. L: <i>Pinus sylvestris</i> .					
252. <i>Athrips mouffetella</i> (LINNAEUS, 1758)				- VI-VII in 1 G		Forest biotopes. L: <i>Lonicera xylosteum</i> .					
253. <i>Athrips nigricostella</i> (DUPONCHEL, 1842)				- V-VIII in 1-2 G		Forest-steppe biotopes. L: <i>Medicago sativa</i> , <i>Caragana frutex</i> .					
254. <i>Athrips spiraeae</i> (STAUDINGER, 1871) (fig. 8)				- VI-VII in 1 G		TL: Sarepta. Forest-steppe biotopes. L: <i>Spiraea media</i> .					
255. <i>Xenolechia scriptella</i> (HÜBNER, 1796)				- V-VIII in 1-2 G		Forest biotopes. L: <i>Acer</i> .					
256. <i>Agonochaetia intermedia</i> SATTLER, 1968				- VII in 1 G		TL: Sarepta. Steppe biotopes.					
257. <i>Teleiodes vulgella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				+ V-VII in 1 G		Forest-steppe biotopes. L: Rosaceae.					
258. <i>Teleiodes notatella</i> (HÜBNER, [1813])				- V-VI in 1 G		Forest-steppe biotopes. L: <i>Salix caprea</i> , <i>S. rosmarinifolia</i> .					
259. <i>Teleiodes anguinella</i> (HERRICH-SCHÄFFER, 1861) (fig. 8)				+ V in 1 G		TL: Sarepta. Steppe biotopes. Biology unknown.					
260. <i>Teleiodes aenigma</i> SATTLER, 1983 (<i>wagae</i> NOWICKI sensu PISKUNOV, 1973)				- IV-V in 1 G		Forest-steppe biotopes, dry meadows.					
261. <i>Teleiodes semicostella</i> (STAUDINGER, 1870)						TL: Sarepta.					
262. <i>Pseudotelphusa scalella</i> (SCOPOLI, 1763)				+ V-VII, VIII in 2 G		Oak forests. L: Musci.					
263. <i>Pseudotelphusa paripunctella</i> (THUNBERG, 1794) (= <i>triparella</i> ZELLER, 1839)				+ V-VII in 1 G		Forest biotopes. L: <i>Quercus</i> , <i>Betula</i> , <i>Salix caprea</i> .					
264. <i>Pseudotelphusa proximella</i> (HÜBNER, 1796)				IV-VII in 2 G		Rare in forest biotopes. L: <i>Betula</i> , <i>Alnus</i> .					
265. <i>Pseudotelphusa alburnella</i> (ZELLER, 1839)				VI-VIII in 1-2 G		Very common in forests. L: <i>Betula</i> .					

	1	2	3	4	5	6	7	8	9	10	11	12
266.	<i>Pseudotelphusa fugitivella</i> (ZELLER, 1839)			+	+	+	+			VI-VIII in 2 G	From Volgograd Distr. known from the old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Very common in forests. L: <i>Ulmus</i> , <i>Acer</i> , <i>Fraxinus</i> , <i>Tilia</i> .	
267	<i>Pseudotelphusa fugacella</i> (ZELLER, 1839)									VI-VII in 1 G	Rare in forests. L: <i>Ulmus</i> .	
268.	<i>Pseudotelphusa luculella</i> (HÜBNER, 1813)				-	V-VII in 1-2 G				Forest biotopes. L: <i>Quercus</i> , <i>Betula</i> , <i>Salix</i> .		
269.	<i>Teleiopsis diffinis</i> (HAWORTH, 1828)				-	V-VII, VIII-IX in 2 G				Dry forest-steppe biotopes. L: <i>Rumex acetosella</i> .		
270.	<i>Alténia perspersella</i> (WOCKE, 1862)				-	mV-VII in 1 G				Near the water and inundated forests.		
271.	<i>Bryotropha basaltinella</i> (ZELLER, 1839)				+	VI-VIII in 1 G				L: <i>Musci</i> .		
272.	<i>Bryotropha umbrosella</i> ZELLER, 1839 (= <i>oppositella</i> THUNBERG, 1794)				-	VI-VIII in 1 G				Rare in forest biotopes. L: <i>Musci</i> .		
273.	<i>Bryotropha affinis</i> (HAWORTH, 1828)				-	VI-VIII in 1 G				Dry steppe biotopes.		
274.	<i>Bryotropha similis</i> (STAINTON, 1854) (= <i>obscure-cinerea</i> NOLCKEN, 1871)					VI-VIII in 1 G				Forest and forest-steppe biotopes. L: lichens.		
275.	<i>Bryotropha mundella</i> DOUGLAS, 1850				-	VI-VII in 1 G				Forest and forest-steppe biotopes.		
276.	<i>Bryotropha senectella</i> ZELLER, 1839				-	VI-VIII in 1 G				Forest and forest-steppe biotopes. L: lichens.		
277.	<i>Bryotropha rossica</i> ANIKIN - & PISKUNOV, 1996 (fig. 8)				-	VI in 1 G				Known from TL: Chardym is at the Volga river.		
278.	<i>Bryotropha terrella</i> [(DENIS - + & SCHIFFERMÜLLER], 1775)				+	VI-IX in 2 G				Forest-steppe biotopes. L: <i>Poaceae</i> .		
279.	<i>Bryotropha desertella</i> (DOUGLAS, 1850)				-	V-VIII in 1-2 G				Forest-steppe biotopes. L: <i>Musci</i> .		
280.	<i>Chionodes viduella</i> (FABRICIUS, 1794)				-	VI-VII in 1 G				Forest-steppe biotopes. L: <i>Betula</i> , <i>Rubus</i> .		
281.	<i>Chionodes lugubrella</i> (FABRICIUS, 1794)				+	VI-VII in 1 G				Forest biotopes. L: <i>Betula</i> .		
282.	<i>Chionodes tragicella</i> (HEYDEN, 1865)					VI in 1 G				Near the forest park in Kamyshin. L: <i>Larix</i> .		
283.	<i>Chionodes distinctella</i> (ZELLER, 1839)				-	VI-IX in 2 G				The moth occurs in the places of the host-plants. L: <i>Artemisia campestris</i> , <i>Rumex</i> , <i>Thymus</i> .		
284.	<i>Chionodes luctuella</i> (HÜBNER, 1793)				-	VI-VII in 1 G				Steppe biotopes.		
285.	<i>Chionodes fumatella</i> DOUGLAS, 1850					VII-IX in 1 G				Rare in forest-steppe. L: <i>Musci</i> .		

1	2	3	4	5	6	7	8	9	10	11	12
246.	<i>Megacraspedus dolosellus</i> ZELLER, 1839						+ -	V-VI in 1 G	Rare in steppe biotopes. L: <i>Cynodon dactylon</i> , <i>Poa bulbosa</i> , cultural Poaceae.		
247	<i>Stenolechia gemmella</i> (LINNAEUS, 1758)						-	VII-IX in 1-2 G	Oak forests. L: <i>Quercus</i> .		
248.	<i>Parachronistis albiceps</i> (ZELLER, 1839)						-	V-VII in 1 G	Parks and green zone of landscapes, forests. L: <i>Corylus avellana</i> , <i>Ulmus</i> , <i>Malus</i> , <i>Persica</i> .		
249.	<i>Recurvaria nanella</i> ([DENIS & SCHIFFERMÜLLER], 1775)						-	VI-VIII in 2 G	The moth occurs in the places with host-plants. L: woody Rosaceae.		
250.	<i>Recurvaria leucatella</i> (CLERCK, 1759)						-	VI-VIII in 1-2 G	Data as for <i>Recurvaria nanella</i> .		
251.	<i>Exoteleia dodecella</i> (LINNAEUS, 1758)						-	VI-VII in 1 G	Forest biotopes. L: <i>Pinus sylvestris</i> .		
252.	<i>Athrips mouffetella</i> (LINNAEUS, 1758)							VI-VII in 1 G	Forest biotopes. L: <i>Lonicera xylosteum</i> .		
253.	<i>Athrips nigricostella</i> (DUPONCHEL, 1842)						-	V-VIII in 1-2 G	Forest-steppe biotopes. L: <i>Medicago sativa</i> , <i>Caragana frutex</i> .		
254.	<i>Athrips spiraeaee</i> (STAUDINGER, 1871) (fig. 8)						-	VI-VII in 1 G	TL: Sarepta. Forest-steppe biotopes. L: <i>Spiraea media</i> .		
255.	<i>Xenolechia scriptella</i> (HÜBNER, 1796)						-	V-VIII in 1-2 G	Forest biotopes. L: <i>Acer</i> .		
256.	<i>Agonochaeta intermedia</i> SATTLER, 1968						-	VII in 1 G	TL: Sarepta. Steppe biotopes.		
257.	<i>Teleiodes vulgella</i> ([DENIS & SCHIFFERMÜLLER], 1775)						+	V-VII in 1 G	Forest-steppe biotopes. L: Rosaceae.		
258.	<i>Teleiodes notatella</i> (HÜBNER, [1813])						-	VI-VI in 1 G	Forest-steppe biotopes. L: <i>Salix caprea</i> , <i>S. rosmarinifolia</i> .		
259.	<i>Teleiodes anguinella</i> (HERRICH-SCHÄFFER, 1861) (fig. 8)						+ V	in 1 G	TL: Sarepta. Steppe biotopes. Biology unknown.		
260.	<i>Teleiodes aenigma</i> SATTLER, 1983 (<i>wagae</i> NOWICKI sensu PISKUNOV, 1973)						IV-V in 1 G	Forest-steppe biotopes, dry meadows.			
261.	<i>Teleiodes semicastella</i> (STAUDINGER, 1870)								TL: Sarepta.		
262.	<i>Pseudotelphusa scalella</i> (SCOPOLI, 1763)						+ V-VII, VIII	in 2 G	Oak forests. L: Musci.		
263.	<i>Pseudotelphusa paripunctella</i> (THUNBERG, 1794) (= <i>triparella</i> ZELLER, 1839)						+ V-VII	in 1 G	Forest biotopes. L: <i>Quercus</i> , <i>Betula</i> , <i>Salix caprea</i> .		
264.	<i>Pseudotelphusa proximella</i> (HÜBNER, 1796)						IV-VII	in 2 G	Rare in forest biotopes. L: <i>Betula</i> , <i>Alnus</i> .		
265.	<i>Pseudotelphusa alburnella</i> (ZELLER, 1839)						VI-VIII	in 1-2 G	Very common in forests. L: <i>Betula</i> .		

	1	2	3	4	5	6	7	8	9	10	11	12
266.	<i>Pseudotelphusa fugitivella</i> (ZELLER, 1839)			+	+	+	+	-	VI-VIII in 2 G		From Volgograd Distr. known from the old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Very common in forests. L: <i>Ulmus</i> , <i>Acer</i> , <i>Fraxinus</i> , <i>Tilia</i> .	
267	<i>Pseudotelphusa fugacella</i> (ZELLER, 1839)								VI-VII in 1 G		Rare in forests. L: <i>Ulmus</i> .	
268.	<i>Pseudotelphusa luculella</i> (HÜBNER, 1813)							V-VII in 1-2 G			Forest biotopes. L: <i>Quercus</i> , <i>Betula</i> , <i>Salix</i> .	
269.	<i>Teleiopsis diffinis</i> (HAWORTH, 1828)							- V-VII, VIII-IX in 2 G			Dry forest-steppe biotopes. L: <i>Rumex acetosella</i> .	
270.	<i>Altenia perspersella</i> (WOCKE, 1862)							- mV-VII in 1 G			Near the water and inundated forests.	
271.	<i>Bryotropha basaltinella</i> (ZELLER, 1839)							+ VI-VIII in 1 G			L: <i>Musci</i> .	
272.	<i>Bryotropha umbrosella</i> ZELLER, 1839 (= <i>oppositella</i> THUNBERG, 1794)							VI-VIII in 1 G			Rare in forest biotopes. L: <i>Musci</i> .	
273.	<i>Bryotropha affinis</i> (HAWORTH, 1828)							- VI-VIII in 1 G			Dry steppe biotopes.	
274.	<i>Bryotropha similis</i> (STAINTON, 1854) (= <i>obscure-cinerea</i> NOLCKEN, 1871)							VI-VIII in 1 G			Forest and forest-steppe biotopes. L: lichens.	
275.	<i>Bryotropha mundella</i> DOUGLAS, 1850							- VI-VII in 1 G			Forest and forest-steppe biotopes.	
276.	<i>Bryotropha senectella</i> (ZELLER, 1839)							- VI-VIII in 1 G			Forest and forest-steppe biotopes. L: lichens.	
277	<i>Bryotropha rossica</i> ANIKIN - & PISKUNOV, 1996 (fig. 8)							- VI in 1 G			Known from TL: Chardym is at the Volga river.	
278.	<i>Bryotropha terrella</i> ([DENIS - & SCHIFFERMÜLLER], 1775)							+ VI-IX in 2 G			Forest-steppe biotopes. L: Poaceae.	
279.	<i>Bryotropha desertella</i> (DOUGLAS, 1850)							- V-VIII in 1-2 G			Forest-steppe biotopes. L: <i>Musci</i> .	
280.	<i>Chionodes viduella</i> (FABRICIUS, 1794)							- VI-VII in 1 G			Forest-steppe biotopes. L: <i>Betula</i> , <i>Rubus</i> .	
281.	<i>Chionodes lugubrella</i> (FABRICIUS, 1794)							+ VI-VII in 1 G			Forest biotopes. L: <i>Betula</i> .	
282.	<i>Chionodes tragicella</i> (HEYDEN, 1865)							- VI in 1 G			Near the forest park in Kamyshin. L: <i>Larix</i> .	
283.	<i>Chionodes distinctella</i> (ZELLER, 1839)							- VI-IX in 2 G			The moth occurs in the places of the host-plants. L: <i>Artemisia campestris</i> , <i>Rumex</i> , <i>Thymus</i> .	
284.	<i>Chionodes luctuella</i> (HÜBNER, 1793)							VI-VII in 1 G			Steppe biotopes.	
285.	<i>Chionodes fumatella</i> DOUGLAS, 1850							VII-IX in 1 G			Rare in forest-steppe. L: <i>Musci</i> .	

1	2	3	4	5	6	7	8	9	10	11	12
286.	<i>Chionodes continua</i> (ZELLER, 1839)					+	+		VII-VIII in 1 G	Rare in the northern forests of <i>Pinus</i> . L: <i>Cladonia rangiferina</i> .	
287	<i>Chionodes violacea</i> (TENGSTRÖM, 1847) (fig. 8)								VIII in 1 G	Rare in forest-steppe. Noted from Russia for the first time.	
288.	<i>Lita longicornis</i> (CURTIS, 1827)								V-VII in 1 G	Forest-steppe biotopes. L: <i>Musci</i> .	
289.	<i>Lita solutella</i> (ZELLER, 1839)								+ V-VII in 1-2 G	Steppe biotopes. L: <i>Genista</i> .	
290.	<i>Mirificarma mulinella</i> (ZELLER, 1839)								VI-VIII in 1 G	Forest-steppe biotopes.	
291.	<i>Mirificarma lenticiginosella</i> + (ZELLER, 1839)								VII in 1 G	Steppe biotopes. L: <i>Genista tinctoria</i> .	
292.	<i>Mirificarma maculatella</i> (HÜBNER, 1796)								VI-VII in 1 G	Forest-steppe biotopes. L: <i>Coronilla varia</i> .	
293.	<i>Mirificarma cytisella</i> TREITSCHKE, 1833								+ eV-VII in 1 G	Forest-steppe biotopes. L: <i>Cytisus</i> .	
294.	<i>Aroga velocella</i> (ZELLER, 1839) (fig. 9)								- IV-VI, VII-VIII in 2 G	Steppe biotopes. From Samara Distr. it's known from Sergievsk (KRULIKOVSKY, 1915). L: <i>Rumex acetosella</i> .	
295.	<i>Aroga flavicornella</i> (ZELLER, 1839)								+ IV-VI in 1 G	Forest-steppe biotopes. L: <i>Prunus spinosa</i> , <i>Prunus</i> , <i>Cerasus</i> .	
296.	<i>Aroga pascuicola</i> STAUDINGER, 1871								- VII in 1 G	Forest-steppe biotopes.	
297	<i>Neofriseria peliella</i> (TREITSCHKE, 1835)								VI-VIII in 1 G	In dry clearings of forest-steppe biotopes. L: <i>Rumex</i> .	
298.	<i>Neofriseria singula</i> (STAUDINGER, 1876) (= <i>suppiliella</i> WALSINGHAM, 1896)								- V-VI in 1 G	Forest biotopes.	
299.	<i>Lutilabria volgensis</i> ANIKIN - & PISKUNOV, 1996 (fig. 9)								- mV in 1 G	TL: Khvalynsk. The stepped chalk hills.	
300.	<i>Filatima incomptella</i> (HERRICH-SCHÄFFER, 1853)								- V in 1 G	Steppe biotopes.	
301.	<i>Filatima tephriditella</i> (DUPONCHEL, 1843)								IV-VI in 1 G	Forest-steppe biotopes.	
302.	<i>Filatima djakovica</i> ANIKIN & PISKUNOV, 1996 (fig. 9)								V in 1 G	Steppe biotopes. TL: Eruslan River, Dyakowskii reserv.	
303.	<i>Filatima zagulajevi</i> ANIKIN & PISKUNOV, 1996 (fig. 9)								- V in 1 G	Steppe biotopes. Only known from TL: Eruslan River, Dyakowskii reserv.	
304.	<i>Filatima pallipalpella</i> (SNELLEN, 1844)								VI-VII in 1 G	Very rare. Biology unknown.	
305.	<i>Epilechia magnetella</i> (STAUDINGER, 1870)								VI in 1 G	Very rare. Known from the vicinity of Uljanovsk.	

1	2	3	4	5	6	7	8	9	10	11	12
306.	<i>Gelechia rhombella</i> ([DENIS & SCHIFFERMÜLLER], 1775)		+ + +						VI-IX in 2 G	Very common in the gardens. L: woody Rosaceae.	
307	<i>Gelechia rhombelliformis</i> (STAUDINGER, 1870)								+ VI-IX in 2 G	Very common in forest-steppe and steppe biotopes. L: <i>Populus nigra</i> , <i>P. pyramidalis</i> .	
308.	<i>Gelechia scotinella</i> (HERRICH-SCHÄFFER, 1854)								+ IV-VIII in 2 G	The moth occurs in the places of the host-plants. L: ligneous Rosaceae.	
309.	<i>Gelechia nigra</i> (HAWORTH, 1828)								VII-VIII in 1 G	Noted from Sarepta by REBEL (1901). Forest biotopes. L: <i>Populus</i> .	
310.	<i>Gelechia muscosella</i> (ZELLER, 1839)								- VI-VII in 1 G	Not common in forest biotopes. L: <i>Salix caprea</i> .	
311.	<i>Gelechia sestertiella</i> (HERRICH-SCHÄFFER, 1854)								VI-VII in 1 G	Forest biotopes. L: <i>Acer</i> .	
312.	<i>Gelechia turpella</i> ([DENIS & + SCHIFFERMÜLLER], 1775) (= <i>pinguinella</i> TREITSCHKE, 1832)								+ VI-IX in 2 G	Was noted by E as <i>Pinguinella</i> . For- est and forest-steppe biotopes. L: <i>Populus nigra</i> .	
313.	<i>Gelechia sororculella</i> (HÜBNER, [1817])								- VI-VII in 1 G	Forest and forest-steppe biotopes. L: <i>Quercus, Sorbus</i> .	
314.	<i>Gelechia basipunctella</i> (HERRICH-SCHÄFFER, 1854)								- VI-VII in 1 G	Rare in forest biotopes near the river. L: <i>Salix</i> .	
315.	<i>Gelechia jakovlevi</i> KRULIKOVSKY, 1905								- mVI-IX in 2 G	Anthropogenic landscapes. L: <i>Malus, Ribes nigrum</i> .	
316.	<i>Gelechia pistaciae</i> FILIPJEV, 1934								+ VII in 1 G	Noted by MARTYNOWA (1952) from Ural district.	
317.	<i>Gladiovalva badidorsella</i> (REBEL, 1935)								- VIII in 1 G	Steppe biotopes.	
318.	<i>Ornativalva plutelliformis</i> (STAUDINGER, 1859)								- VI-VIII in 2 G	Steppe and desert-steppe biotopes. L: <i>Tamarix</i> .	
319.	<i>Ornativalva heluanensis</i> (DEBSKI, 1913) (= <i>frankeni-vorella</i> CHRETEN, 1916)								- V-VII in 1 G	Steppe and desert-steppe biotopes. L: <i>Frankenia, Tamarix</i> .	
320.	<i>Ornativalva sieversi</i> (STAUDINGER, 1870) (fig. 9)								VI-VII in 1 G	TL: Sarepta. Steppe and desert- steppe biotopes. L: <i>Tamarix</i> .	
321.	<i>Ornativalva ornatella</i> SATTLER, 1967								- V-VII in 1 G	Steppe and desert-steppe biotopes.	
322.	<i>Phthorimaea operculella</i> (ZELLER, 1873)								VI-IX in 1-2 G	1 ♀ was found in Saratov, there it can be introduced. L: Solanaceae.	
323.	<i>Phthorimaea praticolella</i> (CHRISTOPH, 1872)								?	Known after 1 ♂ from TL: Sarepta (ZISP).	
324.	<i>Gnorimoschema</i> <i>epithymellum</i> (STAUDINGER, 1859)								- VII-VIII in 1 G	Forest-steppe biotopes. L: Solanaceae.	

1	2	3	4	5	6	7	8	9	10	11	12
325.	<i>Gnorimoschema robustellum</i> (STAUDINGER, 1870) (= <i>syrphetopa</i> MEYRICK, 1926)		o	+		-	VI in 1 G			TL: Sarepta. Steppe biotopes.	
326.	<i>Gnorimoschema strelicella</i> (HERRICH-SCHÄFFER, 1853) (Fig. 10)					-	VI in 1 G			Chalk hills in the forest-steppe zone. L: <i>Thymus</i> .	
327	<i>Scrobipalpula psilella</i> (HERRICH-SCHÄFFER, 1854)					-	V, VI-VIII in 2 G	Dry steppe biotopes. L: <i>Artemisia</i> , <i>Gnaphalium</i> , <i>Achillea</i> , <i>Tanacetum</i> , <i>Helichrysum arenarium</i> , <i>Aster amellus</i> .			
328.	<i>Scrobipalpula erichi</i> PVOVOLNY, 1966					-	IV, VII-IX in 2 G	Agricultural landscapes. L: Solanaceae.			
329.	<i>Scrobipalpa</i> (<i>Euscrobipalpa</i>) <i>acuminatella</i> (SIRCOM, 1850)					-	V-VII, IX-X in 2 G	Forest-steppe biotopes. L: <i>Cardus</i> , <i>Centaurea</i> , <i>Cirsium</i> , <i>Serratula</i> , <i>Tanacetum</i> .			
330.	<i>Scrobipalpa</i> (<i>E.</i>) <i>ustulatella</i> (STAUDINGER, 1870) (Fig. 10)					-	VII in 1 G	TL: Sarepta. Dry steppe biotopes.			
331.	<i>Scrobipalpa</i> (<i>E.</i>) <i>pauperella</i> (HEINEMANN, 1870)					VII-VIII in 1 G		Desert-steppe biotops.			
332.	<i>Scrobipalpa</i> (<i>E.</i>) <i>nitentella</i> (FUCHS, 1902) (= <i>seminella</i> PIERCE & METCALFE, 1935)					-	VIII-IX in 1 G	Desert-steppe biotops.			
333.	<i>Scrobipalpa</i> (<i>E.</i>) <i>klimeschi</i> PVOVOLNY, 1967 (Fig. 10)					-	VIII in 1 G	Steppe biotops.			
334.	<i>Scrobipalpa</i> (<i>E.</i>) <i>rjabovi</i> PISKUNOV, 1990 (Fig. 10)					-	VIII in 1 G	Steppe biotops.			
335.	<i>Scrobipalpa</i> (<i>E.</i>) <i>artemisiella</i> (TREITSCHKE, 1833)					-	VI-VIII in 1-2G	Forest-steppe biotopes. L: <i>Thymus</i> , <i>Mentha</i> , <i>Centaurea</i> , <i>Artemisia</i> .			
336.	<i>Scrobipalpa</i> (<i>E.</i>) <i>murinella</i> (HERRICH-SCHÄFFER, 1854)					-	V-VI in 1 G	Dry steppe biotopes. L: <i>Antennaria dioica</i> .			
337.	<i>Scrobipalpa</i> (<i>E.</i>) <i>ocellatella</i> (BOYD, 1858)					-	V-VIII in 1-2G	L: <i>Beta</i> .			
338.	<i>Scrobipalpa</i> (<i>E.</i>) <i>obsoletella</i> (FISCHER VON RÖSLERSTAMM, 1841) (= <i>miscatella</i> CLARKE, 1932)					V-IX in 2 G		Common everywhere; open places, anthropogenic landscapes. L: <i>Atriplex</i> , <i>Chenopodium</i> .			
339.	<i>Scrobipalpa</i> (<i>E.</i>) <i>atriplicella</i> (FISCHER VON RÖSLERSTAMM, 1841)					IV-VIII in 2 G		Noted from Sarepta by REBEL (1901). Very common everywhere. L: <i>Atriplex patula</i> , <i>Chenopodium</i> .			
340.	<i>Scrobipalpa</i> (<i>E.</i>) <i>chrysanthemella</i> (HOFMANN, 1867)					-	V-VIII in 2 G	Forest-steppe biotopes. L: <i>Leucanthemum vulgare</i> , <i>Artemisia absinthium</i> .			

1	2	3	4	5	6	7	8	9	10	11	12
341.	<i>Scrobipalpa (E.) salinella</i> (ZELLER, 1847)				+		V-VIII in 1-2 G			Rare in steppe biotopes.	
342.	<i>Scrobipalpa (E.) hyos-</i> <i>cyamivora</i> (GERASIMOV, 1940)				-	VI-VII in 1 G			Forest-steppe biotopes. L: <i>Hyoscyamus</i> .		
343.	<i>Scrobipalpa (E.) phag-</i> <i>nalella</i> (CONSTANT, 1895)					V-VI in 1 G			Desert and desert-steppe biotopes.		
344.	<i>Scrobipalpa (E.) bryo-</i> <i>philoides</i> POVOLNY, 1966								TL: Sarepta. Steppe biotopes. Biology unknown. Some specimens collected by CHRISTOPH in Sarepta and Krasnovodsk (Turkmenistan) are kept in BMNH.		
345.	<i>Scrobipalpa (E.) gregori</i> POVOLNY, 1967 (fig. 10)				-	VI in 1 G			Desert and desert-steppe. L: <i>Holocnemnum strobiloceum</i> *. Noted from Russia for the first time.		
346.	<i>Scrobipalpa (E.) brandti</i> POVOLNY, 1972 (fig. 10)					IX in 1 G			Desert and desert-steppe biotopes. Noted from Russia for the first time.		
347.	<i>Scrobipalpa (E.) hyos-</i> <i>cyamella</i> (STAINTON, 1869)				?				TL: Sarepta. Steppe biotopes. Biology unknown.		
348.	<i>Scrobipalpa (E.) costella</i> (HUMPHREYS & WESTWOOD, 1845)				+ ?				Steppe biotopes. Biology unknown. One specimen collected by CHRISTOPH in "Ural" is kept in BMNH.		
349.	<i>Scrobipalpa (E.) pseud-</i> <i>obsoletella</i> (POVOLNY & GREGOR, 1955) (fig. 11)				-	VII in 1 G			Semi-desert steppe.		
350.	<i>Scrobipalpa (E.) disjectella</i> - + (STAUDINGER, 1859) (fig. 11)					V in 1 G			Very local in desert biotopes.		
351.	<i>Opacopsis inustella</i> (ZELLER, 1839) (= <i>gredosensis</i> REBEL, 1904)				-	VII-VIII in 1 G			Dry-steppe biotopes. L: <i>Festuca ovina</i> .		
352.	<i>Opacopsis deserticolella</i> (STAUDINGER, 1870) (fig. 11)				-	V-IX in 2 G			TL: Sarepta. Steppe and desert-steppe biotopes.		
353.	<i>Opacopsis insulella</i> (HEINEMANN, 1870)				+	IV-VII in 1-2 G			From Volgograd Distr. known from old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Forest-steppe and steppe biotopes.		
354.	<i>Cosmardia moritzella</i> (TREITSCHKE, 1835)				-	VI-IX in 2 G			Rare in virgin steppes. L: <i>Melandrium album</i> .		
355.	<i>Caryocolum fischerella</i> (TREITSCHKE, 1833)				-	VI-IX in 2 G			Forest biotopes.		
356.	<i>Caryocolum visciarella</i> (STAINTON, 1855) (= <i>abbi-</i> <i>faciella</i> HEINEMANN, 1870)				-	VII in 1 G			Steppe biotopes. L: <i>Lychnis</i> , <i>Silene</i> , <i>Cucubalus</i> . For Russia noted for the first time.		
357.	<i>Caryocolum vicinella</i> (DOUGLAS, 1851)					VII-VIII in 1 G			Forest biotopes. L: <i>Silene nutans</i> , <i>S. cucubalis</i> , <i>Tunica saxifraga</i> , <i>Cerastium</i> .		

1	2	3	4	5	6	7	8	9	10	11	12
358.	<i>Caryocolum blandella</i> (DOUGLAS, 1852) (= <i>maculea</i> auct.)		+ +		VII-IX in 2 G						From Volgograd Distr. known from old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Common in forest biotopes. L: <i>Stellaria</i> .
359.	<i>Caryocolum kroesmaniella</i> (HERRICH-SCHÄFFER, 1854) (= <i>huebneri</i> auct., nec HAWORTH, 1828)				VI-IX in 2 G						Forest biotopes. L: <i>Stellaria holostea</i> .
360.	<i>Caryocolum schleichi</i> (CHRISTOPH, 1872) (= <i>syriacum</i> POVOLNY, 1977) (fig. 11)				V-VI, VIII-IX in 2 G						TL: Sarepta. Steppe and desert-steppe biotopes. L: unknown.
361.	<i>Caryocolum pullatella</i> (TENGSTRÖM, 1848)				VIII in 1 G						Only known from Sarepta.
362.	<i>Nothris verbascella</i> ([DENIS & SCHIFFERMÜLLER], 1775)				- V-VI, VII-IX in 2 G						Steppe biotopes. L: <i>Verbascum</i> .
363.	<i>Thiotricha subocellea</i> (STEPHENS, 1834)				- VI-VII in 1 G						Very rare in forest-steppe biotopes. L: <i>Origanum vulgare</i> .
364.	<i>Sophronia semicostella</i> (HÜBNER, [(1813)])				- VI-VII in 1 G						From Volgograd Distr. known from old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB. Steppe biotopes. L: <i>Dianthus</i> , <i>Anthoxanthum odoratum</i> .
365.	<i>Sophronia chilonella</i> (TREITSCHKE, 1833)				+ VI-VII in 1 G						Noted from Sarepta by REBEL (1901). Forest-steppe and steppe biotopes. L: <i>Artemisia campestris</i> .
366.	<i>Sophronia sicariella</i> (ZELLER, 1839)				- VI-VII in 1 G						Noted from Sarepta by REBEL (1901). Forest-steppe and steppe biotopes. L: <i>Artemisia campestris</i> , <i>Tanacetum</i> .
367	<i>Sophronia consanguinella</i> - (HERRICH-SCHÄFFER, 1855)				+ VI-VII in 1 G						Forest-steppe biotopes. L: <i>Artemisia campestris</i> , <i>Potentilla</i> .
368.	<i>Aproaerema anthyllidella</i> (HÜBNER, (1813))				- V-VIII in 2 G						Very common everywhere. L: <i>Fabaceae</i> .
369.	<i>Stomopteryx detersella</i> (ZELLER, 1847)				VI-VIII in 1-2G						Forest-steppe and steppe biotopes. L: <i>Eryngium</i> .
370.	<i>Stomopteryx taeniolella</i> (ZELLER, 1839)				V-VIII in 1 G						Forest-steppe and steppe biotopes. L: <i>Lotus corniculatus</i> , <i>Medicago</i> , <i>Trifolium</i> .
371.	<i>Syncopacma sangiella</i> (STAINTON, 1863)				VI-VIII in 1 G						Forest-steppe biotopes. L: <i>Lotus corniculatus</i> *
372.	<i>Syncopacma coronilletella</i> (TREITSCHKE, 1833)				V-VI in 1 G						Forest-steppe biotopes. L: <i>Coronilla varia</i> .
373.	<i>Syncopacma cinctella</i> (CLERCK, 1759) (= <i>vorticella</i> SCOPOLI, 1763)				+ VI-VII in 1 G						Common everywhere. L: <i>Lotus corniculatus</i> , <i>Genista tinctoria</i> .

1	2	3	4	5	6	7	8	9	10	11	12
374.	<i>Syncopacma vinella</i> (BANKES, 1898)				+			~ VI in 1 G		Steppe biotopes.	
375.	<i>Syncopacma larseniella</i> GOZMANY, 1957 (fig. 11)							- VI-VII in 1 G		Very rare in forest-steppe biotopes.	
376.	<i>Lixodessa ochrofasciella</i> (TOLL, 1936)							- V, VII-VIII in 1-2 G		Chalk hills and chalk steppes.	
377.	<i>Iwaruna biguttella</i> (DUPONCHEL, 1843)							- VI-VIII in 1 G		Steppe biotopes. L: herbaceous Fabaceae.	
378.	<i>Acanthophila alacella</i> (ZELLER, 1839)							- VI-VIII in 1 G		Not common in deciduous forests. L: lichens.	
379.	<i>Acompsia cinerella</i> (CLERCK, 1759)							+ V-VIII in 1-2 G		Common in forest biotopes. L: Lichens.	
380.	<i>Anacampsis populella</i> (CLERCK, 1759)	+	+	+	+	+	+	+	+ VI-X in 2 G	Forest and forest-steppe biotopes. L: <i>Populus tremula</i> , <i>P. nigra</i> , <i>Salix caprea</i> , <i>S. repens</i> , <i>Acer campestre</i> .	
381.	<i>Anacampsis disquei</i> (MEESS, 1907) (= <i>quercellea</i> CHRETIEN, 1907)							+ VI-VIII in 1 G		In old oak forests. L: <i>Quercus robur</i> *	
382.	<i>Anacampsis blattariella</i> (HÜBNER, 1796) (= <i>betulinella</i> VARI, 1941)							VII-IX in 1 G		Forest-steppe biotopes. L: <i>Betula pendula</i> *	
383.	<i>Anacampsis temerella</i> (LIENIG & ZELLER, 1846)							- VI in 1 G		Forest-steppe biotopes.	
384.	<i>Anacampsis subsequella</i> (HÜBNER, 1796)							+ VII-VIII in 1 G		Forest-steppe biotopes. L: ligneous Rosaceae, <i>Salix</i> .	
385.	<i>Anacampsis cincticulella</i> (HERRICH-SCHÄFFER, 1854)							- ?		From Volgograd region known from old collection material (CHRISTOPH) from Sarepta, which is kept in ZMHUB.	
386.	<i>Sitotroga cerealella</i> (OLIVIER, 1789)							- VI-VIII in 2-3 G		Common in anthropogenic landscape. L: Poaceae, Fabaceae.	
387.	<i>Metanarsia modesta</i> STAUDINGER, 1871 (fig. 12)							V-VIII in 1 G		TL: Sarepta. Steppe biotopes.	
388.	<i>Metanarsia incertella</i> HERRICH-SCHÄFFER, 1861 (fig. 12)							+ V-VI in 1 G		TL: Sarepta. Steppe and desert-steppe biotopes.	
389.	<i>Platyedra subcinerea</i> (HAWORTH, 1828) (= <i>vilella</i> ZELLER, 1847)							V-X in 2 G		Steppe biotopes. L: <i>Althaea officinalis</i> .	
390.	<i>Pexicopia malvella</i> (HÜBNER, 1805)	+	+	+	+	+	+	+ VII-VIII in 1-2 G		Forest-steppe biotopes. L: <i>Lavatera</i> , <i>Malva</i> , <i>Alcea rosea</i> , <i>Althaea</i> .	
391.	<i>Anarsia lineatella</i> (ZELLER, 1839)							+ V-VIII in 2 G		Forest and forest-steppe biotopes. L.: ligneous Rosaceae.	

	1	2	3	4	5	6	7	8	9	10	11		12
392.	<i>Anarsia spartiella</i> (SCHRANK, 1802)			+ + +							VI-VIII in 1 G	From Volgograd region known from old collection material (CHRISTOPH) Sarepta, which is kept in ZMHUB. Rare in forest-steppe biotopes. L: <i>Genista tinctoria</i> .	
393.	<i>Anarsia eleagnella</i> KUZNEZOV, 1957			-	eVI-VII in 1 G							Forest-steppe and steppe biotopes. L: <i>Eleagnus</i> .	
394.	<i>Psoricoptera gibbosella</i> (ZELLER, 1839)			+	VII-VIII in 1 G							Forest and forest-steppe biotopes. L: <i>Quercus robur</i> *, <i>Salix</i> , <i>Crataegus</i> , <i>Malus</i> .	
395.	<i>Hypatima rhomboidella</i> (LINNAEUS, 1758)				VII-IX in 1 G							Was listed by E as <i>Conscriptella</i> HBN. Forest biotopes. L: <i>Betula</i> , <i>Corylus</i> , <i>Alnus</i> , <i>Populus tremula</i> .	
396.	<i>Holcophora statices</i> STAUDINGER, 1871			-	V in 1 G							TL: Sarepta. Steppe biotopes.	
397.	<i>Mesophleps silacella</i> (HÜBNER, 1796)			-	V-VII in 1 G							Dry steppe biotopes. L: <i>Helianthemum</i> .	
398.	<i>Telephila schmidtiellus</i> (HEYDEN, 1848)			-	eV-VII in 1 G							Forest and forest-steppe biotopes. L: <i>Origanum vulgare</i> .	
399.	<i>Dichomeris ustalella</i> (FABRICIUS, 1794)			-	V-VII in 1 G							Rare in forest biotopes. L: <i>Betula</i> , <i>Quercus</i> , <i>Acer</i> .	
400.	<i>Dichomeris fasciella</i> (HÜBNER, 1796)			-	V-VIII in 1-2 G							Forest-steppe biotopes. L: Rosaceae.	
401.	<i>Dichomeris limosella</i> (SCHLÄGER, 1849)			+	VI-IX in 2 G							Forest-steppe and steppe biotopes. L: <i>Trifolium pratense</i> , <i>Medicago sativa</i> .	
402.	<i>Dichomeris barbella</i> (HÜBNER, 1796)			+	V in 1 G							L: <i>Prunus</i> .	
403.	<i>Uliaria rasilella</i> HERRICH-SCHÄFFER, 1854			-	VII-VIII in 1 G							Forest-steppe biotopes. L: <i>Centaurea</i> .	
404.	<i>Brachmia dimidiella</i> ([DENIS & SCHIFFERMÜLLER], 1775)			+	VI-VII in 1 G							Forest-steppe biotopes. L: <i>Seceli libanotis</i> .	
405.	<i>Brachmia blandella</i> (FABRICIUS, 1798)			-	VI-VII in 1 G							Dry meadows in deciduous forests.	
406.	<i>Brachmia triannulella</i> (HERRICH-SCHÄFFER, 1854)			-	IV-IX in 2 G							Meadows-steppe and steppe biotopes. L: <i>Convolvulus arvensis</i> , <i>Calystegia sepium</i> .	
407.	<i>Helcystogramma lutatella</i> - (HERRICH-SCHÄFFER, 1854)				VII-IX in 1 G							Steppe biotopes. L: <i>Elytrigia repens</i> .	
408.	<i>Helcystogramma albinervis</i> (GERASIMOV, 1929)			-	VI, VII-IX in 2 G							Not common in virgin salt steppe biotopes.	
409.	<i>Helcystogramma rufescens</i> (HAWORTH, 1828)				VI-VIII in 1 G							Steppe biotopes. L: Poaceae.	
410.	<i>Deroxena venosulella</i> (MÖSCHLER, 1862)			+	IV-VI in 1 G							Steppe biotopes. L: unknown.	

1	2	3	4	5	6	7	8	9	10	11	12
---	---	---	---	---	---	---	---	---	----	----	----

Symmocidae

411. *Oegoconia quadripuncta* (HAWORTH, 1828) (fig. 12) mVI-VIII in 1-2 G Steppe biotopes.
412. *Oegoconia bacescui* POPESCU-GORJ & CAPUSE, 1965 - VI in 1 G Steppe and forest-steppe biotopes.
413. *Eremicamima cedestiella* (ZELLER, 1868) + eV-VIII in 1 G Steppe biotopes.
414. *Symmoca signatella* HERRICH-SCHÄFFER, 1854 - ? Noted from Sarepta by REBEL (1901).

Holcopogonidae

415. *Holcopogon bubulcellus* STAUDINGER, 1859 (fig. 12) - eV-VIII in 1-2 G Steppe biotopes. Noted from Sarepta by REBEL (1901) as *helveolellus* STGR.

Taxonomic appendix

During the work with STAUDINGER's collection in ZMHUB we found the type specimen "*Coleophora unicolorella* TOLL" (+; Sarepta, coll. CHRISTOPH; GP ♀ 136) and "*Coleophora sareptella* TOLL" (+; Sarepta, coll. CHRISTOPH; GP ♀ 132) the descriptions of which were not published by TOLL. After our predetermination of the female of "*C. unicolorella*" (fig. 13) we establish it a synonym of *Casignetella gnaphalii* (ZELLER, 1839) and the female of "*C. sareptella*" is a synonym of *Carpochena salicorniae* (HEINEMANN & WOCKE, 1877).

References

- ANIKIN, V. V. (1995): On the fauna of casebearers (Lepidoptera, Coleophoridae) from Uljanovsk district. – Priroda Uljanovskoj oblasti. Uljanovsk. **6** (2): 50–57 (in russian).
- ANIKIN, V. V. (1998a): The casebearers of the Volgo-Ural inter-river region (Lepidoptera, Coleophoridae). – Entomofauna, Zeit. Entom. **19** (2): 33–44.
- ANIKIN, V. V. (1998b): On the fauna of gelechiid and casebearers (Lepidoptera; Gelechiidae, Coleophoridae) from Uljanovsk district. – Priroda Uljanovskoj oblasti. Uljanovsk. **9**.
- ANIKIN, V. V. & M. I. FALKOVITSH (1997): On the casebearer fauna of the Lower Volga region (Lepidoptera: Coleophoridae). – Zoosyst. Rossica **5**: 303–308.
- ANIKIN, V. V. & V. I. PISKUNOV (1995): On the fauna of gelechiid moths (Lepidoptera, Gelechiidae) from the Lower Volga region. Contribution to the knowledge of the Gelechiidae from the European part of Russia. – Actias **2** (1–2): 3–12.
- ANIKIN, V. V. & V. I. PISKUNOV (1996): New species of gelechiid moths (Lepidoptera, Gelechiidae) from the Lower Volga region (Russia). – Zoosystematica Rossica **3** (1): 171–175.
- ANIKIN, V. V. & S. A. SACHKOV (1998): On the casebearer fauna of Zhiguli preserve (Lepidoptera: Coleophoridae). – Zapovidna sprava in Ukraine **4** (1): 89–90 (in russian).




Fig. 1: Map of the Volga-Ural region: ● *Orthographis uralensis* (TOLL); ▲ *O. ptarmicia* (WLSGHM.); ■ - *O. virgatella* (ZLL.); ♦ *Polystrophia calligoni* (FLKv.).




Fig. 2: Map of the Volga-Ural region: ● - *A. klimeschiella* (TOLI); ▲ - *A. eurasiatica* (BLDZ.); ◆ - *A. lonchodes* FLKv., *A. physophorae* FLKv., *A. dissecta* FLKv., *A. nigridorsella* (AMSEL).




Fig. 3: Map of the Volga-Ural region: ● - *Chnoocera botigurella* (H.-S.); ▲ - *Oedicaula serinipennella* (CHR.).




Fig. 4: Map of the Volga-Ural region: ● - *Multicoloria changaica* (REZN.); ▲ - *M. singreni* (FLKV.); ◆ - *M. incostans* REZN.




Fig. 5: Map of the Volga-Ural region: ● - *Ecebalia pratella* (ZLL.); ■ - *E. bagorella* (FLKV.);
◆ - *E. superlonga* (FLKV.); ▲ - *E. kyffhusana* (PETRY).




Fig. 6: Map of the Volga-Ural region: ● - *Casignetella ancistron* (FLKV.); ▲ - *C. occatella* (STGR.); ◆ - *C. remisella* (BLDZ.); ■ - *Carpochena trientella* (CHR.), *C. aequalella* (CHR.), *C. asperginella* (CHR.), *Klinzigidia phlomidella* (CHR.).




Fig. 7: Map of the Volga-Ural region: ● - *Caulastrocecis furfurella* (Stgr.); ▲ - *Aristotelia cervinella* (Ev.); ◆ - *Megacraspedus argyroneurellus* Stgr.; ■ - *M. attritellus* Stgr.




Fig. 8: Map of the Volga-Ural region: ● - *Athrips spiraeae* (Stgr.); ▲ - *Teleiodes anguinella* (H.-S.); ◆ - *Bryotropha rossica* Anik. & Pisk.; ■ - *Chionodes violacea* (Tngstr.).




Fig. 9: Map of the Volga-Ural region: ● - *Aroga velocella* (ZLL.); ▲ - *Lutilabria volgensis* ANIK. & PISK.; ◆ - *Filatima djakovica* ANIK. & PISK., *F. zagulajevi* ANIK. & PISK.; ■ - *Ornativalva sieversi* (STGR.).




Fig. 10: Map of the Volga-Ural region: ● - *Gelechia strelicella* (H.-S.); ▲ - *Scrobipalpa (Euscrobi-palpa) ustulatella* (STGR.); ▲ - *S. (E.) klimeschi* Pov., *S. (E.) rjabovi* Pisk.; ■ - *S. (E.) gregori* Pov., *S. (E.) brandti* Pov.




Fig. 11: Map of the Volga-Ural region: ● *Scrobipalpa (Eusrobipalpa) pseudobsoletella* (Pov. & Gr.), S. (E.) *disjectella* (STGR.); ▲ *Opacopsis deserticolella* (STGR.); ♦ *Caryocolum schlechi* (CHR.); ■ *Syncopacma larseniella* GOZM.




Fig. 12: Map of the Volga-Ural region: ● - *Metanarsia modesta* STGR.; ▲ - *M. incertella* STGR.; ◆ - *Qeoconia quadripuncta* (Hw.); ■ - *Holcopogon bubulcellus* STGR.




Fig. 13: Female genitalia of "Coleophora unicorella TOLL" – *Casignetella unicorella* (ZELLER, 1839): a - female genitalia; b - abdomen.

BALDIZZONE, G. (1988): Contributions à la connaissance des Coleophoridae XLVIII. Guadre nouvelles espèces du genre *Coleophora* HÜBNER de l'URSS. – Beitr. Ent. 38: 74–82.

BALDIZZONE, G. (1994): Coleophoridae dell'Area Irano-Anatolica e regioni limitrofe (Lepidoptera). – Associazione Naturalistica Piemontese. Memorie 3: 423 pp.

- CZEREPANOV, S. K. (1995) Plantae vasculares Rossicae et civitatum collimitanearum (in limicis URSS olim). – S. Petropolis "Mir i Semia-XCV" 901 pp.
- CHRISTOPH, H. (1862): Vier neue südrussische Schmetterlinge. – Stett. Ent. Zeitung **23**: 220–224.
- CHRISTOPH, H. (1872): Neue Lepidoptera des Europäischen Faunengebiets. – Horae Soc. Ent. Ross. **9**: 3–39, pl. 1, 2A.
- CHRISTOPH, H. 1888. Diagnosen zu einigen neuen Lepidopteren des Palaearktischen Faunengebiets. – Horae Soc. Ent. Ross. **22**: 308–314.
- EVERSMANN, E. (1844): Fauna lepidopterologica Volgo-Uralensis. – Casani, 633 pp.
- FALKOVITSH, M. I. (1989): New species of casebearers (Lepidoptera, Coleophoridae) of the Turanian fauna. – Proc. Zool. inst. Acad. Sci. of USSR. **200**: 40–87 (in russian).
- FALKOVITSH, M. I. (1993): New species of casebearer moths (Lepidoptera, Coleophoridae) from Altai. – Proc. Zool. inst. Acad. Sci. of USSR. **251**: 40–52 (in russian).
- GALASJEVA, T. V. (1986): The investigation of woods Zhiguli preserv. – The results of SIW. Moscow. 125 pp (in russian).
- HERRICH-SCHÄFFER, G. A. W. (1856): Neue Schmetterlinge aus Europa und den angrenzenden Landern. I. P. 1–8, pl. II. – Regensburg.
- HERRICH-SCHÄFFER, G. A. W. (1861): Revision der europäischen Schmetterlingsfauna. – Korrespondenzblatt für Sammler von Insekten. Regensburg **2**: 189 pp.
- HODGES, R. W. (1986): The moths of America north of Mexico including Greenland. Fasc. **7.1**. Gelechioidea, Gelechiidae (Part), Dichomeridinae. Washington. P. I–XII + 1–(196), mon. pl. A = HH, col. pl. 1–4.
- KRULIKOVSKIY, L. (1915): The Lepidopterfauna news of the Sergievsk vicinity of Samara Distr. – Russ. ent. oboz. **15**: 218–222 (in russian).
- LERAUT, P. (1980): Liste systematique et synonymique des Lepidopteres de France, Belgique et Corse. – Paris. 334 pp.
- MARTYNOVA, E. F. (1952): Fauna's peculiarities of Cisurals Lepidoptera and its meaning for steppe forestry. – Proc. Zool. inst. Acad. Sci. of USSR **11**: 66–91 (in russian).
- PISKUNOV, V. I. (1990): A new species and two new synonyms of the Palaeartic gelechiid moth (Lepidoptera, Gelechiidae). – News of faunistics and systematics. Izd-vo "Naukova dumka", Kiev, p. 95–97 (in russian).
- POVOLNY, D. (1966): Neubeschreibungen von palaearktischen *Scrobipalpa*-Arten (Lepidoptera, Gelechiidae). – Acta entomol. Bohemoslov. **63** (5): 394–408.
- POVOLNY, D. (1989): Bemerkungen zur generischen und subgenerischen Bewertung einiger Gnorimoschemini (Insecta, Lepidoptera: Gelechiidae). – Staatl. Mus. Tierk. Dresden **26** (2): 157–167.
- REBEL, H. (1901): Catalog der Lepidopteren des palaearktischen Faunengebiets. II. Theil: Pyralidae-Micropterygidae. Catalog der Lepidopteren des palaearktischen Faunengebiets. Berlin, R. Friedlander und Sohn, 368 pp.
- REZNIK, S. YA. (1976): On the fauna of the genus *Multicoloria* CAP. (Lepidoptera, Coleophoridae) of the Far East. – Trudy zool. Inst. Leningr. **62**: 109–113 (in russian).
- REZNIK, S. YA. (1977): A short review of the genus *Multicoloria* CAP. (Lepidoptera, Coleophoridae). – Systematics and faunistics of insects. – Leningrad. Acad. Sci. of USSR. **78–88**, 11 figs (in russian).
- SACHKOV, S. (1995): A review of the fauna of the gelechiid moths (Lepidoptera, Gelechiidae) of Samara District. – Samarskay Luka. **6**: 218–223 (in russian).

- STAUDINGER, O. (1870): Beschreibung neuer Lepidopteren des europäischen Faunengebiets. Berlin. Ent. Zeit. **14**: 97–132, 193–208, 273–330.
- STAUDINGER, O. (1879): Ueber Lepidopteren des südöstlichen europäischen Russland. Stett. Ent. Zeit. **40** (1–3): 315–328.
- STAUDINGER, O. (1880): Lepidopteren-Fauna Kleinasiens. – Horae Soc. ent. ross. **15**: 369–435.
- STAUDINGER, O. & M. WOCKE (1871): Catalog der Lepidopteren des europäischen Faunengebiets. – Dresden. 426 pp.
- TOLL, S. (1953): Rodzina Eupistidae Polski. – Mater. Zoograph. Kraju. **32**: 292 pp, Tabl. I–XXVIII.
- TOLL, S. (1957): Studien über die Genitalien einiger Coleophoriden, XIV, Lepidoptera. – Acta Zool. Krakow. **2** (6): 119–148, 41 figs.
- TOLL, S. (1961): Studien über die Genitalien einiger Coleophoridae, XVIII (Lepidoptera). – Ann. Zool. **19** (6): 209–277, 43 figs.
- TOLL, S. (1962): Materialien zur Kenntnis der paläarktischen Arten der Familie Coleophoridae (Lepidoptera). – Acta Zool. Krakoviensis **7** (16): 577–720, 133 pls.
- ZELLER, H. (1871): Lepidopterologische Beobachtungen im Jahre 1870. – Stett. Ent. Ztg. **32**: 49–81, Pl. 2.

addresses of the authors

V. V. ANIKIN

University, Dept. of Animal Morphology and Ecology
Astrakhanskaja 83
410071 Saratov
Russia

S. A. SACHKOV

University, Dept. Ecology, Botany and Nature Reserve
ul. Acad. Pavlova 1
443011 Samara
Russia

V. V. ZOLOTUHIN

Pedagogical Institute
Dept. of Zoology
pl. 100-letia Lenina 2
432700 Uljanovsk
Russia

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Atalanta](#)

Jahr/Year: 1999

Band/Volume: [29](#)

Autor(en)/Author(s): Anikin Vasily Victorovich, Sachkov Sergej A., Zolotuhin Vadim V.

Artikel/Article: ["Fauna Lepidopterologica Volgo-Uralensis" 150 years later: changes and additions. Part 4. Coleophoridae, Gelechiidae, Symmocidae and Holcopogonidae \(Insecta, Lepidoptera\) 295-336](#)