

**New systematic status for genus *Dodiopsis*
and some notes about *Dodia* distribution**

(Lepidoptera, Arctiidae, Geometridae)

by

POVILAS IVINSKIS & AIDAS SALDAITIS

received 23.IV.2004

Summary: *Dodiopsis* (described as a genus within the family Arctiidae) is synonymized with *Autotrichia* (Geometridae). New data on the distribution of *Dodia sazonovi* (Arctiidae) are presented.

Introduction

In 2001, IVINSKIS and SALDAITIS described a new genus *Dodiopsis* and a new species *D. solanikovi* (IVINSKIS & SALDAITIS, 2001). The description was based on the morphological characteristics of females, because the authors did not have the possibility to study the male genitalia structures of the new species. Lately, they have collected abundant material of the genera *Dodia* and *Autotrichia*, which allowed to determine a new systematic status for the species *Dodiopsis solanikovi*.

Results

In recent years, the authors have obtained 5 male individuals of the same species of the genus *Autotrichia* (Siberia, E. Sayan Tunkin Mts., Mondy v., Hulugaisha Mnt., 2400 m, 15.–20.VI. 2002, S. ОВУКНОВ leg.) as well as 10 males and 2 females of 2 species (Russia, S.W. Tuva West Tanuola R. Sagly Riv. Valley, 05.–15.VI.2003, h-2700 m leg. VASCHENKO). The individuals of the Geometrid genus flew together with representatives of *Epimydia dialampra* and *Dodia sazonovi* (Arctiidae). These butterflies, though belonging to different families, are very similar externally. The butterflies of the above-mentioned genera are distinguished by dark transparent thinly-scaled wings, whereas the antennae of *Autotrichia* and *Epimydia* males are double pectinate. Females of all these butterflies are characterized by more or less reduced wings.

Dodiopsis IVINSKIS & SALDAITIS, 2001 = *Autotrichia* WEHRLI, 1934 **syn. nov.**

The newly collected material suggests that the above-described genus and the species *Dodiopsis solanikovi* ("Arctiidae") do not differ externally from individuals of the genus *Autotrichia* (Geometridae). The genitalia (a large sacklike bursa with a striped signum, the antrum nearly equal in length to a short ductus bursa) (fig. 1). and legs (foretibia: fig. 2; hindtibia: fig. 3) of *Autotrichia* spec. (*solanikovi*) females are typical for moths of this genus. Antennae, venation of wings and legs of the male of *Autotrichia* spec. (*solanikovi*) are identical to other species of the genus *Autotrichia*. The above-specified features allow the authors to conclude that the described genus *Dodiopsis* should be synonymized with the genus *Autotrichia*. The species status of *Autotrichia solanikovi* is questionable and needs further investigation.




Fig. 1: *Autotrichia* spec. female genitalia.
Fig. 2: *Autotrichia* spec. female foretibia.
Fig. 3: *Autotrichia* spec. female hindtibia.

The two *Autotrichia* species live in damp tundra in Tuva at 2500–2700 m elevation. Males were active during daytime; females were found under stones. *Epimydia dialampra* and *Dodia sazonomi* males, active in daytime, were found together with *Autotrichia*. *Dodia sazonomi* females were detected under stones (pers. com. VASCHENKO). This locality (West Tanuola, Tuva) is new to the species *Dodia sazonomi*, which has been reported from the Altai Mts. so far.

Acknowledgements

We would like to thank Mr. VASHENKO, Mr. S. CHURKIN and Mr. H. OSTRASKAS for material and for valuable advice and help. We are also grateful to Mrs. SIGITA DAUGIRDIENE for translating this article into English.

References

- IVINSKIS P. & A. SALDAITIS (2001): A new genus, *Dodiopsis* gen. nov., and a new species, *Dodiopsis solanikovi* spec. nov., from Mongolia (Lepidoptera, Arctiidae). – *Atalanta* **32** (1/2): 227–232, plate VIb.
- IVINSKIS P. & A. SALDAITIS (2001): A correction to IVINSKIS P. & A. SALDAITIS: A new genus, *Dodiopsis* gen. nov., and a new species, *Dodiopsis solanikovi* spec. nov., from Mongolia (Lepidoptera, Arctiidae). *Atalanta* **32** (1/2): 227–232). – *Atalanta* **32** (3/4): 484.
- PROUT, L. B. (1934–1935d, additions 1938a): Brepinae, Oenochrominae, Hemitheinae, Sterrhinae, Larentinae. In: SEITZ, A. (ed.): *Die Gross-Schmetterlinge der Erde, Suppl. 4*. Verlag A. Kernen, Stuttgart: 2–3, plate I.
- VII DALEPP, J. (1996): Checklist of the Geometridae (Lepidoptera) of the former U.S.S.R. – Apollo Books, Stenstrup, 111 pp.

Explanation of colour plate IXa (p. 169):

- Fig. 1: *Dodia diaphana* EVERSMANN, ♂, Mongolia, Arhangaj Aimak, Hangaj Mts., 15 km South from Bulgan vill., 19.–20.VI.2003, h–2040 m, N 47° 16', E 101° 03', leg. A. SALDAITIS.
- Fig. 2: *Dodia albertae* DYAR, ♀, Russia, Khabarovsk district, Myaochan Mts., Gorny vill., Silinka river, 15.VII.1998.
- Fig. 3: *Dodia kononenkoi* TSHISTJAKOV & LAFONTAINE, ♀, Russia, Burjatia, Mondy vill., 10.VI.2001, h–2000 m, leg. KARDASHOV.
- Fig. 4: *Epimydia dialampra* STAUDINGER, ♂, Mongolia, Khovd Aimak, Mongolian Altai Mts., Sutai uul (N.W. slopes), 2700–2900 m, 12.–14.VII.2003, leg. S. CHURKIN.
- Fig. 5: *Epimydia dialampra* STAUDINGER, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h–2700 m, leg. VASCHENKO.
- Fig. 6: *Epimydia dialampra* STAUDINGER, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h–2700 m, leg. VASCHENKO.
- Fig. 7: *Dodia sazonomi* DUBATOLOV, ♀, Russia, Altai Mts., 50° 16'–20' N, 87° 50'–55' E, Kuraisky khrebet, h–3000 m, 29.VI.2000, T. and K. NUPPONEN leg.
- Fig. 8: *Dodia sazonomi* DUBATOLOV, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 24.VI.2003, h–2700 m, leg. VASCHENKO.

Fig. 9: *Dodia sazonomi* DUBATOLOV, ♀, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 01.VII.2003, h-2700 m, leg. VASCHENKO.

Fig. 10: *Dodia sazonomi* DUBATOLOV, ♂, Russia, Altai Mts., 50° 16–20' N, 87° 50–55' E, Kuraisky khrebet, h-3000 m, 02.VII.2000, T. and K. NUPPONEN leg.

Fig. 11: *Epimydia dialampra* STAUDINGER, ♀, Russia, Transbaikalien, Middle Temnik River, h-700 m, 1.–2.VI.1993. M. L. PROKOFIEV leg.

Fig. 12: *Autotrichia* spec., ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h-2700 m, leg. VASCHENKO.

Fig. 13: *Autotrichia* spec., ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h-2700 m, leg. VASCHENKO.

Fig. 14: *Autotrichia* spec., ♂, Sibiria, E. Sayan, Tunkin Mts., Mondy vill., Hulugaisha Mnt., 15.–20.VI.2002, h-2400 m, S. ОВУКHOV leg.

Fig. 15: *Autotrichia* spec., ♀, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h-2700 m, leg. VASCHENKO.

1	2	3	
4	5	6	7
8	9	10	11
12	13	14	15

Addresses of the authors

POVILAS IVINSKIS
Akademijos 2
Vilnius University Institute of Ecology
2600 Vilnius, Lithuania
e-mail: entlab@centras.lt

AIDAS SALDAITIS
Bendoriai 252
Avizieniai, Vilnius reg
e-mail: saldrasa@gmx.net

Colour plate IXa

IVINSKIS, P. & A. SALDAITIS: New systematic status for genus *Dodiopsis* and some notes about *Dodia* distribution (Lepidoptera, Arctiidae, Geometridae). – *Atalanta* 35 (1/2): 105–108.

Fig. 1: *Dodia diaphana* EVERSMAHN, ♂, Mongolia, Arhangaj Aimak, Hangajin Mts., 15 km South from Bulgan vill., 19.–20.VI.2003, h-2040 m, N 47° 16', E 101° 03', leg. A. SALDAITIS. Fig. 2: *Dodia albertae* DYAR, ♀, Russia, Khabarovsk district, Myoachan Mts., Gorny vill., Silinka river, 15.VII.1998. Fig. 3: *Dodia kononenkoi* TSHISTIAKOV & LAFONTAINE, ♀, Russia, Burjatia, Mondy vill., 10.VI.2001, h-2000 m, leg. KARDASHOV. Fig. 4: *Epimydia dialampra* STAUDINGER, ♂, Mongolia, Khovd Aimak, Mongolian Altai Mts., Sutaï uul (N.W. slopes), 2700–2900 m, 12.–14.VII.2003, leg. S. CHURKIN. Fig. 5: *Epimydia dialampra* STAUDINGER, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h-2700 m, leg. VASCHENKO. Fig. 6: *Epimydia dialampra* STAUDINGER, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI.2003, h-2700 m, leg. VASCHENKO. Fig. 7: *Dodia sazonomi* DUBATOLOV, ♀, Russia, Altai Mts., 50° 16'–20' N, 87° 50'–55' E, Kuraisky khrebet, h-3000 m, 29.VI.2000, T. and K. NUPPONEN leg. Fig. 8: *Dodia sazonomi* DUBATOLOV, ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 24.VI.2003, h-2700 m, leg. VASCHENKO. Fig. 9: *Dodia sazonomi* DUBATOLOV, ♀, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 01.VII.2003, h-2700 m, leg. VASCHENKO. Fig. 10: *Dodia sazonomi* DUBATOLOV, ♂, Russia, Altai Mts., 50° 16'–20' N, 87° 50'–55' E, Kuraisky khrebet, h-3000 m, 02.VII.2000, T. and K. NUPPONEN leg. Fig. 11: *Epimydia dialampra* STAUDINGER, ♀, Russia, Burjatia, S.W. Transbaikalien, Middle Temnik River, h-700 m, 1.–2.VI.1993, M. L. PROKOPIEV leg. Fig. 12: *Autotrichia* spec., ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI. 2003, h-2700 m, leg. VASCHENKO. Fig. 13: *Autotrichia* spec., ♂, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI. 2003, h-2700 m, leg. VASCHENKO. Fig. 14: *Autotrichia* spec., ♂, Sibiria, E. Soyan, Tunkin Mts., Mondy vill., Hulugaisha Mt., 15.–20.VI.2002, h-2400 m, S. OBUKHOV leg. Fig. 15: *Autotrichia* spec., ♀, Russia, S.W. Tuva, West Tanuola Mts., Sagly Riv. valley, 05.–15.VI. 2003, h-2700 m, leg. VASCHENKO.

1	2	3	
4	5	6	7
8	9	10	11
12	13	14	15

Farbtafel IXb

Freina, J. J. & T. J. Witt: *Paidia elegantia* spec. nov., eine neue Flechtenbärenart aus dem südlichen Iran (Lepidoptera, Arctiidae, Lithosiinae). – *Atalanta* 35 (1/2): 109–113.

Abb. 1: *Paidia elegantia* spec. nov., Holotypus ♂; Iran, Fars, Straße Ardekan–Talo-chosroo, Corné, 2600 m, 5.VIII.1937, [leg. et] coll. BRANDT, in Museum WITT, München (MWM). Abb. 2: *Paidia elegantia* spec. nov., Paratypus ♂; wie Abb. 1, jedoch 7.VIII.1937 (MWM). Abb. 3: *Paidia elegantia* spec. nov., Paratypus ♂; S. Iran, Miyan Kotal, 1900 m, östl. Kazerun, 51°40'1/29°30'B, 4.–7.VI.1969, leg. VARTIAN (NHMW). Abb. 4: *Paidia elegantia* spec. nov., Paratypus ♂ aberrativ; wie Abb. 3 (MHMW). Abb. 5: *Paidia conjuncta conjuncta* (STAUDINGER, 1891) ♂; W-Iran, Bala-vi-Taq, Berge v. Kasri-Shirin, 1100 m, 3.X.1965, leg. E. & A. VARTIAN (MHMW). Abb. 6: *Paidia conjuncta conjuncta* (STAUDINGER, 1891) ♂; Türkei, Anatolien, 25 km südl. Sivas, 1500 m, 24.+26.VII.1978, leg. W. THOMAS (MWM). Abb. 7: *Paidia conjuncta major* DANIEL, 1963 ♂; Iran, 70 km S v. Teheran, 1300 m, 5.V.1965, [leg.] KASY & VARTIAN (NHMW). Abb. 8: *Paidia conjuncta major* DANIEL, 1963 ♂; Pakistan, Prov. Jammu & Kaschmir, Gilgit valley, 3 km E of Gakuch, 1870 m, 26.VII.1998, leg. CSÓVÁRI & MIKUS (MWM).

1	2	3	4
5	6	7	8
	d	c1	c2

Farbtafel IXc

FREINA, J. J. DE: Erstnachweis von *Amata (Syntomis) nigricornis* (ALPHÉRAKY, 1883) für die iranische Fauna (Lepidoptera, Arctiidae, Syntominiæ). – *Atalanta* 35 (1/2): 114–116.






Amata (Syntomis) nigricornis nigricornis (ALPHÉRAKY, 1883). Abb. 1: ♂, Nordiran, Provinz Azarbaygan-e-Sarqi, N Ahar, Umg. Kaleybar, 1750 m, 14.VIII. 2003, leg. TEN HAGEN, in coll. DE FREINA. Abb. 2: ♂, gleicher Fundort (Maßstab 1:1,05).

Colour plate IXd

STAYKOV, D.: Kurze Übersicht über die Art *Colias erate* (ESPER, [1805]) in Bulgarien mit Beschreibung eines ungewöhnlichen aberranten männlichen Exemplars (Lepidoptera, Pieridae). – *Atalanta* 35 (1/2): 33–36.

Abb. 1: *Colias erate* ab. nov., leg. & coll. STAYKOV.

Colour plate/Farbtafel IXa-d



ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Atalanta](#)

Jahr/Year: 2004

Band/Volume: [35](#)

Autor(en)/Author(s): Ivinskis Povilas, Saldaitis Aidas

Artikel/Article: [New systematic status for genus Dodiopsis and some notes about Dodia distribution 105-108](#)