

A new subspecies of *Neope ramosa* LEECH, 1890 from Yunnan, S.W. China

(Lepidoptera, Nymphalidae, Satyrinae)

by

SONG-YUN LANG

received 29.IX.2023

**Abstract:** In this paper, a new subspecies, *Neope ramosa nanzhao* subspec. nov., is described from southern Yunnan Province, Southwest China.

*Neope ramosa* LEECH, 1890 (= *swinhonis* MELL, 1939; *daguanica* SUGIYAMA, 2015) (Satyrini: Lethina), which had been considered as a variation or a subspecies of *Neope pulaha* (MOORE, [1858]) for a long time (LEECH, 1892; SEITZ, 1907; d'ABRERA, 1990), was known with only its nominate subspecies from E. China (Zhejiang & Fujian) to W. China (W. Sichuan) (OKANO & OKANO, 1984; HUANG, 2002, 2003; LANG, 2017). *Neope ramosa daguanica* SUGIYAMA, 2015 was described from Daguan, N.E. Yunnan, and latter, LANG (2017) sunk it as a junior synonym of the nominate subspecies *ramosa* LEECH basing upon the following reasons: first, no very obvious differences can be found between them; second, no geographical barrier can be found between *daguanica* SUGIYAMA from Daguan County, which is located at southwestern fringe of Sichuan Basin, and the typical *ramosa* LEECH from W. Hubei and C. & E. Sichuan (Provincial borderlines in China had been made based mainly upon political reasons but not barely upon natural geographical lines since the late 13th century AD, namely the Yuan dynasty, so N.E. Yunnan shares similar natural features with mountainous areas of southern Sichuan Basin). SAITO & VU (2020) first discovered *Neope ramosa* LEECH from the Indo-Chinese fauna at Hoang Lien, N. Vietnam, and identified the Vietnamese population as *N. ramosa daguanica* SUGIYAMA. However, specimen illustrated by SAITO & VU (2020: p. 28, figs. 2, 2v) is not *N. ramosa daguanica* SUGIYAMA which is only a junior synonym of the nominate subspecies, but in fact, it should be an undescribed subspecies belonging to the northern Indo-Chinese fauna. In this study, a specimen of *Neope ramosa* LEECH was collected from Lincang, S. Yunnan, S.W. China, and it is similar to the population from N. Vietnam. Considering that S. Yunnan often shares a common fauna with N. Indo-China, the present author here describes a new subspecies of *Neope ramosa* LEECH which is the inhabitant of S. Yunnan and N. Indo-China.

Materials in this study are kept in Chongqing Museum of Natural History, Beibei, CHINA (CMNH) and Dr. SONG-YUN LANG's private collection, Beibei, CHINA (LSY). Photos of the typus of *Neope goschkewitschi swinhonis* MELL kept in Zoologische Forschungsinstitut und Museum Alexander Koenig, Bonn, GERMANY (ZFMK) was provided by Mr. S. Y. HUANG (Bonn).

*Neope ramosa ramosa* LEECH, 1890 (figs: 1-2, 6, 9a)

*Neope ramosa* LEECH, 1890: 29. Type locality (TL): Chang-yang, [Hubei].

*Neope goschkewitschi swinhonis* MELL, 1939: 144. TL: Kuatun, Fukien [Guadun, Fujian].

*Neope ramosa daguanica* SUGIYAMA, 2015: 33, pl. 2: 4, 5, 12, 13. TL: Daguan, N.E. Yunnan.

**Material:** Photo of ♂, "Typus/ Kuatun (2300 m) 27, 40 n. Br. 117, 40 ö. L. J. KLAPPERICH 20. 6. 1938 (Fukien)/*N. goschkewitschi swinhonis* MELL Typus" (ZFMK); 2♂♂, CHINA, Sichuan, Omei, Jiulinggang, 1700 m, 8.VIII.2013, leg. SONG-YUN LANG (LSY); 1♂, CHINA, Sichuan, Nanjiang, Mt. Micangshan, Daba, 1500 m, 12.VI.2016, leg. SONG-YUN LANG (LSY); 1♂, CHINA, Chongqing, Jiangjin, Mt. Simianshan, 1500-2000 m, 29.VI.2008, leg. LI-JIANG WANG (CMNH); 3♂♂, "ginfu [Nanchuan, Mt. Jifoshan] 1400 m 1932 JUL 15 (CMNH)"; 1♂ "ginfu shan 1200 m 1932 JUL 1" (CMNH); 1♂, "ginfo 1000 m JUL11 1932" (CMNH).

**Distribution:** China (S. Henan, S. Shaanxi, Zhejiang, Fujian, Hubei, Hunan, Sichuan, Chongqing, Guizhou, N.E. Yunnan).

*Neope ramosa nanzhao* subspec. nov. (figs: 3, 7, 9b)

*Neope ramosa daguanica*: SAITO & VU (nec SUGIYAMA), 2020: 27, 28, figs. 2, 2v.

**Holotype:** ♂, CHINA, Yunnan, Lincang, Wulao-shan (Lincang Daxue Shan), 2450 m, 16.VII.2023, leg. SONG-YUN LANG, kept in CMNH.

**Diagnosis:** A) The new subspecies can be distinguished from the only known subspecies of *Neope ramosa* LEECH, viz. the nominate subspecies, by the combination of the following characters: 1) it is smaller than the nominate subspecies; 2) on the hindwing underside, the postdiscal ocelli in the spaces 1b to 3 are much closer to the termen than in the nominate subspecies; 3) on the hindwing underside, the darkish submarginal line is located medially between the postdiscal ocelli and the marginal line in the spaces 2 and 3, whereas it is located far away from the ocellus in the nominate subspecies; 4) ♂ uncus (fig. 7) is thinner than in the nominate subspecies (fig. 6); 5) the lamina of the androconium (fig. 9b) is shorter and much narrower than in the nominate subspecies (fig. 9a).

B) The new subspecies is similar to and sympatric with *Neope pulaha pulaha* (MOORE) (figs: 4, 8, 9c-2), but it can be distinguished from *N. pulaha* (MOORE) (figs: 4-5, 8, 9c-d) by the combination of the following characters: 1) it is larger than *N. pulaha* (MOORE); 2) on the hindwing underside, the postdiscal ocelli are obviously larger than in *N. pulaha* (MOORE); 3) on the hindwing underside, the whitish pupils of the postdiscal ocelli are prominent, whereas they are somewhat obscure in *N. pulaha* (MOORE); 4) ♂ uncus (fig. 7) is swelled mediodorsally, whereas in *N. pulaha* (MOORE) (fig. 8) it is straight throughout and only sharply acute near the tip; 5) the lamina of the androconium (fig. 9b) is longer and slightly wider than in *N. pulaha* (MOORE) (figs. 9c-d).

**Etymology:** The subspecific name *nanzhao* is named after Nanzhao (738A.D.-902A.D.), which was a dynastic kingdom including nowadays Yunnan and several of its surrounding areas. Nanzhao was an independent neighbour of the Chinese Tang Empire (618A.D.-907A.D.).

**Distribution:** China (S. Yunnan), N. Vietnam.


**Acknowledgements:** The present author thanks Mr. V. V. TSHIKOLOVETS (Pardubice), Mr. H. SUGIYAMA (Gifu), Mr. S. Y. HUANG (Bonn) and Ms. J. HOU (Beibei) for their various helps.

#### References

- D'ABRERA, B. (1990): Butterflies of the Holarctic Region. I. Papilionidae, Pieridae, Danaidae & Satyridae (Partim). - Hill House Publishers, Melbourne.
- HUANG, H. (2002): Some new satyrids of the tribe Lethini from China (Lepidoptera, Satyridae). - Atalanta **33** (3/4): 361-372, Würzburg.
- HUANG, H. (2003): A list of butterflies collected from Nujiang (Lou Tse Kiang) and Dulongjiang, China with descriptions of new species, new subspecies, and revisional notes (Lepidoptera, Rhopalocera). - Neue Entomologische Nachrichten **55**: 3-114, 160-177, Marktleuthen.
- LANG, S. Y. (2017): The Nymphalidae of China (Lepidoptera, Rhopalocera). Part II. Satyrinae (partim): Tribe Satyrini (partim): Subtribes Eritina, Ragadiina, Lethina, Parargina. - Tshikolovets Publications, Pardubice.
- LEECH, J. H. (1890): New species of Lepidoptera from China. - Entomologist **23**: 26-50, 81-83, 109-114, 187-192, London.
- LEECH, J. H. (1892): Butterflies from China, Japan, and Corea 1, 2. - R. H. Porter, London.
- MELL, R. (1939): Beitraege zur Fauna sinica. xviii. Noch unbeschriebene chinesische Lepidopteren (V). - Deutsche Entomologische Zeitschrift Iris **52**: 135-152, Dresden.
- MONASTYRSKII, A. L. & A. L. DEVYATKIN (2016): Butterflies of Vietnam (an illustrated checklist) 2<sup>nd</sup> edition. - Planorama Media Co., Ltd., Hanoi.
- OKANO, M. & T. OKANO (1984): Notes on some species of *Neope* from China (Lepidoptera: Satyridae). - Tokurana **6/7**: 119-123, Mishima.
- SAITO, T. & V. L. VU (2020): New records of two species of Satyrinae from northern Vietnam. - Butterflies **83**: 27-29, Tokyo.
- SEITZ, A. (1907-1909): The Macrolepidoptera of the world **1**. - Alfred Kernen, Stuttgart.
- SUGIYAMA, H. (2015): Notes on New and Interesting Sino-Himalayan Butterflies. - Pallarge **9**: 29-53, Gifu.

#### Address of the author

SONG-YUN LANG  
Chongqing Museum of Natural History,  
400700, Beibei, China  
Email: langsongyun@gmail.com



*Neope ramosa ramosa*


*N. ramosa nanzhao*



*N. pulaha pulaha*

*N. pulaha emeensis*

**Fig. 1-2:** *Neope ramosa ramosa* LEECH, 1890: (1) ♂, Typus of *Neope goschkevitschi swinhonis* MELL, 1939, Fukien, Kuatun, ZFMK; (2) ♂, Sichuan, Nanjiang, SATY0442, ANDR0077, LSY. **Fig. 3:** *Neope ramosa nanzhao* subspec. nov., holotype, ♂, Yunnan, Lincang, SATY1206, ANDR0315, CMNH. **Fig. 4:** *Neope pulaha pulaha* (MOORE, [1858]), ♂, Yunnan, Nanjian, SATY1207, ANDR0316, LSY. **Fig. 5:** *Neope pulaha emeensis* Li, 1995, ♂, Sichuan, Nanjiang, SATY0441, ANDR0076, LSY.



**Fig. 6-8:** ♂ genitalia. g: ♂ genitalia in lateral view with left valva and aedeagus removed; t: tegumen+uncus+gnathos in lateral view; al: aedeagus in lateral view; ad: aedeagus in dorsal view; v: tip of left valva in dorsal view. **Fig. 6:** *Neope ramosa ramosa* LEECH, 1890, Sichuan, Nanjiang, SATY0442, ANDR0077, LSY. **Fig. 7:** *Neope ramosa nanzhao* subsp. nov., holotype, Yunnan, Lincang, SATY1206, ANDR0315, CMNH. **Fig. 8:** *Neope pulaha pulaha* (MOORE, [1858]), Yunnan, Nanjian, SATY1207, ANDR0316, LSY.

**Fig. 9:** Androconia. **a:** *Neope ramosa ramosa* LEECH, 1890, Sichuan, Nanjiang, SATY0442, ANDR0077, LSY. **b:** *Neope ramosa nanzhao* subsp. nov., holotype, Yunnan, Lincang, SATY1206, ANDR0315, CMNH. **c1-2:** *Neope pulaha pulaha* (MOORE, [1858]): (c1) Tibet, Medog, SATY0491, ANDR0078, LSY; (c2) Yunnan, Nanjian, SATY1207, ANDR0316, LSY. **d:** *Neope pulaha emeensis* Li, 1995, Sichuan, Nanjiang, SATY0441, ANDR0076, LSY.

**Fig. 10:** Distribution map of *Neope ramosa* LEECH, 1890 (Sources of data: LEECH, 1890; MELL, 1939; SUGIYAMA, 2015; LANG, 2017; SAITO & VU, 2020; specimens kept in LSY, CMNH).

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Atalanta](#)

Jahr/Year: 2023

Band/Volume: [54\\_3-4](#)

Autor(en)/Author(s): Lang Song-Yun

Artikel/Article: [A new subspecies of Neope ramosa Leech, 1890 from Yunnan, S.W. China \(Lepidoptera, Nymphalidae, Satyrinae\) 369-372](#)